

BNL-209351-2018-TECH EIC/59

An ergodic approach to polarization in eRHIC electron storage ring

F. Meot

October 2018<br>
Collider Accelerator Department

# **Brookhaven National Laboratory**

# **U.S. Department of Energy**

USDOE Office of Science (SC), Nuclear Physics (NP) (SC-26)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-SC0012704 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof*.* 

# An ergodic approach to polarization in eRHIC electron storage ring

F. Méot Collider-Accelerator Department, BNL, Upton, NY 11973

October 26, 2018

## Abstract

Electron polarization in a storage ring is subject to two long term effects: Sokolov-Ternov polarization and depolarization by diffusion, which over a long time scale lead to an equilibrium state. Simulation-wise, this can be highly CPU time and memory consuming. Simulations aimed at determining optimal ring storage energy in an electron-ion collider use to track thousand particle bunches, for a long time - yet still short compared to depolarization time scales, due to HPC limitations. Based on considerations of ergodicity of electron bunch dynamics in the presence of synchrotron radiation, tracking a single particle instead is investigated. This allows substantial saving in teh required HPC volume, "CPU-time×Memory-allocation". The concept is illustrated with polarization lifetime and equilibrium polarization simulations at the eRHIC electron-ion collider.

> eRHIC Note <sup>59</sup> BNL C-AD 2018

# **Contents**



#### 1 INTRODUCTION 3

## 1 Introduction

The eRHIC installation is briefly described in Fig. 1 [1]. The 18 GeV eRHIC electron storage lattice used in the present spin polarization



Figure 1: The eRHIC electron-ion collider complex, an 18 GeV-255 Gev/nucleon electron-ion collider installation.

simulations has been provided by S. Tepikian [1], optical parameter values relevant to the present simulations will be introduced in due place. The eRHIC lattice includes a double non-planar rotator system (Fig. 2) at the interaction point (IP), comprised of strong solenoids and series of bends, which allows to locally move the stable spin precession direction  $\vec{n}_0$ , from vertical in the arcs to longitudinal at the IP. In a defect-free



Figure 2: Half of the spin rotator system at eRHIC (the system is symmetric with respect to the IP, at the origin here). In green: solenoids.

ring, this region of off-vertical  $\vec{n}_0$  is a major contribution to spin diffusion.

Bunches are injected in the storage ring with alternately up and down polarization, and replaced every 6 min in order to ensure an average polarization of 70% over the hundreds of bunches stored [1]. A proper lattice should maintain bunch depolarization below 20% (absolute) over the 6 min storage. The evolution of the polarization, from  $P_0 = \pm 0.85$  at injection to  $P_{eq}$  at equilibrium (an asymptotic quantity to be determined), satisfies

$$
P(t) = P_{eq}(1 - e^{-t/\tau_{eq}}) + P_0 e^{-t/\tau_{eq}} \tag{1}
$$

This results from (i) synchrotron-radiation (SR) self-polarization and (ii) polarization loss by diffusion, with time constant  $\tau_D$ , such that

$$
1/\tau_{\text{eq}} = 1/\tau_{\text{SP}} + 1/\tau_{\text{D}} \tag{2}
$$

Sokolov-Ternov (ST) self polarization in a flat ring has a time constant  $\tau_{ST[sec.]} \approx 99 \rho_{\rm [m]}^2 {\rm R}_{\rm [m]}/{\rm E}_{\rm [GeV]}^5$  [2], about 30 min at eRHIC at 18 GeV, 10 hrs at 10 GeV, with asymptotic value  $P_{ST} = 92.4\%$ ; the asymptotic self-polarization is taken instead  $P_{SP} = 90\%$  here to account for the non-planar spin rotator, and with time constant  $\tau_{\text{SP}}$ , such that [2]

$$
P_{eq} = P_{SP} \times \tau_{eq} / \tau_{SP}
$$
 (3)

The goal in tracking spin motion is (i) to validate a ring design, including preservation of polarization under the effect of defects, corrections, etc. and (ii) to determine an optimal working point a $\gamma_{ref}$  (a = 1.15965 10<sup>-3</sup> is the electron anomalous magnetic moment).

#### 2 STOCHASTIC MOTION 4

In the following, a method based on single-particle tracking is discussed. First, basic aspects of the stochasticity of particle and spin motions are recalled. Then tracking outcomes are displayed and the single-particle method is discussed.

The numerical simulations discussed in this paper have strongly benefited from NERSC means and environment [3].

## 2 Stochastic motion

The dynamical system of a high energy stored electron bunch at equilibrium is ergodic: over a long time interval, trajectories in the system cover all parts of the 6D phase space. Time averages over one or more trajectories are equivalent to phase space averages,

$$
\lim_{T \to \infty} \int_{t_0}^{t_0 + T} f(\vec{X}(t)) dt = \int f(\vec{X}) \rho(\vec{X}) d^N \vec{X} \Big|_{time = t}
$$
\n(4)

For all three motions, transverse and longitudinal, the evolution of the bunch emittance with time, t, satisfies

$$
\bar{\epsilon}_{n}(t) = \epsilon_{n,eq} \left( 1 - e^{-t/\tau_{n}} \right) + \epsilon_{n,i} e^{-t/\tau_{n}}
$$
\n(5)

(n stands for x, y, or 1) with  $\epsilon_{n,i}$  and  $\epsilon_{n,eq}$  respectively the starting and equilibrium emittances,  $\tau_n = \frac{T_{rev} E_s}{T}$  $\frac{U_{\rm rev} L_{\rm s}}{U_{\rm s}}$  the damping time constant.  $J_{n=x,y,l}$  are the partition numbers,  $J_x + J_y + J_l = 4$ ,  $J_l \approx 2$ . Eq. 5 indicates that after a few damping times, the bunch dynamical system can be considered at equilibrium, bunch emittances have reached their asymptotic values. In the following  $\tau_{SR} = \tau_x \approx \tau_y$  denotes the transverse SR damping time constant.

At 18 GeV the energy loss amounts to 38.7 MeV/turn (a result from prior tracking of a 2000 particle bunch with Monte Carlo SR), thus the damping time amounts to  $\tau_{\rm SR} = 18$ <sub>GeV</sub>/38.7<sub>MeV/turn</sub> = 465 turns, 6 ms.

Figs. 3, 4 display the stochastic motion of a single particle over  $10^3 \tau_{SR}$  and by comparison the instantaneous horizontal and vertical phase spaces of a 10<sup>3</sup>-electron bunch observed at time =  $10^3 \tau_{\rm SR}$ . In this example, statistical variable values such as *rms* coordinates, emittances, either single particle projected over a long tracking time, or multiparticle at time  $t=10^3\tau_{\rm SR}$ , resulting from both methods, essentially satisfy Eq. 4. Over a sufficiently long time interval, an electron has explored the all 6D phase-space, which is a necessary condition for ergodicity to be satisfied.

Out of equilibrium,  $\vec{X}(t) - \vec{X}(t)$  can be taken as the statistical vaiable, with  $\vec{X}(t)$  the average value.  $\vec{X}(t)$  can be determined from a fit using the theoretical damping, for instance in the single particle case, see below. Spin motion is not at equilibrium, the polarization decays with time, fast in resonant conditions. Both the decay time constant and the asymptotic polarization are zero on the resonance, as  $\tau_{\rm eq}\sim \delta^2\times\tau_{\rm SP},\;{\rm P_{eq}}\sim \delta\times {\rm P_{SP}},$  with  $\delta={\rm a}\gamma_{\rm Res.}-{\rm a}\gamma$  the distance to the resonance [2, p. 125]. Fig. 5 displays typical stochastic spin motion in eRHIC storage ring at 18 GeV. In a similar way that  $\tau_{\rm SR}$  can be obtained from the observation of the damped motion of a single electron far from equilibrium,  $\tau_D$  can be obtained from long enough observation of spin motion.

## 3 Polarization

In order to assess polarization properties of a storage ring depending on its energy setting, spin tracking simulations are performed over an ensemble of a $\gamma_{\rm ref}$  rings covering some a $\Delta\gamma_{\rm ref}$  < 1 range of interest. In these hypothesis, all these rings have the same optics: bend strengths  $1/\rho$ , focusing strengths  $G/B\rho$ , chromaticities, etc., are unchanged, what changes is the energy  $a\gamma_{\text{ref}}$  at which each ring is run.

Particles are all launched with their initial spin direction parallel to the local nominal stable spin precession direction  $\vec{n}_0$  (*i.e.*, longitudinal if the origin is taken at IP6, vertical at IP8). Spin tracking only starts after 10 damping times about (5,000 turns) when the bunch has reached its equilibrium emittance. If the motion happens to neighbor a depolarizing resonance, spin will tilt away from the nominal direction toward possibly large angles depending on the strength of/distance to the resonance (in the absence of SR and at constant energy, the spin would steadily rotate around the local tilted  $\vec{n}_{\delta}$ ). Away from any resonance, the spin is expected to only be subject to slow diffusion.

The single particle "depolarization landscape" is expected to look as shown in Fig. 6, obtained in HERA-e conditions, which include a spin rotator which introduces strong depolarization in the  $a\gamma$  = integer regions. Single particle tracking here yields Fig. 7 (18 GeV) and Fig. 8 (10 GeV), which appears qualitatively similar to DESY simulation outcomes, Fig. 6. This is the outcome of the tracking, over a time interval  $[0, t]$ , of a single particle in each one of 1024 (or 2048) rings, all operated with the same optics but with each its particular operation energy  $a\gamma_{ref}$ . These distributions feature similar topology, at both timings. Zooming in on any reduced  $a\Delta\gamma_{ref}$  interval also shows a similar spin distribution (sort of "fractal"). The energy excursion over that time interval  $[0, t]$  is displayed in Fig. 9. Note a property that will be referred to later: the equilibrium energy spread is  $\sigma_E \approx 10^{-3}E$  at 18 GeV, or an extent  $\sigma_{a\gamma} \approx 0.04$ , thus the beam covers ≈40 (80) bins of a 1024 (2048) bin a $\Delta\gamma_{\rm ref} = 1$  interval.

Spin diffusion has to be a slow process for a lattice to be viable, in particular this cannot be the case if, during its energy excursion, a particle neighbors (is within a few resonance strengths from) a depolarizing spin resonance ( $\nu_s \pm \nu_x \pm \nu_y \pm \nu_z =$  integer). In such case, the depolarization is not slow (the orientation of the spin vector changes substantially during the tracking: the vertical spin component  $S_y$ moves towards  $S_y = -1$  in the present representation, Fig. 5). Thus, the working point of concern,  $a\gamma_{ref}$  near resonance, is not optimal. A contrario, observing only slow change in the spin vector means absence of harmful resonance in the energy interval that the particle spans due to SR, and potentially a viable working point.



Figure 3: Left: transverse particle excursion (red: x; blue: y), over time interval  $t/\tau_{SR}: 1 \to 10^3$  (450,000 turns about). Right: transverse phase spaces, matching ellipses and histograms; blue: projection of the single particle motion of the left plot; red: for comparison, case of a  $10^3$  particle bunch, observed at time  $t = 10^3 \tau_{SR}$ . Note that the  $\epsilon_y/\epsilon_x$  ratio represents a 27% coupling, of which the source is the spin rotator in IR6 which includes solenoids (Fig. 2).



Figure 4: Left: stochastic energy excursion over time interval  $t/\tau_{SR}$  :  $1 \to 10^3$ .  $\sigma_{\delta p/p} = 1.14\,10^{-3}$ . Right: longitudinal phase space; blue: projection of the multiturn single particle motion of the left plot; red: case of a 10<sup>3</sup> particle bunch observed at t =  $10^3 \tau_{\rm SR}$ ;  $\sigma_{\delta p/p} = 1.13 \, 10^{-3}$ .



Figure 5: Left: stochastic spin motion observed at IP8 ( $\vec{n}_0$  vertical there), single particle, a few different cases of ring rigidity settings in the 18 GeV region. Right: monitor individual spins, a linear regression on  $P/P_0 = \exp(-t/\tau_D) \approx 1 - t/\tau_D$  provides the diffusion time constant  $\tau_D$ .

#### Diffusion time constant

Spin tracking here does not include the self-polarization process, it is assumed that  $\tau_{SP}$  in Eq. 2 is obtained from the lattice. Thus, a bunch keeps depolarizing, due to diffusion, polarization tends toward  $P_{eq} = 0$  ( $\tau_{SP} = \infty$ ) with a time constant  $\tau_D = \tau_{eq}$ .

In order to ensure the required polarization survival (70% on average over the ensemble of bunches in the ring, stored 6 min each)  $\tau_D$  has to be sufficiently long compared to the store duration; this eliminates, for a viable rigidity setting of the ring, the regions  $a\gamma_{\text{ref}} < 40.45$  and  $> 40.75$  in Fig. 7 (18 GeV), a $\gamma_{\text{ref}} < 22$  and  $> 22.4$  in Fig. 8 (10 GeV). Finally, with  $\tau_{\text{D}}$  much larger than the time interval covered by the tracking (of the order of seconds at best, whereas  $\tau_D$  has to amount to tens of minutes for a lattice to be viable), one can use

$$
P(t)/P_0 = \exp(-\frac{t}{\tau_D}) \approx 1 - t/\tau_D \tag{6}
$$

Single particle spin tracking data are displayed in Fig. 5, a fit of these data provides  $\tau_D$ . From that  $\tau_{eq}$  can be derived (Eq. 2) yielding in turn  $P_{eq}$  (Eq. 3) and  $P(t = 6 \text{ min})$  (Eq. 1).

In order to assess the method, in the following for simplicity, and  $P_{eq}$  being a sub-product, primary tracking outcomes are considered, namely, spin orientation or  $\tau_D$  landscapes.

## 4 A metric

Typically, the energy dependence of particle spins over a  $a\Delta\gamma_{\rm ref}$  interval looks as shown in Figs. 7, 8. A different criterion to quantify the depolarization could be instead, Fig. 9, the energy dependence of the minimal value that spins reached in the course of the tracking. This minimum may happen earlier in the tracking, as observed in Fig. 5, as spins oscillate around a given tilted local  $\vec{n}_{\delta}$ , as long as the latter does not change due to photon emission.

In order to allow comparisons between lattices a metric is required. However, Figs. 7, 8 styles of data do not lend themselves to straightforward comparisons, essentially due to the stochastic aspect. A couple of different possibilities are assessed here instead, based on sliding averaging.

### 4.1 Spin

A sliding average of the data of Fig. 7, with a small sliding interval  $\alpha\delta\gamma_{\rm ref}$  (a few particles / bins), namely

$$
\langle S_{\mathbf{y}} \rangle_{\mathbf{N}} (a \gamma_{\text{ref}, \frac{\mathbf{N}}{2}}) = \frac{1}{\mathbf{N}} \sum_{i}^{i+\mathbf{N}-1} S_{\mathbf{y}} (a \gamma_{\text{ref}, i}) \tag{7}
$$

greatly smooths the fluctuations, as observed in Fig. 10. The local excursion of S<sub>y</sub> over a small  $\delta\gamma_{\rm ref}$  interval in the optimal region (P<sub>f</sub>/P<sub>i</sub>  $\approx$ 0.9983) are grossly below  $\pm 210^{-4}$ . This corresponds to a fluctuation of  $\tau_D$  of less than  $\pm 7$  min around an average  $\approx 60$  min.,  $\approx \pm 10\%$ 



Figure 6: Left: asymptotic polarization at HERA, using SITROS [4]. This graph was produced by tracking bunches of a few hundred polarized electrons, including Monte Carlo SR, over a few SR damping times, for a series of evenly spaced ring rigidity values over an a $\Delta \gamma = 1$  interval  $(37 \leq a\gamma \leq 38)$ . Each a $\gamma$  value represents a particular operation rigidity of the ring, however with unchanged optics. Right, obtained from the present eRHIC simulations: turn-by-turn average spin value over the complete bunch population yields the self-polarization time constant, from what the asymptotic polarization, similarly to the representation in the left plot, can be drawn (average over just a few particles is shown here).



Figure 7: Spin rotation landscape (a photo of the spins,  $S_1(a\gamma_{ref})|_{time=t}$ ) at IP8 where  $\vec{n}_0$  is vertical), at either (red)  $t = 80 \times \tau_{SR}$  (10<sup>4</sup> turns) or (blue)  $t = 900 \times \tau_{SR}$  (4.5 10<sup>5</sup> turns or 5 s). Left: the 2048 rings tracked cover over a a $\Delta\gamma = 1.2$  interval, encompassing integer a $\gamma_{\rm ref}$  values where full spin flip is observed, as expected. Right: a zoom in on a reduced  $40.5 < a\Delta\gamma_{\rm ref} < 40.75$  interval; no strong resonances in that region instead, and spins remain close to  $S_y = 1^-$ .

relative, a good first approach. This can be further improved by increasing the number of bins, for instance, once determined that the region  $40.6 < a\gamma_{\text{ref}} < 40.7$  is viable, the computation can be reiterated.

Why allow a sliding average:

(i) with the present extent a $\Delta\gamma_{\rm ref} \approx 1$  covered in 1024 bins (or 2048), the energy extent of the bunch  $\sigma_{\gamma_{\rm ref}}/\gamma_{\rm ref} \approx 10^{-3}$  or a $\delta\gamma_{\rm ref} = 0.04$ , covers about 40 (or 80) bins,

(ii) thus, a set of a few neighboring bins almost belong in the same ring, averaging over a few bins is not so different from averaging over a few particles in the same bin,

(iii) in any case a possible strong, nearby resonance would cause a dip in the distribution, indicating a non viable  $a\delta\gamma_{\rm ref} = 1$  region, which has to be avoided.

The strong smoothing effect of a sliding average suggest to apply it directly to the final spin distribution, this is done in Fig. 10, case of a  $\Delta\gamma$  : 40.4  $\rightarrow$  40.9 interval covered in 1024 bins. In the present conditions, this series converges when increasing the width of the sliding interval  $[a\delta\gamma_{\rm ref,i}, a\delta\gamma_{\rm ref,i+N}]$ . At some point however, increasing the sliding interval would cause it to reach  $a\gamma_{\rm ref}$  regions where the fluctuations change in a sensible manner (as in Fig. 7-left), for instance featuring a different average, or including high amplitude spikes, so abruptly changing the sum of the series, however the eRHIC lattices of interest have to satisfy  $\langle S_{v} \rangle (a\gamma_{ref}) \approx 1$ , which prohibits such changes. In the present case of 1024 ring samples over  $\Delta \gamma$  : 40.4  $\rightarrow$  40.9, N=40 appears appropriate; this is a sliding window of full width  $a\Delta\gamma_{ref} = 40 \times (40.9 - 40.4)/1024 \approx 0.02$ . Fig. 10 confirms that, with the  $40.59 - 40.63$  interval yielding a final  $\tau_D \approx 55\%$  within about  $\pm 5\%$  of convergence values for both 40.60 – 40.62 and 40.56 – 40.66 sliding windows. It may also be thought of increasing the bins density in the  $a\gamma_{ref}$  of interest to obtain a better homogeneous distribution.



Figure 8: Spin rotation landscape (a photo of the spins,  $S_1(a\gamma_{ref})|_{time=t}$ ) at IP6 where  $\vec{n}_0$  is longitudinal), at 160 SR damping times ( $\approx 5 \times 10^5$  turns). The right and left plots differ by the a $\Delta\gamma$  interval width. Integer a $\gamma_{\rm ref}$  regions show full spin flip as expected. In the reduced 22.33 < a $\Delta\gamma_{\rm ref}$  < 22.47 region no strong resonance is observed, spins remain close to  $S_y = 1^-$ . The right vertical scale in both plots is the rms width of the energy interval explored by a particle during the tracking.



Figure 9: Left, blue curve: depolarization landscape  $S_{y,min}(a\gamma_{ref})|_{t\in[0,T]}$ , over  $39.8 < a\gamma_{ref} <$ 40.9 (18 GeV region), observed at IP8 (nominal  $\vec{n}_0$  is vertical there) at time T =  $10^3 \tau_{\rm SR}$ . This is the smallest value of the projection of particle spin  $\tilde{S}$  on the vertical axis, reached over the time interval [0, T]. The right vertical scale (green curve) is the rms width of the energy interval explored by a particle during the tracking. Right: a zoom in on a reduced  $40.5 < a\Delta\gamma_{ref} < 40.8$  interval, showing  $S_{y,\min}(a\gamma_{ref})|_{t\in[0,T]}$  at (red)  $T = 20 \times \tau_{SR}$  (9500 turns) and (blue)  $T = 1000 \times \tau_{SR}$ (450,000 turns or 6 s ).



Figure 10: Derived from Fig. 7, by applying a sliding average (Eq. 7). The four curves differ by the sampling: average over N= either 10, 50, 90 or 130 a $\delta\gamma_{\rm ref}$  samples, centered at given  $a\gamma_{ref}$  In the present conditions, the distribution converges when increasing the number of samples, N, i.e., the width of the sliding interval,  $[a\delta\gamma_{\rm ref,i}, a\delta\gamma_{\rm ref,i+N}]$ .

#### 5 SUMMARY 9

#### 4.2 Diffusion time constant

The diffusion time constant is derived from single particle motion using a linear regression (Eq. 6), an illustration can be found in Fig. 5-right. Applying to each of the 2048 rings over the a $\Delta\gamma_{\rm ref}$  interval, yields the scan in Fig. 11. Tracking over several SR damping times is needed to draw  $\tau_D$  from individual spin motion, It can be seen in Fig. 11 that, in the region of great  $\tau_D$  values of interest the statistics over 80, 160 and  $10^3 \times \tau_{SR}$  superimpose. This indicates that the required tracking time is comparable with that needed to determine  $\tau_{SR}$  from particle motion, which is a few tens of damping times or less. On the other hand, in case the strong fluctuations of the spin would cause too strong a dependence of the  $\tau_{SR}$  value (from the fit) on the fit sample, rather than increasing the damping time a possibility is instead to launch a few particles per ring: the smoothing effect is immediate, this can be seen by comparison of the spin motion in Figs. 5-right and the averaging over a few particles in Fig. 6-right. These considerations matter as to the interest of the single particle method, this is discussed in the next Section.





Figure 11: Energy scan of the diffusion time constant in the single particle tracking method. 2048 bins cover  $a\Delta\gamma$  : 39.8 – 41.2.  $\tau_D$ values interpolated from 80 (red), 160 (green) and  $10^3 \times \tau_{\rm SR}$  tracking are plotted.

Figure 12: Sliding over energy scan of the diffusion time constant in the single particle tracking method. 2048 bins.

Again a sliding average, applied to the data of Fig. 11, greatly smooths the fluctuations, as observed in Fig. 12. The distance between  $\tau_D$ distributions can be derived from Fig. 12 type of data, which are thus usable for comparing polarization performances of different versions of the eRHIC e-storage ring.

## 5 Summary

Assume similar resolution using both methods, "HPC-Hungry" and "Ergodic", namely, the same number of reference rings, nRings, over the same interval a $\Delta\gamma$ .

In the present hypotheses (eRHIC lattice, energy, etc.):

- first method: the HPC volume is nRings  $\times 10^3$  [particles/bunch]  $\times$  a few  $\tau_{\rm SR}$ ,

- second method: the HPC volume is nRings  $\times$  a few  $\tau_{\text{SR}}$ .

This is a 3 orders of magnitude difference in the HPC volume. On the other hand, greater HPC volume translates in one or the other of, longer queues, longer computing time, more processors, greater volume of I/Os, larger data analysis HPC volume.

It remains to determine how close the single particle method can get to the accuracy of the bunch method (an ongoing work). However it already appears an efficient first approach to the diffusion time constant, in view of qualifying an evolution of a lattice design, the efficiency of error correction and other spin matching schemes. Because it is faster it allows a greater reach (for instance in terms of parameter space exploration) in machine simulations and design optimizations.

## 6 Conclusion

Obviously these results are very preliminary, they are essentially indications that the HPC volume could be reduced. More simulations are required, for further inspection, comparisons between the two methods, etc. Mathematical background and support is in order.

The simulations discussed here were performed on NERSC [3], using stepwise ray-tracing tools for spin motion accuracy [5]. Electron dynamics and spin diffusion in the presence of Monte Carlo SR is a long installed and, needless to say, thoroughly benchmarked feature of the code [6].

# Acknowledgements

This study has benefited from fruitful discussions with D. Abell (Radiasoft), A. Cerfon (CIMS-NYU) and B. Nash (Radiasoft).

# References

- [1] eRHIC p-CDR, BNL (2018)
- [2] S.Y. Lee, Spin dynamics and snakes in synchrotrons, World Scientific (1997).
- [3] NERSC computing, on web : http://www.nersc.gov/
- [4] J. Kewisch, Depolarization der Elektronspins in Speicherringen Durch Nichtlineaire Spin-Bahn-Copplung, DESY 85-109, Okt. 1985.
- [5] https://zgoubi.sourceforge.io/ZGOUBI DOCS/Zgoubi.pdf; https://www.osti.gov/scitech/biblio/1062013-zgoubi-users-guide
- [6] Simulation of radiation damping in rings, using stepwise ray-tracing methods, F. Méot, JINST 10 T06006 (2015).