

3D Vlasov theory of the plasma cascade instability

M. Blaskiewicz

October 2019

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC), Nuclear Physics (NP) (SC-26)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-SC0012704 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

3D Vlasov Theory of the Plasma Cascade Instability

M. Blaskiewicz

BNL 911B, Upton, NY 11973, USA

The plasma cascade instability (PCI) [1, 2] is a proposed mechanism for microbunching in electron beams without dipole magnets. Existing theory is limited to wave propagation that is orthogonal to the advective compression direction. This note provides a theory allowing for wave propagation in arbitrary directions.

The plasma cascade instability (PCI) [1, 2] is a proposed mechanism for microbunching in electron beams without dipole magnets. If the theory bears out this process may well be very widespread, contributing to enhanced noise in a variety of systems employing electron beams. This note shows how one can reduce the Vlasov equation to a set of Volterra equations of the second kind, which are amenable to accurate numerical solution.

Consider a homogeneous, infinite, electron plasma. We use Cartesian coordinates x_1, x_2, x_3, t . The unperturbed plasma has an average velocity

$$\mathbf{v}_0(\mathbf{x}, t) = \sum_{j=1}^3 \hat{x}_j x_j \omega_j(t) \quad (1)$$

This average velocity is driven by the acceleration

$$\mathbf{A}_0(\mathbf{x}, t) = \sum_{j=1}^3 \hat{x}_j x_j k_j(t) \quad (2)$$

We wish to find a distribution with these parameters. We will take an unperturbed distribution of the form $f_0(\mathbf{x}, \mathbf{v}, t) = f_0(H(\mathbf{x}, \mathbf{v}, t))$ with

$$H = \sum_{j=1}^3 \frac{\alpha_j(t)}{2} (v_j - \omega_j x_j)^2. \quad (3)$$

The Vlasov equation is

$$\frac{\partial f_0}{\partial t} + \mathbf{v} \cdot \frac{\partial f_0}{\partial \mathbf{x}} + \mathbf{A}_0(\mathbf{x}, t) \cdot \frac{\partial f_0}{\partial \mathbf{v}} = 0, \quad (4)$$

and since $f_0 = f_0(H)$, H satisfies equation (4) as well. Insert H for f_0 in equation (4). The resulting terms proportional to x_j and v_j are

$$-\alpha_j(v_j - \omega_j x_j) \dot{\omega}_j x_j + \dot{\alpha}_j(v_j - \omega_j x_j)^2/2 - v_j \alpha_j \omega_j (v_j - \omega_j x_j) + x_j k_j \alpha_j (v_j - \omega_j x_j) = 0. \quad (5)$$

Setting the coefficients of x_j^2 , $x_j v_j$ and v_j^2 to zero we find equation (4) is satisfied if

$$\dot{\alpha}_j = 2\alpha_j \omega_j, \quad \dot{\omega}_j + \omega_j^2 = k_j \quad (6)$$

for $j = 1, 2, 3$. If equations (6) are satisfied then any function $f_0(H)$ will satisfy equation (4). For physical solutions we require $f_0(H) d^3 x d^3 v$ to be the number of electrons in the phase space volume $d^3 x d^3 v$.

We will use first order perturbation theory with $f = f_0 + f_1$ so that

$$\frac{\partial f_1}{\partial t} + \mathbf{v} \cdot \frac{\partial f_1}{\partial \mathbf{x}} + \mathbf{A}_0(\mathbf{x}, t) \cdot \frac{\partial f_1}{\partial \mathbf{v}} + \mathbf{A}_1(\mathbf{x}, t) \cdot \frac{\partial f_0}{\partial \mathbf{v}} = 0, \quad (7)$$

where \mathbf{A}_1 is the acceleration created by f_1 and by the ion seeding the instability. Using the results of [3] we introduce time dependent spatial wave numbers and assume a perturbation where the spatial density of the electrons varies as

$$n_{\mathbf{P}}(\mathbf{x}, t) = \hat{n}_{\mathbf{P}}(t) \exp \left(i \sum_{j=1}^3 P_j \lambda_j(t) x_j \right) \equiv \hat{n}_{\mathbf{P}}(t) \exp(i\Psi(\mathbf{x}, t)) \quad (8)$$

where $\dot{\lambda}_j + \omega_j \lambda_j = 0$ and the P_j, s are constant in time. The total perturbed density from all wavenumbers is

$$n_1(\mathbf{x}, t) = \int n_{\mathbf{P}}(\mathbf{x}, t) d^3 P.$$

Consider an ion implanted at $t = 0$ and located at $\mathbf{x} = \mathbf{x}_0 + \mathbf{v}_0 t$. The acceleration it generates is given by (cgs units)

$$\begin{aligned} \mathbf{A}_I(\mathbf{x}, t) &= \frac{qQ}{m} \frac{\mathbf{x} - \mathbf{x}_0 - \mathbf{v}_0 t}{|\mathbf{x} - \mathbf{x}_0 - \mathbf{v}_0 t|^3} \\ &= -i \frac{4\pi qQ}{m(2\pi)^3} \int d^3 P \lambda_1 \lambda_2 \lambda_3 \frac{(P_1 \lambda_1, P_2 \lambda_2, P_3 \lambda_3)}{\sum_j P_j^2 \lambda_j^2} \exp \left(i \sum_m P_m \lambda_m (x_m - x_{0m} - v_{0m} t) \right), \end{aligned} \quad (9)$$

where $q = -|e|$ is the electron charge, m is its mass, and Q is the charge on the ion. The net acceleration for wavenumber \mathbf{P} due to both the ions and electrons is $\mathbf{A}_{\mathbf{P}}(x, t) = \tilde{\mathbf{A}}(t) \exp(i\Psi(\mathbf{x}, t))$ with [5]

$$\tilde{\mathbf{A}}(t) = -4\pi i \frac{(P_1 \lambda_1, P_2 \lambda_2, P_3 \lambda_3)}{\sum_m P_m^2 \lambda_m^2} \left\{ \frac{q^2}{m} \hat{n}_{\mathbf{P}}(t) + \frac{qQ}{(2\pi)^3 m} \lambda_1 \lambda_2 \lambda_3 \exp \left(-i \sum_m P_m \lambda_m (x_{0m} + v_{0m} t) \right) \right\}. \quad (10)$$

To solve the Vlasov equation we consider a single \mathbf{P} . Consider the Ansatz

$$f_{\mathbf{P}}(\mathbf{x}, \mathbf{v}, t) = \frac{df_0}{dH} g(v_1 - \omega_1 x_1, v_2 - \omega_2 x_2, v_3 - \omega_3 x_3, t) \exp(i\Psi(\mathbf{x}, t)).$$

Notice that the x_j dependence in g and f_0 only shows up as $v_j - \omega_j x_j$ so it drops out after integrating over v_j . This generates the correct spatial dependence for $n_{\mathbf{P}}(\mathbf{x}, t)$. For convenient notation define $u_j = v_j - \omega_j x_j$ and remember that

$$\frac{\partial g(\mathbf{x}, \mathbf{v}, t)}{\partial t} = \frac{\partial g(\mathbf{u}, t)}{\partial t} + \frac{\partial g(\mathbf{u}, t)}{\partial \mathbf{u}} \cdot \frac{\partial \mathbf{u}(\mathbf{x}, \mathbf{v}, t)}{\partial t}.$$

Plugging into the Vlasov eq one finds

$$\frac{\partial g(\mathbf{u}, t)}{\partial t} + \sum_{j=1}^3 i g P_j \lambda_j u_j - \omega_j u_j \frac{\partial g}{\partial u_j} + \alpha_j u_j \tilde{A}_j = 0 \quad (11)$$

To proceed we multiply the last term on the right of equation (11) by $\delta(t - t_0)$ with the intention of integrating over t_0 later.

$$\frac{\partial \tilde{g}(\mathbf{u}, t, t_0)}{\partial t} + \sum_{j=1}^3 i \tilde{g} P_j \lambda_j u_j - \omega_j u_j \frac{\partial \tilde{g}}{\partial u_j} + \alpha_j u_j \tilde{A}_j \delta(t - t_0) = 0 \quad (12)$$

We look for solutions of the form

$$\tilde{g}(\mathbf{u}, t, t_0) = H(t - t_0) \mathbf{q}(t) \cdot \mathbf{u} \exp(i\mathbf{K}(t) \cdot \mathbf{u})$$

where $H(t - t_0)$ is 1 for $t \geq t_0$ and zero otherwise. Inserting this expression into (11) and using the same sort of tricks used to solve (5) we find that (12) is satisfied if

$$\dot{K}_j + P_j \lambda_j(t) - \omega_j(t) K_j = 0, \quad \text{with } K_j(t_0) = 0 \quad (13)$$

$$\dot{q}_j - \omega_j(t) q_j = 0, \quad \text{with } q_j(t_0) = -\alpha_j(t_0) \tilde{A}_j(t_0). \quad (14)$$

To bring the pieces together define the general solution $M_j(t)$ as the solution to equation (13) but with the boundary condition $M_j(0) = 0$. Also define the phases $\Phi_j(t)$ so that $\omega_j(t) = \dot{\Phi}_j(t)$. With these definitions $K_m(t, t_0) = M_m(t) - M_m(t_0) \exp[\Phi_m(t) - \Phi_m(t_0)]$ and $q_j(t, t_0) = -\alpha_j(t_0) \tilde{A}_j(t_0) \exp[\Phi_j(t) - \Phi_j(t_0)]$. This yields

$$g(\mathbf{u}, t) = \sum_{j=1}^3 \int_0^t dt_0 q_j(t, t_0) u_j \exp \left(i \sum_{m=1}^3 u_m K_m(t, t_0) \right) \quad (15)$$

To close the equations we note that the only unknown in $\tilde{A}_j(t)$ is

$$\hat{n}_{\mathbf{P}}(t) = \int d^3u \frac{\partial f_0}{\partial H}(\mathbf{u}, t) g(\mathbf{u}, t) \quad (16)$$

$$= - \sum_{j=1}^3 \int_0^t dt_0 \alpha_j(t_0) \tilde{A}_j(t_0) \exp(\Phi_j(t) - \Phi_j(t_0)) \int d^3u f'_0 \left(\sum_{m=1}^3 \alpha_m(t) u_m^2 / 2 \right) u_j \exp \left(i \sum_{k=1}^3 u_k K_k(t, t_0) \right), \quad (17)$$

$$= - \sum_{j=1}^3 \int_0^t dt_0 \tilde{A}_j(t_0) G_j(t, t_0), \quad (18)$$

$$= \int_0^t dt_0 \left[\frac{q^2}{m} \hat{n}_{\mathbf{P}}(t_0) + D_I(t_0) \right] \sum_{j=1}^3 \left(\frac{\frac{4\pi i P_j \lambda_j(t_0)}{3}}{\sum_{m=1}^3 \lambda_m^2(t_0) P_m^2} G_j(t, t_0) \right) \quad (19)$$

where

$$G_j(t, t_0) = \alpha_j(t_0) \exp(\Phi_j(t) - \Phi_j(t_0)) \int d^3u f'_0 \left(\sum_{m=1}^3 \alpha_m(t) u_m^2 / 2 \right) u_j \exp \left(i \sum_{k=1}^3 u_k K_k(t, t_0) \right), \quad (20)$$

$$= - \frac{i \alpha_j(t_0) K_j(t, t_0)}{\alpha_j(t)} \exp(\Phi_j(t) - \Phi_j(t_0)) \int d^3u f_0 \left(\sum_{m=1}^3 \alpha_m(t) u_m^2 / 2 \right) \exp \left(i \sum_{k=1}^3 u_k K_k(t, t_0) \right). \quad (21)$$

and

$$D_I(t) = \frac{qQ}{(2\pi)^3 m} \lambda_1(t) \lambda_2(t) \lambda_3(t) \exp \left(-i \sum_m P_m \lambda_m(t) (x_{0m} + v_{0m} t) \right). \quad (22)$$

We now have a Volterra equation of the second kind for $\hat{n}_{\mathbf{P}}$. While exact solutions look hopeless a numerical solution should be straightforward. We note that changing the integration variable in equation (21) to $z_i = \sqrt{\alpha_i} u_i$ turns it into the Fourier transform of a spherically symmetric function for which a wide range of exact solutions are available.

CONNECTION TO PREVIOUS WORK

If we set $\omega_i = 0$ then these results should reduce to those in [4]. To show this we set

$$d_i(t) = \frac{Z}{(2\pi)^3} \lambda_1(t) \lambda_2(t) \lambda_3(t) \exp \left(-i \sum_m P_m \lambda_m(t) (x_{0m} + v_{0m} t) \right),$$

$$g(\mathbf{K}(t, t_0)) = \frac{1}{n_0} \int d^3u f_0 \left(\sum_{m=1}^3 \alpha_m(t) u_m^2 / 2 \right) \exp \left(i \sum_{k=1}^3 u_k K_k(t, t_0) \right),$$

$$R(t, t_0) = \frac{1}{\sum_{m=1}^3 \lambda_m^2(t_0) P_m^2} \sum_{j=1}^3 \frac{P_j \lambda_j(t_0) \alpha_j(t_0) K_j(t, t_0)}{\alpha_j(t)} \exp(\Phi_j(t) - \Phi_j(t_0)),$$

where Z is the atomic number of the ion. Now we have

$$\hat{n}_{\mathbf{P}}(t) = \frac{4\pi q^2 n_0}{m} \int_0^t dt_0 [\hat{n}_{\mathbf{P}}(t_0) - d_i(t_0)] R(t, t_0) g(\mathbf{K}(t, t_0)). \quad (23)$$

When $\omega_i = 0$, $R(t, t_0) = t_0 - t$, $\lambda_i = 1$, and $\mathbf{K}(t, t_0) = (t_0 - t)\mathbf{P}$. Make these substitutions, account for a difference in Fourier transform conventions, and include the fact that there is a constant velocity offset between reference frames. One finds that equation (23) here is equivalent to equation (8) in [4].

- [1] V. N. Litvinenko, G. Wang, D. Kayran, Y. Jing, J. Ma, I. Pinayev, arxiv:1802.08677, February 2018
- [2] V. N. Litvinenko, G. Wang, Y. Jing, D. Kayran, J. Ma, I. Petrushina, I. Pinayev, K. Shih, arxiv:1902.10846, February 2019
- [3] M. Blaskiewicz, BNL-211960-2019-TECH, (2019)
- [4] Gang Wang and Michael Blaskiewicz, *Phys. Rev. E* **78**, 026413 (2008).
- [5] a factor of $(2\pi)^3$ is missing in [3]