

BNL-223622-2022-TECH C-A/AP/669

Off-Axis Magnetic Fields Extrapolated from On-Axis Multipoles

S. Brooks

July 2013

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC), Nuclear Physics (NP) (SC-26)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Off-Axis Magnetic Fields Extrapolated from On-Axis Multipoles

Stephen Brooks

July 29, 2013

1 Potential Term

In cylindrical polar coordinates (r, θ, z) , consider the magnetic scalar potential

$$\phi = \sin(n\theta + \psi)r^k f(z)$$

for some integers n, k, angle ψ and function f. The associated magnetic field is

$$\mathbf{B} = \nabla\phi = \frac{\partial\phi}{\partial r}\mathbf{e}_r + \frac{1}{r}\frac{\partial\phi}{\partial\theta}\mathbf{e}_\theta + \frac{\partial\phi}{\partial z}\mathbf{e}_z = \begin{bmatrix} \sin(n\theta + \psi)kr^{k-1}f(z)\\ n\cos(n\theta + \psi)r^{k-1}f(z)\\ \sin(n\theta + \psi)r^kf'(z) \end{bmatrix}_{r\theta z}$$

which automatically satisfies $\nabla \times \mathbf{B} = \nabla \times \nabla \phi = \mathbf{0}$. The only remaining condition on ϕ is

$$0 = \nabla \cdot \mathbf{B} = \nabla \cdot \nabla \phi = \frac{\partial^2 \phi}{\partial r^2} + \frac{1}{r} \frac{\partial \phi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \theta^2} + \frac{\partial^2 \phi}{\partial z^2}$$
$$= \sin(n\theta + \psi) \left(k(k-1)r^{k-2}f(z) + kr^{k-2}f(z) - n^2r^{k-2}f(z) + r^k f''(z) \right)$$
$$= \sin(n\theta + \psi)r^{k-2} \left((k^2 - n^2)f(z) + r^2 f''(z) \right).$$

1.1 Long Multipole

If there is no z behaviour (f = 1, say), then for this to hold for all r, θ requires $k^2 = n^2$. Since $k \leq 0$ cases have a singularity at the origin, put $k = n \geq 1$. This gives the field for an infinitely long multipole:

$$\phi = \sin(n\theta + \psi)r^n, \qquad \mathbf{B} = \begin{bmatrix} n\sin(n\theta + \psi)r^{n-1} \\ n\cos(n\theta + \psi)r^{n-1} \\ 0 \end{bmatrix}_{r\theta z} = nr^{n-1} \begin{bmatrix} \sin((n-1)\theta + \psi) \\ \cos((n-1)\theta + \psi) \\ 0 \end{bmatrix}.$$

n = 1 corresponds to a dipole, n = 2 to a quadrupole, etc. Also $\psi = 0$ gives these in their normal orientation and $\psi = \frac{\pi}{2}$ in their skew orientation.

1.2 Normalisation

Although the rest of the paper will evaluate fields for the potential containing r^n above, this produces fields with magnitude $|\mathbf{B}| = nr^{n-1}$. If multipole strengths are defined as the values of polynomial coefficients of the field function, then $|\mathbf{B}| = k_n r^n$ may be obtained from the potential $\phi = \sin(n\theta + \psi) \frac{k_n}{n+1} r^{n+1}$. If strengths are defined as repeated derivatives of the field function, then $|\mathbf{d}^n \mathbf{B}/dx^n| = d_n$ (i.e. $|\mathbf{B}| = \frac{d_n}{n!} r^n$) may be obtained from $\phi = \sin(n\theta + \psi) \frac{d_n}{(n+1)!} r^{n+1}$.

2 Series Solution

If f'' is nonzero, the $\nabla \cdot \nabla \phi = 0$ equation cannot be satisfied by a single term for all r. However, consider a sum of such terms with the same n but different k:

$$\phi = \sin(n\theta + \psi) \sum_{k=n}^{\infty} r^k f_k(z).$$

Note the lowest term with k = n, which dominates the sum near the r = 0 axis, corresponds to a long 2*n*-pole modulated by $f_n(z)$. The potential must satisfy

$$0 = \nabla \cdot \nabla \phi = \sin(n\theta + \psi) \sum_{k=n}^{\infty} r^{k-2} \left((k^2 - n^2) f_k(z) + r^2 f_k''(z) \right)$$

for all θ , so the RHS sum must be zero (for all r, z). Equating coefficients of r^k gives

$$((k+2)^2 - n^2)f_{k+2}(z) + f_k''(z) = 0$$

for $k \ge n$ and the remaining coefficients of r^{n-2} and r^{n-1} are 0 and $((n+1)^2 - n^2)f_{n+1}(z)$ respectively, so $f_{n+1}(z) = 0$. The above relation gives f_{k+2} as a scaled second derivative of f_k , so $f_{n+2j+1}(z) = 0$ for all $j \ge 0$ and

$$f_{n+2j}(z) = \left(\prod_{i=1}^{j} \frac{-1}{(n+2i)^2 - n^2}\right) f_n^{(2j)}(z).$$

The coefficients will be written as

$$C_{nj} = \prod_{i=1}^{j} \frac{-1}{(n+2i)^2 - n^2} = \prod_{i=1}^{j} \frac{-1}{4(n+i)i} = \frac{(-\frac{1}{4})^j n!}{(n+j)!j!}.$$

Setting $f_n = f$, the full potential satisfying Maxwell's equations in free space is:

$$\phi = \sin(n\theta + \psi) \sum_{j=0}^{\infty} r^{n+2j} f_{n+2j}(z) = \sin(n\theta + \psi) \sum_{j=0}^{\infty} C_{nj} r^{n+2j} f^{(2j)}(z)$$

2.1 Magnetic Field

Using the formula for the gradient of a single term $(\sin(n\theta + \psi)r^k f(z))$ given at the start of this note, the magnetic field associated with the series solution is

$$\begin{split} \mathbf{B} &= \nabla \phi = \sum_{j=0}^{\infty} C_{nj} \begin{bmatrix} \sin(n\theta + \psi)(n+2j)r^{n+2j-1}f^{(2j)}(z) \\ n\cos(n\theta + \psi)r^{n+2j-1}f^{(2j)}(z) \\ \sin(n\theta + \psi)r^{n+2j}f^{(2j+1)}(z) \end{bmatrix}_{r\theta z} \\ &= \sum_{j=0}^{\infty} C_{nj}r^{n+2j-1} \begin{bmatrix} (n+2j)\sin(n\theta + \psi)f^{(2j)}(z) \\ n\cos(n\theta + \psi)f^{(2j)}(z) \\ r\sin(n\theta + \psi)f^{(2j+1)}(z) \end{bmatrix}_{r\theta z} \\ &= \sum_{j=0}^{\infty} C_{nj}r^{n+2j-1} \left(\begin{bmatrix} n\sin(n\theta + \psi)f^{(2j)}(z) \\ n\cos(n\theta + \psi)f^{(2j)}(z) \\ n\cos(n\theta + \psi)f^{(2j)}(z) \\ r\sin(n\theta + \psi)f^{(2j+1)}(z) \end{bmatrix}_{r\theta z} + \begin{bmatrix} 2j\sin(n\theta + \psi)f^{(2j)}(z) \\ 0 \end{bmatrix}_{r\theta z} \right) \end{split}$$

$$=\sum_{j=0}^{\infty} C_{nj} r^{n+2j-1} \left(\begin{bmatrix} n \sin((n-1)\theta + \psi) f^{(2j)}(z) \\ n \cos((n-1)\theta + \psi) f^{(2j)}(z) \\ r \sin(n\theta + \psi) f^{(2j+1)}(z) \end{bmatrix} + \begin{bmatrix} 2j \sin(n\theta + \psi) f^{(2j)}(z) \cos \theta \\ 2j \sin(n\theta + \psi) f^{(2j)}(z) \sin \theta \\ 0 \end{bmatrix} \right)$$
$$=\sum_{j=0}^{\infty} C_{nj} r^{n+2j-1} \begin{bmatrix} (n \sin((n-1)\theta + \psi) + 2j \sin(n\theta + \psi) \cos \theta) f^{(2j)}(z) \\ (n \cos((n-1)\theta + \psi) + 2j \sin(n\theta + \psi) \sin \theta) f^{(2j)}(z) \\ r \sin(n\theta + \psi) f^{(2j+1)}(z) \end{bmatrix}.$$

2.2 Computation Without Trigonometric Formulae

The product formulae for sin and cos give

$$\sin(n\theta + \psi)\cos\theta = \frac{1}{2}\sin((n-1)\theta + \psi) + \frac{1}{2}\sin((n+1)\theta + \psi)$$
$$\sin(n\theta + \psi)\sin\theta = \frac{1}{2}\cos((n-1)\theta + \psi) - \frac{1}{2}\cos((n+1)\theta + \psi),$$

which enables the Cartesian field formula to be written as

$$\mathbf{B} = \sum_{j=0}^{\infty} C_{nj} r^{n+2j-1} \begin{bmatrix} ((n+j)\sin((n-1)\theta+\psi) + j\sin((n+1)\theta+\psi))f^{(2j)}(z) \\ ((n+j)\cos((n-1)\theta+\psi) - j\cos((n+1)\theta+\psi))f^{(2j)}(z) \\ r\sin(n\theta+\psi)f^{(2j+1)}(z) \end{bmatrix}.$$

Note that every instance of $\cos \operatorname{or} \sin(n\theta + \psi)$ is multiplied by a large power of r usually including r^n . This can be used to convert fully to Cartesian coordinates using the complex formulae

$$re^{i\theta} = r(\cos\theta + i\sin\theta) = x + iy$$

$$\Rightarrow \quad (re^{i\theta})^n = r^n e^{in\theta} = r^n(\cos n\theta + i\sin n\theta) = (x + iy)^n$$

whence setting $c_n = \operatorname{Re}(x + iy)^n$ and $s_n = \operatorname{Im}(x + iy)^n$ gives

$$r^n \cos n\theta = c_n$$
 and $r^n \sin n\theta = s_n$

These values can be calculated without trigonometry from the recurrence relation

 $c_0 = 1;$ $s_0 = 0;$ $c_{n+1} = xc_n - ys_n;$ $s_{n+1} = yc_n + xs_n,$

which comes from the definition of complex multiplication $(x + iy)^{n+1} = (x + iy)(x + iy)^n$ in terms of components. The formula requires a slightly more general form

$$\hat{c}_n = r^n \cos(n\theta + \psi) = \operatorname{Re} r^n e^{\mathrm{i}n\theta} e^{\mathrm{i}\psi}, \qquad \hat{s}_n = r^n \sin(n\theta + \psi) = \operatorname{Im} r^n e^{\mathrm{i}n\theta} e^{\mathrm{i}\psi}.$$

These satisfy a similar recurrence because the repeated multiplication by $re^{i\theta} = x + iy$ starts with $e^{i\psi}$ rather than 1:

$$\hat{c}_0 = \cos\psi;$$
 $\hat{s}_0 = \sin\psi;$ $\hat{c}_{n+1} = x\hat{c}_n - y\hat{s}_n;$ $\hat{s}_{n+1} = y\hat{c}_n + x\hat{s}_n$

The evaluations of $\cos \psi$ and $\sin \psi$ are not a problem because they can be precalculated for each multipole, unlike \hat{c}_n and \hat{s}_n that depend on x, y. The field formula can now be rewritten as

$$\begin{split} \mathbf{B} &= \sum_{j=0}^{\infty} C_{nj} r^{2j} \begin{bmatrix} ((n+j)\hat{s}_{n-1} + j\frac{1}{r^2}\hat{s}_{n+1})f^{(2j)}(z) \\ ((n+j)\hat{c}_{n-1} - j\frac{1}{r^2}\hat{c}_{n+1})f^{(2j)}(z) \\ \hat{s}_n f^{(2j+1)}(z) \end{bmatrix} \\ &= \sum_{j=0}^{\infty} C_{nj} \begin{bmatrix} ((n+j)(r^2)^j\hat{s}_{n-1} + j(r^2)^{j-1}\hat{s}_{n+1})f^{(2j)}(z) \\ ((n+j)(r^2)^j\hat{c}_{n-1} - j(r^2)^{j-1}\hat{c}_{n+1})f^{(2j)}(z) \\ (r^2)^j\hat{s}_n f^{(2j+1)}(z) \end{bmatrix}, \end{split}$$

where powers of $r^2 = x^2 + y^2$ have been emphasised as they can be calculated without a square root. Note that negative powers never arise from $(r^2)^{j-1}$ when j = 0 because it is multiplied by j, so disappears.

2.3 Computation Using a 2D Field Map

The only parts of the above formula that depend on θ are the various \hat{c}_n and \hat{s}_n , which are relatively quick (and unavoidable) to calculate for each evaluation point. Calculating the repeated derivatives of f and waiting for the infinite sum to converge are, on the other hand, quite slow but may be precalculated as three functions of just r and z in the following way:

$$\mathbf{B} = \begin{bmatrix} \hat{s}_{n-1}F(r,z) + \hat{s}_{n+1}G(r,z) \\ \hat{c}_{n-1}F(r,z) - \hat{c}_{n+1}G(r,z) \\ \hat{s}_nH(r,z) \end{bmatrix},$$

where

$$F(r,z) = \sum_{j=0}^{\infty} C_{nj}(n+j)(r^2)^j f^{(2j)}(z); \qquad G(r,z) = \sum_{j=1}^{\infty} C_{nj}j(r^2)^{j-1}f^{(2j)}(z);$$
$$H(r,z) = \sum_{j=0}^{\infty} C_{nj}(r^2)^j f^{(2j+1)}(z).$$

2.3.1 Universal Field Map

Considering just one end of the magnet, if the fringe field is related to a universal fringe field function via $f(z) = \hat{f}(z/l)$, then $f^{(n)}(z) = l^{-n} \hat{f}^{(n)}(z/l)$ and

$$F(r,z) = \sum_{j=0}^{\infty} C_{nj}(n+j) \left(\frac{r}{l}\right)^{2j} \hat{f}^{(2j)}\left(\frac{z}{l}\right) = \hat{F}_n\left(\frac{r}{l},\frac{z}{l}\right);$$
$$G(r,z) = \frac{1}{r^2} \sum_{j=1}^{\infty} C_{nj} j \left(\frac{r}{l}\right)^{2j} \hat{f}^{(2j)}\left(\frac{z}{l}\right) = \frac{1}{r^2} \hat{G}_n\left(\frac{r}{l},\frac{z}{l}\right);$$
$$H(r,z) = \frac{1}{r} \sum_{j=0}^{\infty} C_{nj} \left(\frac{r}{l}\right)^{2j+1} \hat{f}^{(2j+1)}\left(\frac{z}{l}\right) = \frac{1}{r} \hat{H}_n\left(\frac{r}{l},\frac{z}{l}\right),$$

where the new functions \hat{F}_n , \hat{G}_n , \hat{H}_n depend only on n and the form of \hat{f} , not l. In terms of these new functions,

$$\mathbf{B} = \begin{bmatrix} \hat{s}_{n-1}\hat{F}_n(r/l,z/l) + \hat{s}_{n+1}\frac{1}{r^2}\hat{G}_n(r/l,z/l) \\ \hat{c}_{n-1}\hat{F}_n(r/l,z/l) - \hat{c}_{n+1}\frac{1}{r^2}\hat{G}_n(r/l,z/l) \\ \hat{s}_n\frac{1}{r}\hat{H}_n(r/l,z/l) \end{bmatrix}.$$

Unfortunately use of the universal function \hat{H}_n incurs a square root to calculate $\frac{1}{r}$, which the non-universal forms do not have, assuming they are stored as functions of r^2 and z.

2.3.2 Field Map for Magnet Ends Only

In the interior of a long magnet, the field tends towards the expression given earlier:

$$\mathbf{B} = nr^{n-1} \begin{bmatrix} \sin((n-1)\theta + \psi) \\ \cos((n-1)\theta + \psi) \\ 0 \end{bmatrix} = \begin{bmatrix} n\hat{s}_{n-1} \\ n\hat{c}_{n-1} \\ 0 \end{bmatrix}.$$

This corresponds to F(r, z) = n and G(r, z) = H(r, z) = 0. The lowest-order terms up to r^2 in these sums are:

$$F(r,z) = C_{n0}nf(z) + C_{n1}(n+1)r^2f''(z) + \dots = nf(z) - \frac{1}{4}r^2f''(z) + \dots;$$

$$G(r,z) = C_{n1}f''(z) + C_{n2}2r^2f''''(z) + \dots = \frac{-f''(z)}{4(n+1)} + \frac{r^2f'''(z)}{32(n+1)(n+2)} + \dots;$$

$$H(r,z) = C_{n0}f'(z) + C_{n1}r^2f'''(z) + \dots = f'(z) - \frac{r^2f'''(z)}{4(n+1)} + \dots,$$

so when f(z) = 1 and f'(z) = f''(z) = 0 to some precision, the field calculation may be replaced by the simple multipole expression. Similarly, when f = f' = f'' = 0 to a good approximation, the field may be given as zero. This means F, G, H only need to be calculated for the transition region where 0 < f(z) < 1 for each magnet (or end of magnet).

2.3.3 Magnets with Symmetrical Ends

It is common to use the same fringe field for both ends of a magnet of length L, so that f(z) = g(z) + g(L-z), where g is a sigmoid function going from $-\frac{1}{2}$ to $\frac{1}{2}$, representing one end of the magnet (e.g. $g(z) = \frac{1}{2} \tanh(z/l)$). If F, G, H are now calculated using g in place of f, the magnetic field is given by

$$\mathbf{B} = \begin{bmatrix} \hat{s}_{n-1}(F(r,z) + F(r,L-z)) + \hat{s}_{n+1}(G(r,z) + G(r,L-z)) \\ \hat{c}_{n-1}(F(r,z) + F(r,L-z)) - \hat{c}_{n+1}(G(r,z) + G(r,L-z)) \\ \hat{s}_n(H(r,z) - H(r,L-z)) \end{bmatrix},$$

noting the sign change for H(r, L-z) because it contains odd derivatives of g.

For the regions where g(z) is almost constant at $-\frac{1}{2}$ or $\frac{1}{2}$, $F(r, z) = -\frac{1}{2}n$ or $\frac{1}{2}n$ may be used respectively and G(r, z) = H(r, z) = 0. If both g(z) and g(L - z) are in a constant region and they have opposite sign (i.e. far outside the magnet), the **B** field will be zero and the calculation should be aborted before calculating \hat{s}_{n-1} etc.