

Transition Jump Using ? ? Produced by the Polarized Proton Fast Quadrupoles

L. Ahrens

July 1986

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

AGS Studies Report

Date(s) July 11, 1986 Time(s) 0800-1600
 Experimenter(s) L.A. Ahrens, E. Raka, L.G. Ratner, S.P. Yamin
 Reported by L.G. Ratner
 Subject Transition Jump Using Δv Produced by the Polarized
Proton Fast Quadrupoles

Observations and Conclusion

This method was successfully used at CERN many years ago at a level of a few 10^{12} protons. They produced a $(\Delta v_H)_{TR} \approx 0.2$ with about a 1-2 msec rise time. Since $\Delta v_H \approx \Delta \gamma$, this gave them a $(\Delta \gamma)_{TR}$ of 200/sec, an enhancement factor of 7, and allowed a clean jump at about $2-3 \times 10^{12}$.

The polarized proton fast quads are capable of producing a $\Delta v_H \approx 0.2$ in 2 μsec or a $\Delta \gamma \approx 10^5/\text{sec}$, an enhancement of 2000. However, it is not only rate that is necessary, but one must not destroy the beam by this non-adiabatic pulse.

In order to produce a $\Delta v_H \approx 0.2$, since the quads are located at vertical β -max positions, one has to apply a $\Delta v_V \approx 0.4$. This would drive the tune to cross $v_V = 9$ and so to prevent this, one must also energize the slow quads to compensate for this.

As we set up conditions for making these tests, it appeared that 2 μsec was too fast. We therefore turned off the HV 2 μsec pulse and used only the LV 50 μsec pulse from the fast quad modulators. This rise time factor will be reinvestigated in our next study. We used the vertical slow quads to compensate for the fast quad pulse and measured that our $(\Delta v_H)_{TR} = -0.17 \pm 0.01$ with about a 50 μsec rise time. This gives an enhancement of 68 which would be good for 2×10^{13} protons, but one must not only have crossing speed but also we think significantly higher $\Delta \gamma$. We were operating the AGS at this time with 2×10^{12} circulating protons and we were able to cross transition very smoothly with this technique. Our attempt to increase the intensity

was masked by other problems and finally lack of time. It is not clear what the intensity limit of this method is and we will plan to continue this study during the next AGS operating period. The following pictures on Page 5 of the γ_{TR} Studies Book show the effect at transition produced by the fast tune shift. A much smoother passage is produced using the fast quads.

Measured with 9 Mods

	Base	Slow Q	Fast Q
Q _V	8.780	- .20 = 8.580	+ .30 = 8.88
$\times \frac{10}{9}$		- .22 = 8.560	+ .33 = <u>8.89</u>
Q _H	8.713	+ .10 = 8.813	- .16 = 8.66
		+ .11 = 8.823	- .18 = <u>8.64</u>
		$\pm .01$	$\pm .01$



$$(\Delta V_H)_{\text{fast}} = -.17 \pm .01 \approx (\Delta \delta)$$

$$\Delta t = 50 \times 10^{-6} \text{ sec}$$

$$\frac{\Delta V}{\Delta t} = \frac{.17 \times 10^6}{50} = 3.4 \times 10^3$$

$$\left(\frac{\Delta \delta}{\Delta t}\right)_0 = \frac{30}{.6 \text{ sec}} = 50/\text{sec}$$

$$\rightarrow f = 68$$



FAST Q OFF. TIMING OF
 ϕ JUMP OPTIMIZED. NO
 RADIAL SHIFT.

Q₂ 10¹²

FAST Q ON. TIMING OPTIMIZ
 NO RADIAL SHIFT
 ϕ JUMP @ 15650
 FQ @ 15800