

BNL-224586-2023-TECH C-A/AP/696

Integrals of Polynomial Functions over Spheres and Balls

S. Brooks

June 2023

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC), Nuclear Physics (NP) (SC-26)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-SC0012704 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Integrals of Polynomial Functions over Spheres and Balls

Stephen Brooks

June 27, 2023

1 Definitions and Problem

Let $B_n = {\mathbf{x} : |\mathbf{x}| \le 1}$ be the unit *n*-dimensional ball and $S_{n-1} = {\mathbf{x} : |\mathbf{x}| = 1}$ be the unit (n-1)-dimensional sphere. Note that this sphere is the surface of the ball: $S_{n-1} = \partial B_n$.

We want to evaluate the integral

$$I_{a_1...a_n}^{B_n} = \int_{B_n} \left(\prod_{i=1}^n x_i^{a_i}\right) \,\mathrm{d}^n \mathbf{x}$$

of a general monomial term over the n-dimensional ball. It will be helpful to define a similar integral

$$I_{a_1...a_n}^{S_{n-1}} = \int_{S_{n-1}} \left(\prod_{i=1}^n x_i^{a_i}\right) \,\mathrm{d}^{n-1}\mathbf{x}$$

over the surface of the sphere. Finally, an integral over all space but weighted by an n-dimensional unit Gaussian distribution will also be useful:

$$I_{a_1...a_n}^{g_n} = \int_{\mathbb{R}^n} \left(\prod_{i=1}^n x_i^{a_i} \right) \frac{1}{\sqrt{2\pi^n}} e^{-\frac{1}{2}|\mathbf{x}|^2} \, \mathrm{d}^n \mathbf{x}.$$

2 Separability of Gaussian Integral

The Gaussian integrand can be written as a product

$$I_{a_1...a_n}^{g_n} = \int_{\mathbb{R}^n} \left(\prod_{i=1}^n x_i^{a_i} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_i^2} \right) \, \mathrm{d}^n \mathbf{x}.$$

Since each term in the product only depends on x_i , this is a separable integral that is the product of one-dimensional integrals:

$$I_{a_1...a_n}^{g_n} = \prod_{i=1}^n \int_{\mathbb{R}} x_i^{a_i} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x_i^2} \, \mathrm{d}x_i = \prod_{i=1}^n I_{a_i}^{g_1}.$$

3 Values of One-Dimensional Gaussian Integral $I_a^{g_1}$

The one-dimensional Gaussian integrals can be evaluated by noting

$$\frac{\mathrm{d}}{\mathrm{d}x} \left[x^a \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \right] = (ax^{a-1} - x^{a+1}) \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2}$$

and therefore by integrating both sides over \mathbb{R} ,

$$0 = a I_{a-1}^{g_1} - I_{a+1}^{g_1}$$

The recurrence of values can be started by noting $I_0^{g_1} = 1$ because the Gaussian is normalised and $I_1^{g_1} = 0$ because it is the integral of an odd function. Writing the recurrence as $I_{a+2}^{g_1} = (a+1)I_a^{g_1}$ makes it clear that $I_a^{g_1} = 0$ for a odd. For even values,

$$I_2^{g_1} = 1, \quad I_4^{g_1} = 3, \quad I_6^{g_1} = 3 \times 5, \quad I_8^{g_1} = 3 \times 5 \times 7, \quad \cdots,$$

giving the general formula

$$I_{2a}^{g_1} = \prod_{b=1}^{a} (2b-1) = (2a-1)!! = \frac{(2a)!}{2^a a!}.$$

3.1 Integrals on the Half Real Line $I_a^{h_1}$

The following calculations will also need 1D Gaussian integrals on the half real line defined by

$$I_a^{h_1} = \int_0^\infty x^a \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \,\mathrm{d}x.$$

Starting as before but integrating over $[0,\infty)$ gives

$$\left[x^a \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2}\right]_{x=0}^{\infty} = aI_{a-1}^{h_1} - I_{a+1}^{h_1}$$

The left hand side is equal to $\frac{-1}{\sqrt{2\pi}}$ when a = 0 and zero otherwise. The recurrence can still be rewritten $I_{a+2}^{h_1} = (a+1)I_a^{h_1}$ valid for $a \ge 0$. The original a = 0 case gives

$$\frac{-1}{\sqrt{2\pi}} = -I_1^{h_1} \qquad \Rightarrow \qquad I_1^{h_1} = \frac{1}{\sqrt{2\pi}}.$$

For even functions we have $I_{2a}^{h_1} = \frac{1}{2}I_{2a}^{g_1}$ and in particular $I_0^{h_1} = \frac{1}{2}$. The recurrence gives the general formulae

$$I_{2a}^{h_1} = \frac{1}{2}(2a-1)!!$$
 and $I_{2a+1}^{h_1} = \frac{1}{\sqrt{2\pi}}\prod_{b=1}^a 2b = \frac{1}{\sqrt{2\pi}}(2a)!! = \frac{1}{\sqrt{2\pi}}2^aa!.$

4 Relation of Gaussian to Spherical Integral

The Gaussian integrand can also be split into parts that depend on radius $r = |\mathbf{x}|$ and parts that do not:

$$I_{a_1\dots a_n}^{g_n} = \int_{\mathbb{R}^n} \frac{1}{\sqrt{2\pi}^n} e^{-\frac{1}{2}r^2} r^{\sum a_i} \left(\prod_{i=1}^n \left(\frac{x_i}{r}\right)^{a_i} \right) \, \mathrm{d}^n \mathbf{x}.$$

Defining $\mathbf{x} = r\mathbf{u}$ so that $\mathbf{u} \in S_{n-1}$ gives $d^n \mathbf{x} = r^{n-1} dr d^{n-1} \mathbf{u}$ and

$$I_{a_1...a_n}^{g_n} = \int_0^\infty \frac{1}{\sqrt{2\pi}^n} e^{-\frac{1}{2}r^2} r^{\sum a_i} r^{n-1} \, \mathrm{d}r \int_{S_{n-1}} \left(\prod_{i=1}^n u_i^{a_i}\right) \, \mathrm{d}^{n-1} \mathbf{u}$$
$$= \frac{1}{\sqrt{2\pi}^{n-1}} I_{n-1+\sum a_i}^{h_1} I_{a_1...a_n}^{S_{n-1}}.$$

5 Evaluating the Spherical Integral

By the formula in the previous section,

$$I_{a_1...a_n}^{S_{n-1}} = \frac{\sqrt{2\pi}^{n-1} I_{a_1...a_n}^{g_n}}{I_{n-1+\sum a_i}^{h_1}}$$

The separability of the Gaussian integrals allows this to be written with only $I_a^{g_1}$ and $I_a^{h_1}$ terms:

$$I_{a_1...a_n}^{S_{n-1}} = \sqrt{2\pi}^{n-1} \frac{\prod_{i=1}^n I_{a_i}^{g_1}}{I_{n-1+\sum a_i}^{h_1}}.$$

6 Evaluating the Ball Integral

Integrating over a smaller sphere of radius r will introduce a factor of r^{n-1} from the change in surface area and a factor of $r^{\sum a_i}$ from the change in the monomial term itself. Thus,

$$I_{a_1...a_n}^{B_n} = \int_0^1 r^{n-1+\sum a_i} I_{a_1...a_n}^{S_{n-1}} \, \mathrm{d}r = \frac{1}{n+\sum a_i} I_{a_1...a_n}^{S_{n-1}}.$$

Using the formula from the previous section, this can be written in terms of $I_a^{g_1}$ and $I_a^{h_1}$:

$$I_{a_1...a_n}^{B_n} = \frac{\sqrt{2\pi}^{n-1}}{n+\sum a_i} \frac{\prod_{i=1}^n I_{a_i}^{g_1}}{I_{n-1+\sum a_i}^{h_1}}.$$