



Brookhaven
National Laboratory

BNL-104134-2014-TECH

AGS.SN258;BNL-104134-2014-IR

Vertical Chromaticity Study at 4.5 GeV/c in the AGS

L. Ratner

May 1989

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

8/3/89

Number 258

AGS Studies Report

Date(s) 5/29/89 Time(s) 0800 - 1200

Experimenter(s) L. Ahrens and L.G. Ratner

Reported by L.G. Ratner and L.A. Ahrens

Subject Vertical Chromaticity Study at 4.5 GeV/c in the AGS

We carried out a study that involved varying the ring sextupole magnets and measuring the resulting AGS chromaticity. The purpose of this test was to study the range of change available without affecting beam loss and beam size. We had seen large losses during the polarized proton run when attempting sextupole adjustment. It is thought that changing the AGS chromaticity during the fast tune jumps needed for polarized proton running could lead to needing less tune jump (see Appendix). The present experiment saw losses only when the tune spread in the beam became reasonably excessive. The study was done at a beam intensity of 3×10^{12} protons per pulse.

Figure 1 is a plot of the vertical tune as a function of radius and sextupole current at 8960 GCC (position of the $G\gamma = \nu$ polarized proton resonance). No changes were made to the horizontal sextupoles and the horizontal chromaticity was not measured. It appears that the radial pivot point and the vertical tune at that point are different for + and - sextupole currents. We do not understand this.

Figure 2 is a plot of the chromaticity as a function of current. The radial extremes were used for this plot. However, using other fits to the data does not significantly change this plot.

The tune measurements ignored the coupling between the horizontal and vertical planes which distorts the measure when the tunes cross, but is not important when they are very different. The radial measure was an average of the full set of horizontal PUEs. A revolution frequency measurement which would confirm this measure must be taken at fixed field and this was not done.

46 1320

K&E
10 X 10 TO 1/2 INCH
KEUFFEL & ESSER CO. MADE IN U.S.A.

$v_g(r) @ 8960 \text{ GCC } (G \rightarrow 2)$




FIG. 1

VERTICAL CLOUDINESS AT GØ=V (8960G00)




FIG. 2

46 1320

K&E 10 X 10 TO 1/4 INCH 7 X 10 INCHES
KIEFFEL & ESSER CO. MADE IN U.S.A.

Appendix

The condition for an intrinsic depolarizing resonance is $G\gamma = kn \pm \nu$, where $G = 1.79$, $\gamma = E/m$, $K = 12$ (for AGS), $n = \text{integer}$, and $\nu = \text{vertical tune}$. Because of energy spread in the beam, this condition is not normally satisfied for all particles simultaneously. Let (p_0, γ_0, ν_0) refer to the particle at the center of the beam momentum distribution. Consider the off-momentum particle whose momentum is $p = p_0 + \Delta p$, $\gamma = \gamma_0 + \Delta\gamma$, $\nu = \nu_0 + \Delta\nu$. For relativistic particles $\Delta\gamma/\gamma = \Delta p/p$ and by definition of chromaticity $\Delta\nu/\nu = \xi \Delta p/p$.

Now for this off-momentum particle, the two sides of the resonance condition become:

$$G\gamma \rightarrow G\gamma_0 + G\Delta\gamma = G\gamma_0 + G\gamma_0 \frac{\Delta p}{p}$$

$$kn \pm \nu \rightarrow kn \pm \nu_0 \pm \nu \xi \frac{\Delta p}{p}.$$

If these changes are equal, this particle will hit the resonance simultaneously with the on-momentum particle, a situation which makes resonance jumping most effective since the jump occurs at the same time for all the particles.

The change will be equal if:

$$G\gamma_0 \frac{\Delta p}{p} = \pm \nu_0 \xi \frac{\Delta p}{p},$$

a condition independent of $\Delta p/p$ and met for

$$\xi = \mp \frac{G\gamma_0}{\nu_0}.$$

In particular for $G\gamma = 0 + \nu$, $\xi = +1$ and for $G\gamma = 36 - \nu$, $\xi = -3.1$ ($\nu \sim 8.75$).