

## Exact Timestep for a Pairwise Coulomb Collision

S. Brooks

February 2025

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**  
USDOE Office of Science (SC), Nuclear Physics (NP)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-SC0012704 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

# Exact Timestep for a Pairwise Coulomb Collision

Stephen Brooks

February 24, 2025

## 1 Introduction

Standard numerical integrators work well for many-body Coulomb repulsion problems when the timestep is much shorter than the timescale of relative position changes. However, for ‘hard’ collisions in which two particles have a near miss and exchange a lot of momentum within one timestep, they understandably struggle. This note proposes using the exact solution of Keplerian two-body orbits (usually hyperbolic) to calculate the momentum exchange with other particles: either a selection of the ‘closest’ ones or all of them.

## 2 Pairwise Collision in Central Frame

Take the masses of two particles to be  $m_{1,2}$  and their charges to be  $q_{1,2}$ . Adopt the frame where the total momentum  $\mathbf{p} = m_1\mathbf{v}_1 + m_2\mathbf{v}_2 = \mathbf{0}$  and also the centre of mass  $m_1\mathbf{x}_1 + m_2\mathbf{x}_2 = \mathbf{0}$  is at the origin. From this, it is always true that  $\mathbf{x}_2 = -\frac{m_1}{m_2}\mathbf{x}_1$  and  $\mathbf{x}_1 - \mathbf{x}_2 = \frac{m_1+m_2}{m_2}\mathbf{x}_1$ . Similar relations hold for  $\mathbf{v}_1$ , meaning the problem is restricted to a two dimensional subspace spanned by  $\mathbf{x}_1$  and  $\mathbf{v}_1$ . The radial Coulomb force on the first particle is

$$\mathbf{F}_1(|\mathbf{x}_1|) = \frac{q_1q_2}{4\pi\epsilon_0} \frac{\mathbf{x}_1 - \mathbf{x}_2}{|\mathbf{x}_1 - \mathbf{x}_2|^3} = \frac{q_1q_2}{4\pi\epsilon_0} \frac{\frac{m_1+m_2}{m_2}\mathbf{x}_1}{\left(\frac{m_1+m_2}{m_2}\right)^3|\mathbf{x}_1|^3} = \frac{q_1q_2}{4\pi\epsilon_0} \left(\frac{m_2}{m_1+m_2}\right)^2 \frac{\mathbf{x}_1}{|\mathbf{x}_1|^3}.$$

Defining  $r = |\mathbf{x}_1|$ , this is a central force of magnitude

$$F_1(r) = \frac{q_1q_2}{4\pi\epsilon_0} \left(\frac{m_2}{m_1+m_2}\right)^2 \frac{1}{r^2}.$$

## 3 Binet Equation

The trajectory shape can be conveniently found from the Binet equation. Defining  $u = 1/r$  and using polar coordinates, it states, for a central force  $F$ ,

$$F = -mh^2u^2 \left( \frac{d^2u}{d\theta^2} + u \right),$$

where  $h = r^2\dot{\theta}$  is a conserved angular-momentum-like quantity. For this problem,  $F = ku^2$  where  $k = \frac{q_1q_2}{4\pi\epsilon_0} \left(\frac{m_2}{m_1+m_2}\right)^2$ . Thus,

$$ku^2 = -m_1h^2u^2 \left( \frac{d^2u}{d\theta^2} + u \right) \quad \Rightarrow \quad \frac{d^2u}{d\theta^2} + u = -\frac{k}{m_1h^2}$$

$$\Rightarrow u(\theta) = A \cos \theta + B \sin \theta - \frac{k}{m_1 h^2}$$

for some constants  $A, B$ . The origin for  $\theta$  is free to choose at this point, so it simplifies the calculation to choose it such that  $B = 0$  and  $A \geq 0$ . The solution is now

$$u(\theta) = A \cos \theta + C$$

where  $C = -\frac{k}{m_1 h^2}$  is a constant that is negative for repelling particles with  $q_1 q_2 > 0$  and positive for attracting particles with  $q_1 q_2 < 0$ .

## 4 Initial Conditions

Call the value of  $\theta$  corresponding to the initial condition  $\theta_0$ , so that

$$u(\theta_0) = \frac{1}{|\mathbf{x}_1|} = A \cos \theta_0 + C,$$

where in this section  $\mathbf{x}_1$  and  $\mathbf{v}_1$  are evaluated at the initial condition.

The first derivative satisfies

$$u' = \frac{du}{d\theta} = \frac{du}{dr} \frac{dr}{dt} \frac{dt}{d\theta} = \frac{-1}{r^2} \dot{r} \frac{1}{\dot{\theta}}.$$

Initially we have  $\dot{r} = \mathbf{v}_1 \cdot \mathbf{x}_1 / |\mathbf{x}_1|$  and  $\dot{\theta} = |\mathbf{x}_1 \times \mathbf{v}_1| / |\mathbf{x}_1|^2$ , so

$$u'(\theta_0) = \frac{-1}{|\mathbf{x}_1|^2} \frac{\mathbf{v}_1 \cdot \mathbf{x}_1}{|\mathbf{x}_1|} \frac{|\mathbf{x}_1|^2}{|\mathbf{x}_1 \times \mathbf{v}_1|} = \frac{-\mathbf{x}_1 \cdot \mathbf{v}_1}{|\mathbf{x}_1| |\mathbf{x}_1 \times \mathbf{v}_1|}.$$

These initial conditions also determine

$$h = r^2 \dot{\theta} = |\mathbf{x}_1|^2 \frac{|\mathbf{x}_1 \times \mathbf{v}_1|}{|\mathbf{x}_1|^2} = |\mathbf{x}_1 \times \mathbf{v}_1|,$$

which is a conserved quantity proportional to angular momentum. Using the solution for  $u(\theta)$  gives

$$u'(\theta_0) = \frac{-\mathbf{x}_1 \cdot \mathbf{v}_1}{|\mathbf{x}_1| h} = -A \sin \theta_0.$$

Taking the sum of squares gives

$$\begin{aligned} (A \cos \theta_0)^2 + (-A \sin \theta_0)^2 &= A^2 = \left( \frac{1}{|\mathbf{x}_1|} - C \right)^2 + \left( \frac{-\mathbf{x}_1 \cdot \mathbf{v}_1}{|\mathbf{x}_1| h} \right)^2 \\ \Rightarrow A &= \frac{1}{|\mathbf{x}_1|} \sqrt{(1 - |\mathbf{x}_1| C)^2 + \left( \frac{\mathbf{x}_1 \cdot \mathbf{v}_1}{h} \right)^2}. \end{aligned}$$

Taking the ratio gives

$$\frac{A \sin \theta_0}{A \cos \theta_0} = \tan \theta_0 = \frac{\frac{\mathbf{x}_1 \cdot \mathbf{v}_1}{|\mathbf{x}_1| h}}{\frac{1}{|\mathbf{x}_1|} - C} = \frac{\mathbf{x}_1 \cdot \mathbf{v}_1}{(1 - |\mathbf{x}_1| C) h}.$$

But also

$$\cos \theta_0 = \frac{\frac{1}{|\mathbf{x}_1|} - C}{A} = \frac{1 - |\mathbf{x}_1| C}{|\mathbf{x}_1| A}$$

is useful.

## 5 Time Dependence and the Eccentric Anomaly

As  $h = r^2\dot{\theta}$  is a constant found in the previous section,  $\dot{\theta} = \frac{d\theta}{dt} = hu^2$ . The equation to solve is

$$\frac{d\theta}{dt} = hu^2 = h(A \cos \theta + C)^2.$$

Kepler's eccentric anomaly  $E$  is defined via

$$\cos\{h\} E = \frac{e + \cos \theta}{1 + e \cos \theta}$$

for some eccentricity  $e$  that will be found later. Terms in  $\{\text{brackets}\}$  are used for the hyperbolic form when  $|e| > 1$ . The sign of  $\sin\{h\} E$  is defined to be the same as  $\sin \theta$ . The variable of the differential equation can be changed to  $E$  via the chain rule:

$$\begin{aligned} \frac{dE}{dt} &= \frac{dE}{d \cos\{h\} E} \frac{d \cos\{h\} E}{dt} \\ &= \frac{\{-\}1}{-\sin\{h\} E} \left( \frac{1}{1 + e \cos \theta} \frac{d \cos \theta}{dt} + (e + \cos \theta) \frac{-1}{(1 + e \cos \theta)^2} e \frac{d \cos \theta}{dt} \right) \\ &= \frac{\{-\}1}{-\sin\{h\} E} \left( \frac{1}{1 + e \cos \theta} + \frac{-e(e + \cos \theta)}{(1 + e \cos \theta)^2} \right) \frac{d \cos \theta}{dt} \\ &= \frac{\{-\}1}{-\sin\{h\} E} \frac{1 + e \cos \theta - e^2 - e \cos \theta}{(1 + e \cos \theta)^2} (-\sin \theta) h (A \cos \theta + C)^2 \\ &= \frac{\{-\} \sin \theta}{\sin\{h\} E} (1 - e^2) h \frac{(A \cos \theta + C)^2}{(1 + e \cos \theta)^2} \\ &= \frac{\sin \theta}{\sin\{h\} E} |1 - e^2| h \frac{(A \cos \theta + C)^2}{(1 + e \cos \theta)^2}. \end{aligned}$$

Next,  $\sin\{h\} E$  needs to be evaluated:

$$\begin{aligned} \sin\{h\} E &= \pm \sqrt{\{-\}(1 - \cos\{h\}^2 E)} = \pm \sqrt{\{-\} \left( 1 - \frac{(e + \cos \theta)^2}{(1 + e \cos \theta)^2} \right)} \\ &= \pm \sqrt{\{-\} \left( \frac{1 + 2e \cos \theta + e^2 \cos^2 \theta - e^2 - 2e \cos \theta - \cos^2 \theta}{(1 + e \cos \theta)^2} \right)} \\ &= \pm \frac{\sqrt{\{-\}(1 - e^2 - \cos^2 \theta + e^2 \cos^2 \theta)}}{1 + e \cos \theta} = \pm \frac{\sqrt{|1 - e^2|(1 - \cos^2 \theta)}}{1 + e \cos \theta} = \frac{\sqrt{|1 - e^2|} \sin \theta}{|1 + e \cos \theta|}. \end{aligned}$$

Substituting this back in,

$$\begin{aligned} \frac{dE}{dt} &= \frac{\sin \theta}{\sin\{h\} E} |1 - e^2| h \frac{(A \cos \theta + C)^2}{(1 + e \cos \theta)^2} = \frac{|1 + e \cos \theta|}{\sqrt{|1 - e^2|}} |1 - e^2| h \frac{(A \cos \theta + C)^2}{(1 + e \cos \theta)^2} \\ &= \sqrt{|1 - e^2|} h \frac{(A \cos \theta + C)^2}{|1 + e \cos \theta|} = \sqrt{|1 - e^2|} h C^2 \frac{(1 + \frac{A}{C} \cos \theta)^2}{|1 + e \cos \theta|}. \end{aligned}$$

Setting  $e = \frac{A}{C}$  gives

$$\frac{dE}{dt} = \sqrt{|1 - e^2|} h C^2 |1 + e \cos \theta|.$$

Note that this  $e$  has the same sign as  $C$ , so is the usual definition of eccentricity for attracting particles, but negative for repelling particles.

To express  $\frac{dE}{dt}$  back in terms of  $E$ , note that

$$\begin{aligned} \cos\{h\} E - \frac{1}{e} &= \frac{e - \frac{1}{e}}{1 + e \cos \theta} \\ \Rightarrow 1 + e \cos \theta &= \frac{e - \frac{1}{e}}{\cos\{h\} E - \frac{1}{e}} = \frac{e^2 - 1}{e \cos\{h\} E - 1}. \end{aligned}$$

Therefore

$$\frac{dE}{dt} = \sqrt{|1 - e^2|} h C^2 \frac{|1 - e^2|}{|1 - e \cos\{h\} E|} = \frac{|1 - e^2|^{3/2} h C^2}{|1 - e \cos\{h\} E|}.$$

Kepler's equation suggests an implicit solution of the form

$$qt = E - e \sin\{h\} E$$

for some constant  $q$ . Differentiating both sides with respect to  $t$  gives

$$q = \frac{dE}{dt} - e \cos\{h\} E \frac{dE}{dt} = (1 - e \cos\{h\} E) \frac{dE}{dt}$$

and substituting the  $\frac{dE}{dt}$  found above gives

$$q = (1 - e \cos\{h\} E) \frac{|1 - e^2|^{3/2} h C^2}{|1 - e \cos\{h\} E|} = \sigma |1 - e^2|^{3/2} h C^2,$$

which is indeed a constant. The sign  $\sigma$  of  $1 - e \cos\{h\} E$  is

- 1 for elliptic orbits ( $0 \leq e < 1$ ),
- $-1$  for hyperbolic attractive orbits ( $e > 1, C > 0$ ) and
- 1 for hyperbolic repulsive orbits ( $e < -1, C < 0$ ).

## 6 Time Step

With  $A$  and  $\theta_0$  known, calculate  $e = \frac{A}{C}$  and proceed via

$$\cos\{h\} E_0 = \frac{e + \cos \theta_0}{1 + e \cos \theta_0}, \quad qt_0 = E_0 - e \sin\{h\} E_0$$

to find the initial time  $t_0$  in Kepler's equation.

Suppose the position is required at  $t = t_0 + \delta t$  for some time step  $\delta t$ . A solution of  $qt = E - e \sin\{h\} E$  is needed. This does not have a closed-form solution but Newton–Raphson iterations exist (see Appendices). Then

$$\cos \theta = \frac{e - \cos\{h\} E}{e \cos\{h\} E - 1}, \quad \sin \theta = \frac{|1 + e \cos \theta|}{\sqrt{|1 - e^2|}} \sin\{h\} E$$

and

$$u(\theta) = A \cos \theta + C.$$

A convenient way of handling the frames of reference is to rotate  $\mathbf{x}_1$  by angle  $\theta - \theta_0$  about the  $\mathbf{x}_1 \times \mathbf{v}_1$  axis and then scale by  $r/r_0 = u(\theta_0)/u(\theta)$ .

To find the velocity, it is easier to use the known differential equations  $\dot{\theta} = hu^2$  and  $u'(\theta) = -A \sin \theta$  to get

$$\frac{dr}{dt} = \frac{dr}{du} \frac{du}{d\theta} \frac{d\theta}{dt} = \frac{-1}{u^2} (-A \sin \theta) hu^2 = Ah \sin \theta.$$