

Force Between Two Moving, Non-Accelerating Charges

S. Brooks

October 2024

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy
USDOE Office of Science (SC), Nuclear Physics (NP)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-SC0012704 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Force Between Two Moving, Non-Accelerating Charges

Stephen Brooks

October 28, 2024

1 Non-Moving Charges

The electric potential and field from a stationary charge q_1 placed at the origin are

$$V(\mathbf{x}) = \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}|} \quad \text{and} \quad \mathbf{E}(\mathbf{x}) = -\nabla V = \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}|^3} \mathbf{x}.$$

The force on another charge q_2 at position \mathbf{x} is then

$$\mathbf{F}_2 = q_2(\mathbf{E} + \mathbf{v}_2 \times \mathbf{B}) = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{|\mathbf{x}|^3} \mathbf{x},$$

where $\mathbf{B} = \mathbf{0}$ because the non-moving charge q_1 produces no magnetic field.

2 Transforming the Four-Potential

The electromagnetic four-potential is $A^\mu = (\frac{1}{c}V, \mathbf{A})$, which transforms as a four-vector under Lorentz transformations. Let primes ($'$) denote the rest frame of charge q_1 . In this frame,

$$A'^\mu(x'^\mu) = \left(\frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|}, \mathbf{0} \right) = \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|} (1, \mathbf{0}).$$

If this charge is travelling at velocity $\mathbf{v}_1 = \boldsymbol{\beta}c$, the (inverse) Lorentz transformation is

$$x^\mu = \begin{bmatrix} \gamma & \gamma\boldsymbol{\beta}^T \\ \gamma\boldsymbol{\beta} & I + \frac{\gamma-1}{\beta^2}\boldsymbol{\beta}\boldsymbol{\beta}^T \end{bmatrix} x'^\mu,$$

where $\beta = |\boldsymbol{\beta}|$ and $\gamma = (1 - \beta^2)^{-\frac{1}{2}}$. The same transformation gives $A^\mu(x^\mu)$ from $A'^\mu(x'^\mu)$, so

$$A^\mu(x^\mu) = \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|} (\gamma, \gamma\boldsymbol{\beta}).$$

The forward Lorentz transformation on $x^\mu = (ct, \mathbf{x})$ gives

$$x'^\mu = \left(\gamma ct - \gamma\boldsymbol{\beta} \cdot \mathbf{x}, -\gamma ct\boldsymbol{\beta} + \mathbf{x} + \frac{\gamma-1}{\beta^2}(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} \right),$$

thus $\mathbf{x}' = -\gamma ct\boldsymbol{\beta} + \mathbf{x} + \frac{\gamma-1}{\beta^2}(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta}$.

3 Deriving the Fields

The field tensor is $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$, where $\partial^\mu = (\frac{1}{c}\partial_t, -\nabla)$. The four-potential is a function of \mathbf{x}' , so first calculate the derivatives

$$\begin{aligned}\partial^0 x'^i &= -\gamma\beta^i \\ \partial^j x'^i &= -\left(\delta_{ij} + \frac{\gamma-1}{\beta^2}\beta^i\beta^j\right).\end{aligned}$$

Next observe that

$$\partial^\mu |\mathbf{x}'| = \sum_i \frac{\partial |\mathbf{x}'|}{\partial x'^i} \partial^\mu x'^i = \sum_i \frac{x'^i}{|\mathbf{x}'|} \partial^\mu x'^i$$

and use the chain rule to get

$$\begin{aligned}\partial^\mu A^\nu &= \frac{1}{4\pi\epsilon_0 c} \frac{-q_1}{|\mathbf{x}'|^2} (\gamma, \gamma\boldsymbol{\beta})^\nu \sum_i \frac{x'^i}{|\mathbf{x}'|} \partial^\mu x'^i \\ &= \frac{1}{4\pi\epsilon_0 c} \frac{-q_1}{|\mathbf{x}'|^3} (\gamma, \gamma\boldsymbol{\beta})^\nu \sum_i x'^i \left(-\gamma\beta^i, -\left(\mathbf{e}_i + \frac{\gamma-1}{\beta^2}\beta^i\boldsymbol{\beta}\right)\right)^\mu \\ &= \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|^3} (\gamma, \gamma\boldsymbol{\beta})^\nu \left(\gamma\mathbf{x}' \cdot \boldsymbol{\beta}, \mathbf{x}' + \frac{\gamma-1}{\beta^2}(\mathbf{x}' \cdot \boldsymbol{\beta})\boldsymbol{\beta}\right)^\mu.\end{aligned}$$

The fields and forces on the particles will be evaluated at $t = 0$, so there are simpler expressions for

$$\begin{aligned}\mathbf{x}' &= \mathbf{x} + \frac{\gamma-1}{\beta^2}(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta}, \\ \mathbf{x}' \cdot \boldsymbol{\beta} &= \mathbf{x} \cdot \boldsymbol{\beta} + (\gamma-1)(\boldsymbol{\beta} \cdot \mathbf{x}) = \gamma\boldsymbol{\beta} \cdot \mathbf{x}, \\ \mathbf{x}' + \frac{\gamma-1}{\beta^2}(\mathbf{x}' \cdot \boldsymbol{\beta})\boldsymbol{\beta} &= \mathbf{x} + \frac{\gamma-1}{\beta^2}(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} + \frac{\gamma-1}{\beta^2}(\gamma\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} \\ &= \mathbf{x} + \frac{(\gamma-1)(\gamma+1)}{\beta^2}(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} = \mathbf{x} + \gamma^2(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta}.\end{aligned}$$

This means at $t = 0$,

$$\partial^\mu A^\nu = \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|^3} (\gamma, \gamma\boldsymbol{\beta})^\nu (\gamma^2\boldsymbol{\beta} \cdot \mathbf{x}, \mathbf{x} + \gamma^2(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta})^\mu.$$

The electric field is

$$\begin{aligned}\frac{1}{c}E^i &= F^{i0} = \partial^i A^0 - \partial^0 A^i \\ \Rightarrow \mathbf{E} &= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} (\gamma(\mathbf{x} + \gamma^2(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta}) - \gamma\boldsymbol{\beta}(\gamma^2\boldsymbol{\beta} \cdot \mathbf{x})) \\ &= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} (\gamma\mathbf{x} + \gamma^3(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} - \gamma^3(\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta}) \\ &= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \gamma\mathbf{x}.\end{aligned}$$

The x component of the magnetic field is

$$B^x = F^{zy} = \partial^z A^y - \partial^y A^z$$

$$\begin{aligned}
\Rightarrow B^x &= \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|^3} (\gamma\beta^y (z + \gamma^2(\boldsymbol{\beta} \cdot \mathbf{x})\beta^z) - \gamma\beta^z (y + \gamma^2(\boldsymbol{\beta} \cdot \mathbf{x})\beta^y)) \\
&= \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|^3} (\gamma\beta^y z - \gamma\beta^z y).
\end{aligned}$$

This is true cyclically $x \rightarrow y \rightarrow z \rightarrow x$, so

$$\mathbf{B} = \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|^3} \gamma \boldsymbol{\beta} \times \mathbf{x}.$$

4 Evaluating the Force

Taking these results together, the force on charge q_2 at position \mathbf{x} , travelling at velocity \mathbf{v}_2 is

$$\begin{aligned}
\mathbf{F}_2 &= q_2(\mathbf{E} + \mathbf{v}_2 \times \mathbf{B}) \\
&= \frac{1}{4\pi\epsilon_0 c} \frac{q_1 q_2}{|\mathbf{x}'|^3} (c\gamma_1 \mathbf{x} + \mathbf{v}_2 \times (\gamma_1 \boldsymbol{\beta}_1 \times \mathbf{x})) \\
&= \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{|\mathbf{x}'|^3} \gamma_1 (\mathbf{x} + \boldsymbol{\beta}_2 \times (\boldsymbol{\beta}_1 \times \mathbf{x})) \\
&= \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{|\mathbf{x}'|^3} \gamma_1 (\mathbf{x} + (\boldsymbol{\beta}_2 \cdot \mathbf{x})\boldsymbol{\beta}_1 - (\boldsymbol{\beta}_2 \cdot \boldsymbol{\beta}_1)\mathbf{x}),
\end{aligned}$$

where $\mathbf{x}' = \mathbf{x} + \frac{\gamma_1 - 1}{\beta_1^2} (\boldsymbol{\beta}_1 \cdot \mathbf{x})\boldsymbol{\beta}_1$.

5 Check: using the E, B field transformation rules

In the rest frame of charge q_1 ,

$$\mathbf{E}'(\mathbf{x}') = \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \mathbf{x}'$$

and $\mathbf{B}' = \mathbf{0}$. The field transformation rules are

$$\begin{aligned}
\mathbf{E}_{\parallel} &= \mathbf{E}'_{\parallel}, \\
\mathbf{B}_{\parallel} &= \mathbf{B}'_{\parallel}, \\
\mathbf{E}_{\perp} &= \gamma(\mathbf{E}'_{\perp} - \mathbf{v} \times \mathbf{B}'), \\
\mathbf{B}_{\perp} &= \gamma\left(\mathbf{B}'_{\perp} + \frac{1}{c^2} \mathbf{v} \times \mathbf{E}'\right),
\end{aligned}$$

where \parallel and \perp denote the components parallel and perpendicular to \mathbf{v} respectively.

$$\begin{aligned}
\mathbf{E} &= \mathbf{E}_{\parallel} + \mathbf{E}_{\perp} = \mathbf{E}'_{\parallel} + \gamma\mathbf{E}'_{\perp} \\
&= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \left(\frac{1}{\beta^2} (\mathbf{x}' \cdot \boldsymbol{\beta})\boldsymbol{\beta} + \gamma \left(\mathbf{x}' - \frac{1}{\beta^2} (\mathbf{x}' \cdot \boldsymbol{\beta})\boldsymbol{\beta} \right) \right) \\
&= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \left(\gamma\mathbf{x}' - \frac{\gamma - 1}{\beta^2} (\mathbf{x}' \cdot \boldsymbol{\beta})\boldsymbol{\beta} \right).
\end{aligned}$$

Using the formulae for \mathbf{x}' in terms of \mathbf{x} at $t = 0$ from a previous section,

$$\begin{aligned}
\mathbf{E} &= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \left(\gamma \left(\mathbf{x} + \frac{\gamma - 1}{\beta^2} (\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} \right) - \frac{\gamma - 1}{\beta^2} (\gamma\boldsymbol{\beta} \cdot \mathbf{x})\boldsymbol{\beta} \right) \\
&= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \gamma \mathbf{x}.
\end{aligned}$$

For the magnetic field,

$$\begin{aligned}\mathbf{B} &= \mathbf{B}_{\parallel} + \mathbf{B}_{\perp} = 0 + \gamma \frac{1}{c^2} \mathbf{v} \times \mathbf{E}' \\ &= \frac{1}{4\pi\epsilon_0} \frac{q_1}{|\mathbf{x}'|^3} \gamma \frac{1}{c^2} \mathbf{v} \times \mathbf{x}' \\ &= \frac{1}{4\pi\epsilon_0 c} \frac{q_1}{|\mathbf{x}'|^3} \gamma \boldsymbol{\beta} \times \mathbf{x},\end{aligned}$$

where we have used $\mathbf{v} \times \mathbf{x}' = \mathbf{v} \times \mathbf{x}$ since $\mathbf{v} \parallel \boldsymbol{\beta}$ and \mathbf{x}' and \mathbf{x} only differ by a multiple of $\boldsymbol{\beta}$.