

## Emittance growth from electron beam modulation

M. Blaskiewicz,

December 2009

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

C-A/AP/#363  
December 2009

# Emittance Growth from Electron Beam Modulation

M. Blaskiewicz



**Collider-Accelerator Department  
Brookhaven National Laboratory  
Upton, NY 11973**

Notice: This document has been authorized by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this document, or allow others to do so, for United States Government purposes.

# Emittance Growth from Electron Beam Modulation

M. Blaskiewicz\*  
BNL, Upton NY 11973, USA

In linac ring colliders like MeRHIC and eRHIC a modulation of the electron bunch can lead to a modulation of the beam beam tune shift and steering errors [1–3]. These modulations can lead to emittance growth. This note presents simple formulas to estimate these effects which generalize some previous results.

## I. THEORY OF QUADRUPOLE GROWTH

Consider one dimensional motion in the  $x$  direction with bare tune  $Q$ . We use the evolution variable  $\theta$  which updates by  $2\pi$  each turn. The interaction point is at  $\theta = 0$ , the modulating beam beam force is denoted by  $\epsilon(\theta)$  and the hamiltonian is given by

$$H(x, p, \theta) = \frac{p^2 + Q^2 x^2}{2} + \delta_p(\theta) \frac{Q x^2}{2} \epsilon(\theta), \quad (1)$$

where the periodic delta function is

$$\delta_p(\theta) = \sum_{k=-\infty}^{\infty} \delta(\theta - 2\pi k).$$

The equations of motion are  $dx/d\theta = \partial H/\partial p$  and  $dp/d\theta = -\partial H/\partial x$ . Do a canonical transformation to action angle variables with  $x = \sqrt{2J/Q} \sin \psi$  and  $p = \sqrt{2JQ} \cos \psi$  yielding

$$H(\psi, J, \theta) = QJ + \delta_p(\theta) J \sin^2 \psi \epsilon(\theta). \quad (2)$$

Averaging over  $\psi$  and  $\epsilon$  with  $\langle \delta_p(\theta) \rangle = 1/2\pi$  one finds  $\langle H(J) \rangle = QJ + J \langle \epsilon \rangle / 4\pi$  so the average beam beam tune shift is  $\langle \delta Q \rangle = \langle \epsilon \rangle / 4\pi$ . We redefine the tune to be  $Q \rightarrow Q + \langle \delta Q \rangle$  and introduce the fluctuating part of the noise.  $\delta\epsilon = \epsilon - \langle \epsilon \rangle$ . The hamiltonian becomes

$$H(\psi, J, \theta) = QJ + \delta_p(\theta) J \sin^2 \psi \delta\epsilon(\theta). \quad (3)$$

The emittance growth is due to the modulation in  $\delta\epsilon$  and the equations of motion are

$$\frac{dJ}{d\theta} = -\frac{\partial H}{\partial \psi} = -\delta_p(\theta) J \sin(2\psi) \delta\epsilon(\theta)$$

and

$$\frac{d\psi}{d\theta} = \frac{\partial H}{\partial J} = Q + \delta_p(\theta) \sin^2 \psi \delta\epsilon(\theta).$$

The expansion parameter is  $\delta\epsilon$ . To zeroeth order  $\psi_0(\theta) = Q\theta + \hat{\psi}$ . To first order in  $\delta\epsilon$

$$\psi_1(\theta) = Q\theta + \hat{\psi} + \int_0^\theta \delta_p(s) \sin^2(Qs + \hat{\psi}) \delta\epsilon(s) ds \quad (4)$$

$$\approx Q\theta + \hat{\psi} - \frac{1}{2} \int_0^\theta \delta_p(s) \cos(2Qs + 2\hat{\psi}) \delta\epsilon(s) ds. \quad (5)$$

---

\*Electronic address: [blaskiewicz@bnl.gov](mailto:blaskiewicz@bnl.gov)

The evolution of the action is given by

$$\frac{d \ln J}{d\theta} = -\delta_p(\theta) \sin(2\psi) \delta\epsilon(\theta) \quad (6)$$

$$\approx -\delta_p(\theta) \sin(2\psi_1(\theta)) \delta\epsilon(\theta) \quad (7)$$

$$\approx -\delta_p(\theta) \sin(2Q\theta + 2\hat{\psi}) \delta\epsilon(\theta) + \delta_p(\theta) \cos(2Q\theta + 2\hat{\psi}) \delta\epsilon(\theta) \int_0^\theta \delta_p(s) \cos(2Qs + 2\hat{\psi}) \delta\epsilon(s) ds \quad (8)$$

$$\approx -\delta_p(\theta) \sin(2Q\theta + 2\hat{\psi}) \delta\epsilon(\theta) + \frac{1}{2} \frac{d}{d\theta} \left[ \int_0^\theta \delta_p(s) \cos(2Qs + 2\hat{\psi}) \delta\epsilon(s) ds \right]^2. \quad (9)$$

Integrating equation (9) one finds  $J(\theta) = J(0) \exp(u(\theta))$  where

$$u(\theta) = - \int_0^\theta \delta_p(s) \sin(2Qs + 2\hat{\psi}) \delta\epsilon(s) + \frac{1}{2} \left[ \int_0^\theta \delta_p(s) \cos(2Qs + 2\hat{\psi}) \delta\epsilon(s) ds \right]^2.$$

As long as  $u$  is small compared to one, which is necessary for eq (8), the first term on the right hand side dominates. Hence  $u$  is effectively the sum of a large number of uncorrelated random variables and therefore nearly a gaussian random variable. Notice that

$$\langle u(\theta) \rangle = \left\langle \frac{1}{2} \left[ \int_0^\theta \delta_p(s) \cos(2Qs + 2\hat{\psi}) \delta\epsilon(s) ds \right]^2 \right\rangle \approx \sigma_u^2/2$$

Now since  $\langle \exp(u) \rangle = \exp(\langle u \rangle + \sigma_u^2/2)$  for a gaussian, we have  $\langle J(\theta) \rangle = J(0) \exp(2 \langle u(\theta) \rangle)$  and the problem is to evaluate  $\langle u(\theta) \rangle$ . Start by using the periodic delta function to turn the integral to a sum and define  $\delta\epsilon(2\pi k) = \delta\epsilon_k$  so that

$$\langle 2u_n \rangle = \sum_{k=0}^n \sum_{m=0}^n \cos(4\pi Qk + 2\hat{\psi}) \cos(4\pi Qm + 2\hat{\psi}) \langle \delta\epsilon_k \delta\epsilon_m \rangle \quad (10)$$

$$\approx \sum_{k=0}^n \sum_{m=0}^n \frac{1}{2} \cos(4\pi Q[k - m]) \langle \delta\epsilon_k \delta\epsilon_m \rangle \quad (11)$$

$$\approx n \sum_{m=-\infty}^{\infty} \frac{1}{2} \cos(4\pi Qm) \langle \delta\epsilon_k \delta\epsilon_{m+k} \rangle \quad (12)$$

where  $\delta\epsilon_k$  is assumed to be stationary noise. As a simple model take  $\langle \delta\epsilon_k \delta\epsilon_{m+k} \rangle = \sigma_\epsilon^2 \exp(-\alpha|m|)$ . In this case the sum can be done resulting in

$$\ln \left\langle \frac{J(\theta)}{J(0)} \right\rangle = \langle 2u_n \rangle = n \frac{\sigma_\epsilon^2}{2} \frac{1 - e^{-2\alpha}}{1 + e^{-2\alpha} - 2 \cos(4\pi Q) e^{-\alpha}} \quad (13)$$

## II. COMPARISON WITH SIMULATIONS

For simulation purposes I take a one turn matrix followed by a thin lens beam beam kick,

$$x_n(k+1) = \cos \psi_n x_n(k) + \sin \psi_n p_n(k) \quad (14)$$

$$p_n(k+1) = \cos \psi_n p_n(k) - \sin \psi_n x_n(k) + \epsilon_k x_n(k+1) \quad (15)$$




FIG. 1: Comparison of simulation and equation 12. The correlation suppression factor is 0.06.

where  $k$  is the time like variable and each of the  $n$  particles has a slightly different phase advance to create filamentation. The beam beam kick  $\epsilon_k$  is normalized as above. Comparison of a simulation and equation (13) is shown in Fig 1. The correlation time of the noise was  $\alpha = 0.1$  and  $Q = 0.2$  resulting in a growth rate only 0.06 of what it would be for white noise of the same amplitude. As one can see from the figure the agreement is very good over many e-folding times. Of course in a real beam the electron kick is not purely linear so the growth would be reduced. In any case the formula seems reliable for an e-folding time.

### III. GROWTH DUE TO STEERING ERRORS

Along with variations in electron beam intensity and emittance there can also be steering errors. We start by writing the one turn map in the linear approximation

$$x(k+1) = \cos \psi x(k) + \sin \psi x'(k) \beta^* \quad (16)$$

$$x'(k+1) = \cos \psi x'(k) - \sin \psi x(k) / \beta^* - \epsilon [x(k+1) - y(k+1)] / \beta^* \quad (17)$$

where  $y(k)$  is the electron beam offset,  $\beta^*$  is the beta function at the crossing point, and the beam beam tune shift is  $\epsilon/4\pi$  as before. We neglect the beam beam tune shift and define  $z(k) = x(k) + i\beta^* x'(k)$  then

$$z(k+1) = e^{-i\psi} z(k) + i\epsilon y(k+1). \quad (18)$$

Set  $U(k) = z(k) \exp(ik\psi)$  yielding

$$U(k+1) = U(k) + i\epsilon y(k+1) \exp(i(k+1)\psi), \quad (19)$$

so that

$$U(k) = U_0 + i\epsilon \sum_{m=0}^k y(m) \exp(im\psi). \quad (20)$$

Taking averages

$$\langle |U(k)|^2 \rangle = |U_0|^2 + \epsilon^2 \sum_m \sum_\ell \langle y(m)y(\ell) \rangle \exp(i(m-\ell)\psi) \quad (21)$$

$$\approx |U_0|^2 + k\epsilon^2 \sum_{m=-\infty}^{\infty} \langle y(m+\ell)y(\ell) \rangle \cos(m\psi) \quad (22)$$

Taking  $\langle y(m + \ell)y(\ell) \rangle = \sigma_y^2 \exp(-\alpha|m|)$  one finds

$$2\epsilon_{rms}\beta^* = \langle |U(k)|^2 \rangle = |U_0|^2 + k(\epsilon\sigma_y)^2 \frac{1 - e^{-2\alpha}}{1 + e^{-2\alpha} - 2\cos(2\pi Q)e^{-\alpha}}, \quad (23)$$

where  $\epsilon_{rms} = \langle x^2 \rangle / \beta^*$  is the rms geometric emittance of the ion beam.

### Acknowledgements

Thanks to Vadim Ptitsyn for useful discussions and encouragement.

- 
- [1] C. Montag, RHIC/AP/143, 2004 also C. Montag NIM A, Volume 581, p581, 2007
  - [2] See Eduard Pozdeyev's MeRHIC discussion September 7, 2006.
  - [3] Y. Hao, V. Litvinenko, C. Montag, E. Pozdeyev, V. Ptitsyn, PAC07, p1865 (2007).