



BNL-104054-2014-TECH

AGS.SN177;BNL-104054-2014-IR

Further Studies of the Fast Injection Bump and Comparison with Analytic Expressions and with Results from the BEAM Program

L. Ahrens

February 1985

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

AGS Studies Report

Date(s) February 27, 1985 Time(s) 0800-1000
 Experimenter(s) L. Ahrens and C. Gardner
 Reported by C. Gardner
 Subject Further Studies of the Fast Injection Bump and
Comparison with Analytic Expressions and with
Results from the BEAM Program

Observations and Conclusion

The purpose of this study was to examine in some detail the dependence of both the sine and cosine components of the equilibrium orbit (E.O.) harmonics on the relative magnitudes of the kicks at A13 and B7. Data were obtained in the same way as in the previous studies (AGS Studies Reports 171, 172) using the ring PUE system and the ORBED/NORB programs. The NORB program analyzes the E.O. and arbitrarily defines the angular zero for its fourier analysis as straight section (S.S.) L20. With the magnitude of the A13 kick fixed at 1.5 V the B7 kick was swept through a series of values ranging from 0.6 V to 3.0 V. The sine and cosine components of the 9th harmonic of the E.O. obtained for each case are listed in Table I and are plotted in Figure 1. These data were obtained with a horizontal tune of 8.75. Note that the sine and cosine components appear to vary linearly with the magnitude of the kick at B7. Fourier analysis of the analytic expression for the deformed E.O. shows that this is indeed the case, which we now show.

The general expression for the deformed E.O. is¹

$$\eta(\phi) = \frac{v}{2 \sin \pi v} \int_{\phi}^{\phi+2\pi} F(\psi) \cos v(\pi+\phi-\psi) d\psi \quad (1)$$

where

$$\phi = \phi(s) = \int_0^s \frac{ds}{v\beta(\bar{s})}, \quad (2)$$

$$f(\psi) = \beta^{3/2}(s) F(s), \quad (3)$$

¹E.D. Courant and H.S. Snyder, Annals of Physics: 3, 1-48 (1958).

s is the distance along the undeformed orbit of circumference c , $F(s)$ is the perturbation term due to perturbing dipoles in the ring, and the other symbols have their usual meanings. For the case of two delta function kicks at s_1 and s_2 respectively ($0 < s_1 < s_2 < c$), $F(s)$ is a periodic function of the form

$$F(s) = -\theta_1 \delta(s-s_1) - \theta_2 \delta(s-s_2), \quad 0 < s < c \quad (4)$$

where θ_1 (θ_2) is the magnitude of the kick at s_1 (s_2). Inserting (4) into (3) and (1), the radial displacement

$$y(s) = \beta^{1/2}(s) \eta(\phi) \quad (5)$$

of the E.O. due to the perturbing kicks is

$$y(s) = \frac{\beta^{1/2}(s)}{2 \sin \pi \nu} [\Delta_1 \cos \nu(\phi - \psi_1 + \pi) + \Delta_2 \cos \nu(\phi - \psi_2 + \pi)], \quad 0 < s < s_1$$

$$y(s) = \frac{\beta^{1/2}(s)}{2 \sin \pi \nu} [\Delta_1 \cos \nu(\phi - \psi_1 - \pi) + \Delta_2 \cos \nu(\phi - \psi_2 + \pi)], \quad s_1 < s < s_2 \quad (6)$$

$$y(s) = \frac{\beta^{1/2}(s)}{2 \sin \pi \nu} [\Delta_1 \cos \nu(\phi - \psi_1 - \pi) + \Delta_2 \cos \nu(\phi - \psi_2 - \pi)], \quad s_2 < s < c$$

where

$$\Delta_1 \equiv -\theta_1 \beta^{1/2}(s_1), \quad \Delta_2 \equiv -\theta_2 \beta^{1/2}(s_2) \quad (7)$$

$$\psi_1 \equiv \psi(s_1) = \int_0^{s_1} \frac{d\bar{s}}{\nu \beta(\bar{s})}, \quad \psi_2 \equiv \psi(s_2) = \int_0^{s_2} \frac{d\bar{s}}{\nu \beta(\bar{s})} \quad (8)$$

Now we consider the specific case at hand in which s_1 is the location (along the unperturbed E.O.) of the A13 kick and s_2 is the location of the B7 kick. θ_1 and θ_2 are respectively the magnitudes of these kicks and $\beta(s_1)$ and $\beta(s_2)$ are the values of the betatron function at A13 and B7. The locations of these kicks have been chosen so that nominally they produce a half-lambda bump. Thus, we have

$$\nu \psi_2 = \nu \psi_1 + \pi. \quad (9)$$

The location of the A13 kick (W.R.T. S.S. L20) is also such that one has nominally

$$\nu\psi_1 = \pi. \quad (10)$$

(This is because we happen to inject at A20, one superperiod away from our angular zero reference (S.S. L20) or $1/12 (2\pi\nu) \approx 3\pi/2$ away, and the kick wants to be $\lambda/4 = \pi/2$ away from injection--hence placing the A13 kick at π .)

Putting (9) and (10) into (6) and letting the tune $\nu = 8.75$ we find

$$y(s) = \frac{\beta^{1/2}(s)}{2 \sin \pi\nu} (\Delta_1 - \Delta_2)(\sqrt{2}/2) [\cos \nu\phi + \sin \nu\phi], \quad 0 < s < s_1 \quad (11)$$

$$y(s) = \frac{\beta^{1/2}(s)}{2 \sin \pi\nu} (\sqrt{2}/2) [(\Delta_1 - \Delta_2) \cos \nu\phi - (\Delta_1 + \Delta_2) \sin \nu\phi], \quad s_1 < s < s_2$$

$$y(s) = \frac{\beta^{1/2}(s)}{2 \sin \pi\nu} (\Delta_1 - \Delta_2)(\sqrt{2}/2) [\cos \nu\phi - \cos \nu\phi], \quad s_2 < s < c.$$

The fourier sine and cosine components of the perturbed orbit $y(s)$ are respectively

$$A_N = \frac{1}{\pi} \int_0^{2\pi} d\phi y(s) \sin N\phi \quad (12)$$

$$B_N = \frac{1}{\pi} \int_0^{2\pi} d\phi y(s) \cos N\phi$$

Taking $N = 9$ and putting (11) into (12) with $\beta^{1/2}(s)$ replaced by the average, $\beta_{AV}^{1/2}$, we find the following values for the ninth harmonic of the deformed E.O.:

$$A_9 = (\beta_{AV}^{1/2}/2 \sin \pi\nu) \frac{1}{\pi} (\sqrt{2}/2) [-0.34 \Delta_1] (\text{sine}) \quad (13)$$

$$B_9 = (\beta_{AV}^{1/2}/2 \sin \pi\nu) \frac{1}{\pi} (\sqrt{2}/2) [3.97(\Delta_1 - \Delta_2) + 0.02(\Delta_1 + \Delta_2)] (\text{cosine})$$

(We note here that Equations (13) are exact only when $v = 8.75$ and (9) and (10) are satisfied. In practice (9) and (10) are only approximately true, so in general there will be small corrections to (13)). Thus we see that with the magnitude (θ_1) of the kick at A13 fixed, the cosine component varies linearly with the magnitude (θ_2) of the kick at B7, while the sine component remains constant. Our observations (see Figure 1) are consistent with these results.

We note from Equations (11) that the residuals will have zero amplitude when $\Delta_1 = \Delta_2$. Thus, using (7), the condition for nulled residuals is

$$(\theta_2/\theta_1)^2 = \beta(s)_1/\beta(s)_2. \quad (14)$$


Comparing (13) and Figure 1, we see that the value of the B7 kick for which the cosine component is zero is very nearly the value for which $\Delta_1 = \Delta_2$. Thus, our measured results show that the residuals are nulled when $(\theta_2/\theta_1)^2 = (1.95/1.5)^2 = 1.7$, which agrees with the result obtained in AGS Studies Report No. 172, and as mentioned there is slightly less than the value of 2.1 one expects if $\beta_H(A13) = \beta_{\max} \approx 22 \text{ m}$, $\beta_H(B7) = \beta_{\min} \approx 10.5 \text{ m}$.

Figure 2 shows the E.O.'s obtained from the ring PUE system for three cases. In case (b) the magnitude of the A13 and B7 kicks have been adjusted to null the residuals. In case (a) the B7 kick is smaller than the value required to null the residuals while in case (c) it is larger than the value required for a null. Equilibrium orbits obtained by the BEAM program for the three corresponding cases are shown in Figure 3. Comparison of the corresponding cases in the two figures shows that the general features of the orbits agree quite well. (Note especially the behavior near the A13 and B7 kicks).




Table I

<u>Kick Magnitudes</u> (volts)		<u>Ninth Harmonic of</u> <u>Deformed E.O.</u> (mm)	
<u>A13</u>	<u>B7</u>	<u>Sin</u>	<u>Cos</u>
1.5	0.6	-0.06	+0.45
1.5	1.0	-0.07	+0.33
1.5	1.3	-0.08	+0.22
1.5	1.5	-0.07	+0.14
1.5	1.8	-0.08	+0.02
1.5	2.0	-0.09	-0.02
1.5	2.1	-0.09	-0.07
1.5	2.2	-0.09	-0.10
1.5	2.3	-0.10	-0.15
1.5	2.5	-0.10	-0.23
1.5	3.0	-0.10	-0.36

FIGURE 1



- 7 -
FIGURE 2



- 8 -
FIGURE 3

