

The simulation of Siberian Snakes based on calculated three dimensional magnet field

X. Meiqin

December 1997

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Alternating Gradient Synchrotron Department
Relativistic Heavy Ion Collider Project
BROOKHAVEN NATIONAL LABORATORY
Upton, New York 11973

Spin Note

AGS/RHIC/SN No. 070

**The simulation of Siberian Snakes
based on calculated three dimensional magnet field**

X. Meiqin, T. Katayama
(Center for Nuclear Study, The University of Tokyo/RIKEN)

December 3, 1997

The simulation of Siberian Snakes based on calculated three dimensional magnetic field

Xiao Meiqin T. Katayama

(Center for Nuclear Study, The University of Tokyo/RIKEN)

1. Introduction

As a special kind of transport elements in storage ring, Siberian Snakes in RHIC, which consist of four right-handed helical dipole magnets, are expected to rotate spin direction of polarized proton to 180° , and should not affect the beam orbit at the same time, the axis of spin precession should make an angle $\phi = \pm 45^\circ$ too^{[1][2]}. However, the problem is that actually they can not make the spin direction rotate 180° perfectly, also the fringing magnetic field will affect the beam trajectories.

The simulation in Siberian Snakes based on analytical formulation (such as Blewett-Chasman formula) of magnetic field had been well done^[3]. For better representation of the real Snakes in RHIC, which are still under the construction, the simulation based on three dimensional magnetic field map is needed. The work had been started by A.U.Luccio^[4], but two main problems still remain on the optimization of the spin matrices and of the orbit matrices for the Snakes, they are:

- (1) Numerical spin matrices should be unitary,
- (2) Orbit matrices should be symplectic, essentially to preserve beam emittance, among other motion integration.

On the other hand, the results of magnetic field measurement for the prototype of a helical magnet was well agreed with those of three dimensional numerical calculation by TOSCA^[5]. It was encouraged to work further based on calculated magnetic field map, and to look for different methods to solve the problems mentioned above.

2. The methods for simulation

2.1 The equations of orbit motion and spin rotation

In Coordinate system of Fig.1, the Lorentz orbit motion equations are

$$\frac{ds}{ds} = 1 \quad (1)$$

$$\frac{dx}{ds} = x' = u \quad (2)$$

$$\frac{dy}{ds} = y' = v \quad (3)$$

$$\frac{du}{ds} = w \cdot [uvB_x - (1+u^2)B_y + vB_s] \quad (4)$$

$$\frac{dv}{ds} = w \cdot [(1+v^2)B_x - uvB_y - vB_s] \quad (5)$$




Fig. 1 Coordinate system

where $w = \frac{e \cdot (u^2 + v^2 + 1)^{\frac{1}{2}}}{m_0 \gamma \mathcal{V}}$, V is the velocity of charged particle. B_x , B_y , and B_s are the components of magnetic field in x , y and s direction respectively.

The equations of spin precession^[6] are

$$\frac{dS_x}{ds} = S_s P_y - S_y P_s \quad (6)$$

$$\frac{dS_y}{ds} = S_x P_s - S_s P_x \quad (7)$$

$$\frac{dS_s}{ds} = S_y P_x - S_x P_y \quad (8)$$

where

$$P_x = \frac{h}{B\rho} \cdot [(1+G\gamma)(-uB_s + (1+v^2)B_x - uvB_y) + (1+G)u(uB_x + B_s + vB_y)]$$

$$P_y = \frac{h}{B\rho} \cdot [(1+G\gamma)(-vB_s + (1+u^2)B_x - uvB_y) + (1+G)v(uB_x + B_s + vB_y)]$$

$$P_s = \frac{h}{B\rho} \cdot [(1+G\gamma)(-uB_x - vB_y + (u^2 + v^2)B_s) + (1+G)(uB_x + B_s + vB_y)]$$

$$h = \frac{1}{\sqrt{x'^2 + y'^2 + 1}} = \frac{1}{\sqrt{u^2 + v^2 + 1}}, \quad B\rho = \frac{m_0 \gamma \mathcal{V}}{e}.$$

Four-order Runge-Kutta method in Cartesian Coordinates was used to integrate the equations (1) - (8) in this simulation. In order to calculate the axis of spin precession, another three equations with the same form of (6)-(8) were used^[7]

$$\frac{dS_x^A}{ds} = S_s^A P_y - S_y^A P_s \quad (9)$$

$$\frac{dS_y^A}{ds} = S_x^A P_s - S_s^A P_x \quad (10)$$

$$\frac{dS_s^A}{ds} = S_y^A P_x - S_x^A P_y \quad (11)$$

The idea is the following:

If a spin vector \vec{S} precesses from \vec{S}_o to \vec{S}_f (see Fig. 2.1), the axis of precession must belong to the plane π bisecting the (\vec{S}_o, \vec{S}_f) angle. The axis is then perpendicular to the vector $\delta\vec{S} = \vec{S}_f - \vec{S}_o$. If we repeat the same argument for a second spin orientation \vec{S}^A , the axis must be also perpendicular to $\delta\vec{S}^A = \vec{S}_f^A - \vec{S}_o^A$. The axis of procession \vec{S} is therefore perpendicular both to $\delta\vec{S}$ and to $\delta\vec{S}^A$. Then

$$\vec{\sigma} = \delta\vec{S} \times \delta\vec{S}^A$$




Fig. 2.1 To find the axis of precession




Fig. 2.2 Angles of the axis of precession
φ, horizontal, θ, vertical

The angle are defined as (see Fig. 2.2)

$$\text{tag} \phi = \frac{\sigma_x}{\sigma_z}, \quad \text{tag} \theta = \frac{\sigma_y}{\sqrt{\sigma_x^2 + \sigma_z^2}}$$

2.2 The interpolation of magnetic field

The magnetic field distribution in three components B_x , B_y and B_s of a helical magnet at Cartesian grid $x = x_i$, $y = y_j$ and $s = s_k$ was given. The interpolation for this map was divided into two steps:

(1) Fitting on the (x, y) plane ($s = s_k$)

Bicubic spline function method was used to fit $B_x(x, y, s_k)$, $B_y(x, y, s_k)$, $B_s(x, y, s_k)$ ($k=1, 2, \dots, L$) and their derivatives in Cartesian Coordinates ,

$$B_x(x, y, s_k) = \sum_{k,l=1}^4 A_{xijkl}(x, y)(x - x_i)^{k-1}(y - y_j)^{l-1}$$

$$B_y(x, y, s_k) = \sum_{k,l=1}^4 A_{yijkl}(x, y)(x - x_i)^{k-1}(y - y_j)^{l-1}$$

$$B_s(x, y, s_k) = \sum_{k,l=1}^4 A_{sijkl}(x, y)(x - x_i)^{k-1}(y - y_j)^{l-1}$$

$$x_i \leq x \leq x_{i+1}, \quad y_j \leq y \leq y_{j+1}$$

$$(i=1, 2, \dots, m-1, j=1, 2, \dots, n-1)$$

Where the coefficients A_{ijkl} can be determined by one-dimensional cubic spline function, and the corresponding boundary conditions

$$\frac{\partial B_x(x, y, s_k)}{\partial x}(x = x_1, x_m; j = 1, 2, \dots, n)$$

$$\frac{\partial B_x(x, y, s_k)}{\partial y}(y = y_1, y_n; i = 1, 2, \dots, m)$$

$$\frac{\partial^2 B_x(x, y, s_k)}{\partial x \partial y}(x = x_1, x_m; y = y_1, y_n)$$

could be calculated by two-dimensional 3 - points Lagrangian function

$$B_x(x, y, s_k) = \sum_{i,j=1}^3 \left[\prod_{\substack{m,n=1 \\ m \neq i \\ n \neq j}}^3 \left(\frac{x - x_m}{x_i - x_m} \right) \left(\frac{y - y_n}{y_j - y_n} \right) \right] \bullet B_x(x_i, y_j, s_k)$$

(2) fitting in s - the third direction

Spline function method was used again to fit the coefficients on the plane (x, y) $A_{ijkl}(s_1)$, $A_{ijkl}(s_2), \dots, A_{ijkl}(s_l)$,

$$SA(s) = K_{0i} + K_{1i}(\Delta s) + K_{2i}(\Delta s)^2 + K_{3i}(\Delta s)^3$$

where

$$\Delta s = s - s_i$$

K_{0i} , K_{1i} , K_{2i} , K_{3i} could be calculated by one dimensional spline function fitting.

3. The construction of field distribution for full Snakes and Program SSTRAN

Two sets of magnetic field maps on Cartesian grid with 5 mm (named Map827.grid) and with

2.5 mm step (named Map827fine.grid) of half helical magnet length ($s=0 \sim L/2$) and half space ($y \geq 0, y=0 \sim 4$ cm) was provided by Dr. M. Okamura. The magnetic field on axis ($x=0, y=0$) is given in Fig 3. Then for the region of $y < 0$, the following symmetrical conditions are used:

$$B_x(x, -y, s) = B_x(x, y, s)$$

$$B_y(x, -y, s) = B_y(-x, y, s)$$

$$B_s(x, -y, s) = -B_s(-x, y, s)$$

For another half length ($s = -L/2 \sim 0$), the symmetrical conditions are

$$B_x(-x, y, -s) = -B_x(x, y, s)$$

$$B_y(-x, y, -s) = B_y(x, y, s)$$

$$B_s(-x, y, -s) = -B_s(x, y, s).$$

The field distribution of other three helical magnets takes as follows:

$$B_2 = -BCOEF * B_1 \quad (12)$$

$$B_3 = -B_2$$

$$B_4 = -B_1$$

11/09/97 22.48




Fig. 3 The magnetic field distribution of Map827.grid on axis ($x=0, y=0$)

BCOEF in equation (12) is the ratio of the magnetic field strength of the second helical magnet to that of the first one. The constructed field distribution based on Map827.grid on the axis ($x=0, y=0$) for full Sakes is given in Fig.4.

Program SSTRAN for integrating the equation (1)-(11) based on the calculation methods discussed above was written in FORTRAN.

4. Simulation Results

4.1 Maxwellian property checking

By using Map827.grid and Map827fine.grid, the interpolated magnetic field from Map827.grid was compared with that directly from Map827fine.grid, the difference between them is within 10^{-5} . Two magnetic field maps was checked to see how well they satisfy the Maxwell equations

$$\vec{\nabla} \cdot \vec{B} = \frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_s}{\partial s} = 0$$

$$\vec{\nabla} \times \vec{B} = \left(\frac{\partial B_s}{\partial y} - \frac{\partial B_y}{\partial s} \right) \vec{i} + \left(\frac{\partial B_x}{\partial s} - \frac{\partial B_s}{\partial x} \right) \vec{j} + \left(\frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \right) \vec{k} = 0$$

too. The first derivatives were directly calculated by the coefficients of spline function A_{ijkl} , the results were listed in table 1. It was seen from Table 1 that the scale B is well satisfied (10^{-6}) in the region around the axis, the error of curl B mainly comes from $(\vec{\nabla} \times \vec{B})_x$, it may be due to small value of B_s and rapid change with y .

Table 1. The results of Maxwellian property checking on Map827.grid

Point coordinates (m)	Magnetic field strength (T)	$\vec{\nabla} \cdot \vec{B}$ $\vec{\nabla} \times \vec{B}$ (T/m)
Fringing point: (0., 0., 0.02)	Bx=0.0698596 By=1.33392 Bs=0.000000145	$\vec{\nabla} \cdot \vec{B} = 4.5709596E - 6$ $(\vec{\nabla} \times \vec{B})_x = -6.304323E - 3$ $(\vec{\nabla} \times \vec{B})_y = 4.0081994E - 3$ $(\vec{\nabla} \times \vec{B})_s = -9.4958265E - 5$
inside point: (0. 0. 0.15)	Bx=0.511353 By=-1.23405 Bs=0.000000839	$\vec{\nabla} \cdot \vec{B} = 4.9906576E - 6$ $(\vec{\nabla} \times \vec{B})_x = -2.0883111E - 2$ $(\vec{\nabla} \times \vec{B})_y = -4.0178569E - 3$ $(\vec{\nabla} \times \vec{B})_s = -2.1633758E - 4$
point far from axis (-2.5, 2.5, 0.15)	Bx=0.51472201 By=-1.24078 Bs=-0.047527853	$\vec{\nabla} \cdot \vec{B} = 7.366029E - 3$ $(\vec{\nabla} \times \vec{B})_x = -2.2103902E - 2$ $(\vec{\nabla} \times \vec{B})_y = 2.6094199E - 3$ $(\vec{\nabla} \times \vec{B})_s = -4.9999588E - 2$

4.2 Optimized BCOEF

The ray with initial conditions of $(x_o, x'_o, y_o, y'_o) = (0.0, 0.0, 0.0, 0.0)$ and $(S_{ox}, S_{oy}, S_{os}) = (0, 1, 0)$ was traced through the full Snakes and optimized for BCOEF based on numerical calculated magnetic field of Map827.grid. BCOEF was adjusted automatically so as to get the final orbit angle x'_e, y'_e as small as possible and S_{ey} as closed to -1 as possible (spin direction turn over 180°). Take the fringing magnetic field into account, the magnetic field from 0 ~ 1.312 m (this is the length of Yoke) was taken from Map827.grid. The optimized BCOEF obtained is 3.09557299 when the amplitude of the field strength of the first magnet is 1.33626 T. The results of the final orbit and spin precess were listed in table 2 together with those based on Blewett-Chasman field formulas^[8] for comparison, the trajectory and spin procession in full snake were shown in Fig. 5 and Fig.6 respectively.

Table 2. The results of the trajectory with initial condition of $(x_o, x'_o, y_o, y'_o) = (0.0, 0.0, 0.0, 0.0)$ and $(S_{ox}, S_{oy}, S_{os}) = (0, 1, 0)$ when the field strength of the second magnet is at optimized value.

Filed Results	Map827.grid	Blewett-Chasman formula
B_{m1} (T)	1.336260	1.336260
B_{m2} (T)	4.136490	3.987906
x (mm)	0.24	-17.557
x' (mrad.)	0.01111	-1.58361
y (mm)	-0.46	2.0
y' (mrad.)	-0.08429	0.15085
Sx	0.01553612	-0.06402582
Sy	-0.99973410	-0.99512058
Ss	-0.01704195	0.07528736
$\phi(^{\circ})$	51.47374108	45.98502244
$\theta(^{\circ})$	-0.04410307	0.99667815

4.3 spin matrix

With the same initial orbit of $(x_o, x'_o, y_o, y'_o) = (0.0, 0.0, 0.0, 0.0)$, three representative rays with initial spin $\vec{S}_o = (1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$ respectively were traced, then spin matrix could be directly got by the final precessed spin. At the optimum value of BCOEF=3.09557299, that is $B_1=1.33626$ T, $B_2=4.136490$ T, the spin matrix




Fig. 4 The constructed field distribution of full Snakes on axis ($x=0$, $y=0$)

$$M_s = \begin{bmatrix} 0.22415978 & -0.01312770 & 0.97446405 \\ 0.01553612 & -0.99973410 & -0.01704195 \\ 0.97442862 & 0.01895950 & -0.22389618 \end{bmatrix}$$

The determinant of M_s : $\det[M_s] = 1.00000014777028$

$$|\det[M_s] - 1| = 1.4777028E-7$$

The dependence of the matrix elements on particle position and angle was investigated too. The calculation was repeated for different values of (x_o, x_o', y_o, y_o') . The spin process of some of the rays with large r ($r > 3.0\text{cm}$) rotate unregularly, but most of them turn out to be very close to each other, and the determinant of their matrices have the same order ($1.E-7$).




Fig. 5 The trajectory of the ray with $(x_o, x_o', y_o, y_o') = (0, 0, 0, 0)$




Fig. 6 The spin precession of the ray with $(S_{ox}, S_{oy}, S_{os}) = (0, 1, 0)$

4.4 orbit matrix

The calculation of orbit matrix M_T is more complicated than that of spin matrix. To calculate M_T , a certain number of rays, whose initial conditions were generated by random with Gaussian distribution in the components of (x_o, x'_o, y_o, y'_o) , were tracked. By solving a system of algebraic equations between the final and the initial coordinates of the same ray, a transformation expressed as a power expansion truncated to a pre established degree can be found. In this simulation, we are interested in a 0.th order 4×4 matrix. The problem has the same sense of general linear least squares, the solutions by use of the *normal equations* ^[9] was taken to get elements of the matrix .

It was found that the results largely depend on the number of the rays traced, The more the rays traced, the better the results.

After the number of the rays over 16, the matrix calculated is getting stable, the following is the result from 16 rays traced,

$$M_{or} = \begin{bmatrix} 0.99391892 & 10.82810815 & -0.01694592 & -0.03364342 \\ -0.00429513 & 0.96566983 & 0.00604201 & -0.00743476 \\ 0.00061034 & 0.16304712 & 0.98440460 & 10.67164840 \\ 0.00060338 & 0.06262435 & -0.00489109 & 0.94845089 \end{bmatrix}$$

The determinant of M_{or} : $\det[M_{or}] = 0.99508166E+00$

$$|\det[M_{or}]-1| = 4.91834490E-03$$

5. Conclusion

One set of methods have been setup for simulation in Siberian Snakes, and the results for numerical calculated magnetic field of Map827.grid is good, but further work should be done in next step.

References

- [1] K.Brown (BNL),et al. Conceptual Design for the Acceleration of Polarized Protons in RHIC. Spin Note AGS/RHIC/SN No.001.January 19,1996.
- [2] T. Katayama, Spin Dynamics in Synchrotron and Storage Ring, November,1995.
- [3] H.Wu, et al. Numerical and Analytic Studies on Siberian Snake. Spin Note AGS/RHIC/SN No. 025. April 29,1996.
- [4] A.U.Luccio, Field Map Generated Matrices for Spin Tracking. Spin Note AGS/RHIC/SN No.

041. November 5, 1996.

- [5] M .Okamura, The Hall Probe Measurement and Calculation of the Half Length Helical Dipole Magnet. Spin Note AGS/RHIC/SN No. 060, August 20, 1997.
- [6] V.Bargmann, Louis Michel and V.L.Telegdi, Precession of the Polarization of Particles Moving in a Homogeneous Electromagnetic Field. Physical Review Letters. Vol. 2, No. 10. May 15, 1959.
- [7] A.U.Luccio, Numerical Studies of Siberian Snakes and Spin Rotators for RHIC. Spin Note AGS/RHIC/SN No. 008. January 19, 1996.
- [8] J. P. Blewette and R. Chasman, Orbits and fields in the helical wiggler. Journal of Applied Physics, Vol.48, No. 7, July 1977.
- [9] W.H.Press et al. Numerical Recipes in FORTRAN, The Art of Scientific Computing Second Edition. Cambridge University Press, 1992.