

## Effect on Spin of Systematic Twist in RHIC Dipole Magnets

M. J. Syphers

September 1998

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Alternating Gradient Synchrotron Department  
Relativistic Heavy Ion Collider Project  
BROOKHAVEN NATIONAL LABORATORY  
Upton, New York 11973

*Spin Note*

AGS/RHIC/SN No. 076

**Effect on Spin of Systematic Twist  
In RHIC Dipole Magnets**

M. Syphers, A. Jain  
Brookhaven National Laboratory

September 10, 1998

# Effect on Spin of Systematic Twist in RHIC Dipole Magnets

M. Syphers, A. Jain

September 10, 1998

Magnetic measurements in some of the earlier RHIC dipole magnets indicated a twist of the dipole field along the body of the magnets. In one case in particular, a strong systematic twist through an angle of about 10 mrad over the length of the 9.5 m magnet was observed. While the dipole can be set so that the average field direction is vertical, thus self-canceling (or at least greatly reducing) the effects of a twisting field on the orbit, such a twisting field could in principle have an undesirable effect on the stable spin direction within the machine.

The average field direction was measured and noted in the magnet database for each magnet to be used for surveying purposes. This direction was determined from 10 field direction measurements along the length of the magnet. Recently, this data has been re-examined to look for systematic twist along the length of about 300 RHIC dipole magnets. For each of the magnets, the central 8 measurements in the straight section were used, and a straight line was fit to the field directions as a function of the axial position. The slope of this line is a measure of the twist in the magnet. This slope multiplied by the magnetic length, 9.45 m, is referred to as the "average twist" in the magnet. Fig. 1 shows the data and fitted line for a particular magnet in RHIC (one with a rather large average twist).

The output of the investigation is shown in Fig. 2. Some of the earlier magnets showed large twisting of the field, most likely generated by the welding process of the magnet cold mass shell. This process was subsequently altered, and the typical average twist angle through a magnet was kept below 2.2 mrad rms. The final distribution of twist angles is shown in Fig. 3. As can be seen, the distribution has a systematic twist error of 0.575 mrad.

## 1 Effects on Spin

Suppose that a magnet's field  $\vec{B}_0$  is systematically twisted along its length  $\ell$  through an angle  $\alpha$ . The field is oriented at an angle  $-\alpha/2$  at the entrance of the magnet, and  $+\alpha/2$  at the exit. We assume at this point that the effects on the orbit are

DRG516 (Seq. No. 46)




Figure 1: Field direction, in mrad from vertical, versus axial location along the length of RHIC dipole DRG516. The average twist of the field is 9.4 mrad.

---

DRG(8) Dipoles -- Twist over 9.45 m (Warm)




Figure 2: Average twist angle over the length of RHIC dipole magnets versus dipole magnet number. Larger twists were observed in some of the earlier magnets, which was kept better under control for the remainder of the production.

### Twist in RHIC DRG/DR8 Dipoles




Figure 3: Distribution of average twist angles for 296 RHIC dipole magnets. The mean of the distribution is 0.58 mrad, with a standard deviation of 2.2 mrad.

---

minimal. (See Section 3.) The magnet acts as a helical dipole field of length  $\ell$  and pitch  $k = \alpha/\ell$ . The spin vector  $\vec{S}$  is then transformed according to

$$\begin{pmatrix} S_x \\ S_y \\ S_z \end{pmatrix}_{out} = M_\alpha M_{dpl} M_\alpha \begin{pmatrix} S_x \\ S_y \\ S_z \end{pmatrix}_{in} \quad (1)$$

where the matrix for the “helical” field is given by [1]

$$\begin{aligned} M_{dpl} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \phi & \sin \phi \\ 0 & -\sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \cos \mu & -\sin \mu & 0 \\ \sin \mu & \cos \mu & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \phi & -\sin \phi \\ 0 & \sin \phi & \cos \phi \end{pmatrix} \\ &= \begin{pmatrix} \cos \mu & -\cos \phi \sin \mu & \sin \phi \sin \mu \\ \cos \phi \sin \mu & \cos^2 \phi \cos \mu + \sin^2 \phi & \cos \phi \sin \phi (1 - \cos \mu) \\ -\sin \phi \sin \mu & \cos \phi \sin \phi (1 - \cos \mu) & \sin^2 \phi \cos \mu + \cos^2 \phi \end{pmatrix}, \end{aligned}$$

and where

$$M_\alpha = \begin{pmatrix} \cos(\alpha/2) & \sin(\alpha/2) & 0 \\ -\sin(\alpha/2) & \cos(\alpha/2) & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (2)$$

$$\mu = -k\ell \sqrt{1 + \left(\frac{\kappa}{k}\right)^2}, \quad (3)$$

$$\phi = \tan^{-1}(\kappa/k), \quad (4)$$

$$\kappa \equiv (1 + G\gamma)B_0/(B\rho). \quad (5)$$

## 2 Estimates for RHIC

Ideally, if  $\alpha$  were zero for each RHIC dipole, then the total spin matrix for RHIC would just be the matrix for a rotation of  $2\pi(1 + G\gamma)$  about the vertical axis. For RHIC high energy (250 GeV) and a field of 3.45 T, this rotation angle modulo  $2\pi$  is  $137.53^\circ$ . If each dipole in RHIC has a systematic twist of  $\alpha = 0.58$  mrad, then subsequent matrix for RHIC is altered, and the resulting spin rotation angle is  $137.54^\circ$ . The change in the spin tune of RHIC is thus only  $2 \times 10^{-5}$ .

The above calculation was in the absence of Siberian Snakes. With the two Snakes in RHIC, the spin tune is altered from 0.5 by only about  $2 \times 10^{-6}$  due to the systematic twist error.

Since the twist angle has a distribution with rms 2.2 mrad, an ensemble of 500 RHIC rings was looked at, each ring containing dipoles generated randomly from a Gaussian distribution of twist errors with mean 0.575 mrad and rms 2.2 mrad. For each “ring” the difference in the spin tune from 0.5 was computed. The results are






Figure 4: Distribution of spin tune shift (from the ideal value of 0.5) for an ensemble of 500 “RHIC rings.” The dotted curve is the normalized integrated sum.

shown in Fig. 4. Here, we see that the chances of the spin tune shift due to this effect being greater than about  $7 \times 10^{-5}$  is only 5%.

The twisting of the dipole field in the RHIC dipole magnets should not cause a significant effect on the proton spin at all.

### 3 Effects on Orbit

A twisting of the field through a dipole magnet in the RHIC lattice will contribute to the closed orbit distortion around the accelerator and must be corrected with the dipole corrector system. A changing orientation of the main dipole field, to lowest order in the twist angle, will cause no distortion of the vertical field component, however horizontal field errors will be generated. For a helical twist with pitch  $\alpha$ , where the field begins and ends with angles  $\pm\alpha/2$  to the vertical, the horizontal field component will be

$$B_x(s) = -B_0 \sin(ks - \alpha/2)$$

along the length  $\ell$  of the magnet. Thus, the vertical equation of motion will be

$$y'' = \frac{B_x}{B\rho} = -\frac{B_0}{B\rho} \sin(ks - \alpha/2)$$

and upon passing through the twisted field of the magnet, the resulting vertical displacement and slope is given by

$$y(\ell) = y_0 + y'_0 \ell - \frac{B_0 \ell}{k(B\rho)} \cos(\alpha/2) + 2 \frac{B_0}{k^2(B\rho)} \sin(\alpha/2) \quad (6)$$

$$y'(\ell) = y'_0 \quad (7)$$

where  $y_0$  and  $y'_0$  are the initial positions and slope coming into the magnet.

As can be seen, the action of the twisted dipole is equivalent to passing through a drift of length  $\ell$  and subsequently being given a step in the vertical direction of displacement

$$\Delta y = \frac{1}{12} \theta_0 \alpha \ell$$

where here,  $\theta_0$  is the horizontal bending angle of the dipole magnet and  $\alpha$  is considered to be small.

A “step” in the particle trajectory generates a closed orbit distortion which is given by

$$y(s) = \sqrt{\frac{\beta(s)}{\beta_0}} \frac{\Delta y}{2 \sin \pi \nu} [\alpha_0 \cos(\psi(s) - \pi \nu) - \sin(\psi(s) - \pi \nu)]$$

with  $\alpha$  and  $\beta$  being the Courant-Snyder parameters of the lattice, the subscript “0” designating the location at the end of the helical field.

For a distribution of  $N$  errors around the accelerator, the expected rms closed orbit can then be estimated by

$$\hat{y}_{rms} = \sqrt{\hat{\beta}} \frac{\Delta y_{rms}}{2|\sin \pi\nu|} \sqrt{\langle \frac{\alpha^2}{\beta} \rangle + \langle \frac{1}{\beta} \rangle} \sqrt{\frac{N}{2}}.$$

Noting that the combination  $(1 + \alpha^2)/\beta \equiv \gamma$  (another Courant-Snyder parameter), the expression becomes

$$\hat{y}_{rms} = \sqrt{\hat{\beta}\langle\gamma\rangle} \frac{\Delta y_{rms}}{2\sqrt{2}|\sin \pi\nu|} \sqrt{\frac{N}{2}}$$

For our final expression, we use the fact that  $\gamma$  is a constant within all the arc dipoles in the FODO cells, and so  $\hat{\beta}\langle\gamma\rangle = 1 + \hat{\alpha}^2 = 2/[1 - \sin(\mu/2)]$ , where  $\mu$  is the cell phase advance. Thus, we have finally

$$\hat{y}_{rms} = \frac{\Delta y_{rms}}{2|\sin \pi\nu|\sqrt{1 - \sin(\mu/2)}} \sqrt{N}.$$

We can now put in some numbers. For the RHIC distribution of twist angles  $\alpha$ , we get

$$\Delta y_{rms} = \frac{1}{12} \theta_0 \ell \alpha_{rms} = \frac{1}{12} (40 \text{ mrad})(9.5 \text{ m})(2.2 \text{ mrad}) = 0.07 \text{ mm}.$$

And, assuming approximately 160 RHIC dipole magnets per ring,

$$\hat{y}_{rms} \approx 1.5 \text{ mm}.$$

Thus, the orbit distortion due to the twist is much smaller than the orbit distortion due to the quadrupole alignment errors, which generate uncorrected rms orbit excursions of 10-15 mm or more.

## References

- [1] M. J. Syphers, "Spin Motion through Helical Dipole Magnets," BNL Internal Report AGS/RHIC/SN-020, February, 1996.