

## Alternate AGS - Booster lattices

Z. Parsa

May 1986

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ALTERNATE AGS - BOOSTER LATTICES

*Booster Technical Note*  
No. 32

Z. PARSA AND S. TEPIKIAN

MAY 7, 1986

ACCELERATOR DEVELOPMENT DEPARTMENT  
*Brookhaven National Laboratory*  
Upton, N.Y. 11973

## ABSTRACT

WE HAVE STUDIED THE ALTERNATE CONCEPTUAL LATTICES FOR THE AGS - BOOSTER. WE PRESENT SOME OF OUR RESULTS FOR THE STANDARD BOOSTER LATTICE AND THE ALTERNATE [1/4 AGS, P=12] COMBINED, HYBRID AND [1/3 AGS, P = 8] SEPARATED FUNCTION LATTICES. WE INCLUDE OUR CALCULATION AND ANALYSIS OF THE STOP BANDWIDTHS FOR THE THIRD ORDER RESONANCES OF THESE LATTICES.

## INTRODUCTION:

-----

We have studied the nonlinear effects of the AGS-Booster lattice [1,2] and its alternates. In our investigation of the nonlinear effects for the present Booster lattice in addition to the third order structure resonances given in Section V, we have calculated the fourth order resonances given elsewhere [3,4]. These resonances are crossed at injection because of the space charge effect since the Laslett tune shift may be as large as 1. This has led to our investigation of the following [1/4 AGS, P = 12] combined, hybrid and [1/3 AGS, P = 8] separated function lattices given in Figs. 1-3. In section two we discuss the general treatment of the nonlinear effects following Guignard [5] and Donald [6]. In Sections three and four using program HARMON (MAD403) the nonlinear effects of these alternative lattices are analyzed and given in tables I-III. With space charge tune shifts as large as 1 the third order resonances may be crossed.

The sixth order resonances are important and we will discuss them in subsequent papers using program NONLIN [4] (since HARMON is not equipped to do so). We have used  $B'' = .24$  T/m for the eddy current sextupoles.

## SECTION II

-----

To study the dynamics of an accelerator with chromaticity correcting sextupoles and eddy current sextupoles, we have used the following Hamiltonian [1]:

$$H_0 = \frac{1}{2} (p_x^2 + p_y^2) + \left( \frac{1}{\rho^2} - K(s) \right) \frac{x^2}{2} + K(s) \frac{y^2}{2} \quad (1)$$

$$+ \frac{S(s)}{6} (x^3 - 3xy^2)$$

where,  $x$  and  $y$  are the positions,  $p_x$  and  $p_y$  are the conjugate momentum,  $K$  is the quadrupole focussing strength,  $S$  is the sextupole strength, and  $s$  is the distance along the orbit (the "time" variable for the Hamiltonian).

First we transform the Hamiltonian to action - angle form:

$$H_1 = \frac{2\pi}{C} (Q_x J_x + Q_y J_y) + V(J_x, J_y, \phi_x, \phi_y, s) \quad (2)$$

where, C is the circumference; Jx, Jy are the action variables (directly related to the beam emittances);  $\phi_x$ ,  $\phi_y$  are the angle variables and Qx, Qy are the betatron tunes.

When the system is operating near a resonance, perturbation theory no longer holds. Although, an approximate invariant can be found for system near a single resonance if the contribution from the other resonances are small. This can lead to the Hamiltonian

$$H_2 = \frac{2\pi}{C} e K_x + \sum_{\ell=1}^{\ell} W_{\ell} K_x^{\frac{\ell}{2}+1} + A \cos \psi_x \quad (3)$$

where, e is the bandwidth given in eq. (4); Kx and Ky are the new action variables and  $\psi_x$  is the transformed angle.  $W_{\ell}$  are the stabilizing coefficients and A is the resonance strength. This Hamiltonian is an invariant of the motion.

$$e = N_x Q_x + N_y Q_y - p \quad (4)$$

where the integers  $N_x$ ,  $N_y$  and p define a given resonance.

Following Guignard [5], dynamic properties (such as stop bandwidths,  $\Delta e$ ) can be found from invariant of the motion  $H_2$ . The stop bandwidths ( $\Delta e$ ) are defined to be the smallest bandwidth such that the action in Eq. (3) is still bounded. This can be done by first considering the fixed points of Eq. (3) which are defined to be the points at which there is no motion. These fixed points are the solutions to the following equations (note that Ky is also an invariant of the motion and can be treated as a constant).

$$\begin{aligned} 0 &= \frac{d\psi_x}{ds} = \frac{\partial H_2}{\partial K_x} \\ &= \frac{2\pi}{C} e + \sum_{\ell=1}^{\ell} \left( \frac{\ell}{2} + 1 \right) W_{\ell} K_x^{\frac{\ell}{2}} + \frac{\partial A}{\partial K_x} \cos \psi_x \end{aligned} \quad (5a)$$

$$0 = \frac{d K_x}{d s} = - \frac{\partial H_2}{\partial \psi_x} = A \sin \psi_x \quad (5b)$$

There are many solutions [ $\psi_x = n \pi$ ] to eq. (5b) which lead to two cases [ $\cos(\psi_x) = \pm 1$ ] in eq. (5a). We note that for the smallest positive value of  $K_x$  there is at most one solution for each of these cases corresponding to stable and /or unstable fixed point(s). The nature of these solutions are determined by the bandwidth ( $e$ ) and the stabilizing coefficients ( $W_\ell$ ).

We find the stop bandwidths for these two cases by substituting  $\cos(\psi_x) = \pm 1$  into eq. (3) and obtaining the following two new equations:

$$F_{\pm} = \frac{2\pi}{C} e K_x + \sum_{\ell=1}^{\ell} W_\ell K_x^{\frac{\ell}{2} + 1} \pm A \quad (6)$$

For the given initial conditions, four equations can be deduced from Eq. (6) from which the stop bandwidths and the extrem values of  $K_x$  are found. These equations are listed below:

$$\begin{aligned} 0 &= F_+(K_{x0}) - F_+(K_x) \\ 0 &= F_+(K_{x0}) - F_-(K_x) \\ 0 &= F_-(K_{x0}) - F_+(K_x) \\ 0 &= F_-(K_{x0}) - F_-(K_x) \end{aligned} \quad (7)$$

For  $e < \Delta e$  Eqs. (7) can be satisfied for any value of action  $K$ , where  $\Delta e$  is the stop bandwidth.

We also note that, Eqs. (8) [5] calculates the necessary distance ( $\delta e$ ) between the operating tunes ( $Q_x, Q_y$ ) and the resonance line which must be kept to avoid or limit the relative growth of amplitude or beating of a single particle in a given interval  $\Lambda = [(K_x/K_{x0})^{1/2} - 1]$ . Where for a sum resonance

$$\delta e \geq \frac{\Delta e}{2} \left( 1 + \frac{1}{\Lambda} \frac{\frac{n_x}{2} E_y + \frac{n_y}{2} E_x}{\frac{n_x}{2} E_y + \frac{n_y}{2} E_x} \right) \quad (8a)$$

and for difference resonance

$$\delta e \geq \frac{\Delta e}{2} \frac{1}{\Lambda} \quad (8b)$$

$N_x, N_y$  are the integers (see eq. (4)) defining a given resonance and  $E_x$  and  $E_y$  are the emittances of the beam at injection [directly related to  $K_x$  and  $K_y$ ].

In the following sections we discuss our results including the stop bandwidths (defined above) calculated for the alternate lattices using program HARMON (MAD403 [6]).

### SECTION III

-----

In this section we consider the 1/4 AGS combined function and 1/4 AGS hybrid lattices shown in Fig. 1 and 2 respectively. Both lattices are very similar with operating tunes at  $Q_x = 4.82, Q_y = 4.83$  and a periodicity of 12. If the space charge tune shift can be as large as 1 then we must worry about the third order resonances  $3Q_x = 12$  and  $Q_x + 2Q_y = 12$ .

The results of running MAD403 with HARMON are shown in Tables I and II. Table IA gives the orbit parameters for the combined function lattice at the operating tunes, while Table IB gives the orbit parameters at the tunes  $Q_x = 4.01$  and  $Q_y = 4.11$  near the resonances. The stop bandwidths at the tunes near resonances are given in Table IC. Note  $e$  can be as large as about 0.55 which means the tune shift due to the space charge should be less than about 0.73 from the operating tune.

Tables IIA-C gives similar information on the hybrid lattice. It can be seen in Table IIC that the stop bandwidths are larger for the hybrid lattice ( $\Delta e = 0.65$ ) than the combined function lattice ( $\Delta e = 0.55$ ), but we still can shift the tune as long as the upper limit of tune shift is less than 0.71 (slightly smaller than the above lattice).

### SECTION IV

-----

The third lattice we have studied is the 1/3 AGS separated function design with a periodicity of 8 as shown in Fig. 3. If this machine is run at the operating tunes of  $Q_x = 6.82$  and  $Q_y = 6.83$ , then no resonances of up to fourth order are crossed due to space charge tune shifts. The orbit parameters for this



case are given in Table IIIA.

If this machine is to run at an alternate set of operating tunes of  $Q_x = 5.82$  and  $Q_y = 5.83$  then two third order resonances can be crossed at  $3Q_x = 16$  and  $Q_x + 2Q_y = 16$ . The orbit parameters for this case are shown in Table IIIB. Additionally, the orbit parameters at the tunes near resonances are given in Table IIIC. Table IIID gives the stop bandwidths which are of the order of 0.007. These stop bandwidths are small and it is expected that these resonances are passable (although, they may require some tuning out).

## SECTION V

-----

The standard AGS - Booster lattice is limited by third order resonances as were the alternate lattices described in the previous sections. If space charge can shift the operating tune of this machine near to the tune of 4 then the third order resonances become noticeable. For tunes near the third order resonances, (for instance,  $Q_x = 4.01$  and  $Q_y = 4.11$ ), the orbit parameters are calculated and are given in Table IVA. The stop bandwidths are given in Table IVB which are found to be of the order of 0.076. This suggests that we should limit the space charge tune shifts to within the order of 0.78 before the third order resonance effects becomes important.

## VI. CONCLUSION

-----

Due to the space charge tune shifts at injection, resonances that could cause the beam to blow up in an accelerator can be crossed. These resonances in the standard AGS-Booster lattice and its alternates were investigated. Considering the third order resonances, it is found that the stop bandwidths of the 1/4 AGS Combined function and the 1/4 AGS Hybrid lattices have very strong third order resonances limiting the maximum space charge tune shifts that could be handled before there are adverse effects to the beam dynamics. In the standard AGS-Booster, the stop bandwidths for the third order resonances are strong, but we can accommodate space charge tune shifts as large as 0.78, if the fourth order resonances (calculated elsewhere [3,4]) are passable. Finally, the 1/3 AGS separated function lattice at the normal operating tunes does not cross any resonance up to fourth order. However, if

this lattice is run at an alternate set of tunes then the third order resonances could be crossed. We found that the stop bandwidths for this resonance are small and these resonances are passable (although, tuning them out may be necessary).

The sixth order resonances are important and will be discussed elsewhere, since HARMON is limited to the calculation of fourth order resonances.

#### REFERENCES:

-----

1. E. Courant and Z. Parsa, Booster Lattice with Enlarged Q5 and 1,2,4,7 Sextupole Configuration. Booster Tech. Note No.26, (April 21,1986).
2. Z. Parsa, Booster Parameters with Enlarged Q5, Booster Tech. Note No. 25, (April 17, 1986).
3. Z. Parsa, S. Tepikian, Analysis of Resonances in the AGS - Booster, Booster Tech. Note No. 34 (May 1986)
4. Z. Parsa, S. Tepikian and E. Courant, Fourth Order Resonances in the AGS - Booster Lattice, Booster Tech. Note No. 35
5. M. Donald, D. Schofield, a Users Guide to the Harmon Program, LEP Note 420 (1982); M. Donald private communication (May 1986); using [PARSA1.MAD]MAD403.EXE.
6. G. Guignard, General treatment of Resonances in Accelerators, Cern 78-11 (Nov. 10, 1978)

\*\*

[In the following tables COS, SIN and MODULUS (A in Eq.(3)) are the resonance strengths, and DE(S), DQ(S) and DQ20(S) are the stop bandwidths. Where "S" means Systemmatic and "R" means Random].

TABLE IA

ALTERNATE AGS-BOOSTER LATTICE WITH CHROMATICITY SEXTUPOLES  
AND EDDY CURRENTS  
AT OPERATING TUNE ( $Q_x = 4.82$ ,  $Q_y = 4.83$ )

BOOSTER LATTICE OF COMBINED FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH =	201.780000	NSUP =	12
QX =	4.820000	QY =	4.830000
QX' =	0.000000	QY' =	-0.000003
ALFA =	0.509860E-01	GAMMA(TR) =	4.428684
BETAX =	1.15267E+01	BETAY =	1.15697E+01
		ETAX =	1.56754E+00
BETAX(MAX) =	15.591579	BETAY(MAX) =	15.607346
DX(MAX) =	2.239726	DY(MAX) =	0.000000

-2

NORMALIZED STRENGTHS [m<sup>-2</sup>]  
ID STRENGTH

SFC	-2.76326E-01
SDC	-1.02324E+00
SXV1	4.46595E-02
SXV2	5.58243E-02
SXV3	7.81541E-02
SXV4	3.34946E-02

Q SHIFT EFFECTS [PERTURBATION OF TUNES]

	G22000	DQXDEX	DQX
-1.86911E-01	1.60308E+00	-3.73823E-01	-2.55321E-05
	G00220	DQYDEY	DQY
2.08373E+01	2.19891E+00	4.16746E+01	2.84637E-03
	G11110	DQXDEY	DQYDEX
-2.35393E+01	2.93188E+00	-2.35393E+01	-2.35393E+01
		DQX	DQY
		-1.60773E-03	-1.60773E-03

TABLE IB  
-----

ALTERNATE AGS-BOOSTER LATTICE WITH CHROMATICITY SEXTUPOLES  
AND EDDY CURRENTS  
AT TUNE NEAR RESONANCE (QX = 4.01, QY = 4.11)

BOOSTER LATTICE OF COMBINED FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH =	201.780000	NSUP =	12
QX =	4.010119	QY =	4.110112
QX' =	0.671394	QY' =	3.951067
ALFA =	0.689755E-01	GAMMA (TR) =	3.807611
BETAX =	1.04933E+01	BETAY =	1.04988E+01
		ETAX =	2.15115E+00
BETAX (MAX) =	14.547907	BETAY (MAX) =	14.295551
DX (MAX) =	2.873679	DY (MAX) =	0.000000

NORMALIZED STRENGTHS  
ID STRENGTH

SFC	-2.76326E-01
SDC	-1.02324E+00
SXV1	4.46595E-02
SXV2	5.58243E-02
SXV3	7.81541E-02
SXV4	3.34946E-02

Q SHIFT EFFECTS

	G22000	DQXDEX	DQX
-5.08782E+02	1.34757E+00	-1.01756E+03	-6.94996E-02
	G00220	DQYDEY	DQY
-1.71354E+02	1.88448E+00	-3.42708E+02	-2.34069E-02
	G11110	DQXDEY	DQYDEX
-7.64174E+02	2.51264E+00	-7.64174E+02	-7.64174E+02
		DQX	DQY
		-5.21931E-02	-5.21931E-02

TABLE IC

ALTERNATE BOOSTER LATTICE WITH [COMBINED FUNCTION ELEMENTS]  
 WITH CHROMATICITY SEXTUPOLES AND EDDY CURRENTS  
 DELTA(P)/P= 0.000000 [EX0 =6.8300E-05 EY0 =6.8300E-05]

THIRD ORDER EFFECTS OF SEXTUPOLES [ RESONANCE EFFECTS]  
 3Qx = 12

COSINE	SINE	MODULUS	RANDOM	DE(S)
1.2896E+00	2.7465E-01	1.3185E+00	9.2934E-01	1.9615E-01
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
1.3825E-01	6.5382E-02	4.6083E-02	1.3512E-01	9.5237E-02

Qx+2Qy=12

COSINE	SINE	MODULUS	RANDOM	DE(S)
3.2830E+00	5.8184E+00	6.6807E+00	4.5129E+00	5.5212E-01
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
3.7296E-01	2.4691E-01	1.6679E-01	5.1029E-01	3.4471E-01

TABLE IIA

ALTERNATE AGS-BOOSTER LATTICE WITH CHROMATICITY SEXTUPOLES  
AND EDDY CURRENTS  
AT OPERATING TUNE ( $Q_x = 4.82$ ,  $Q_y = 4.83$ )

hybrid BOOSTER LATTICE OF COMBINED FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH	=	201.780000	NSUP	=	12
QX	=	4.820000	QY	=	4.830000
QX'	=	0.000054	QY'	=	0.000031
ALFA	=	0.478141E-01	GAMMA (TR)	=	4.573220
BETAX	=	1.00758E+01	BETAY	=	9.76910E+00
			ETAX	=	1.43651E+00
BETAX (MAX)	=	14.058073	BETAY (MAX)	=	13.935099
DX (MAX)	=	2.306863	DY (MAX)	=	0.000000

NORMALIZED STRENGTHS  
ID STRENGTH

SFC	-1.89019E-01
SDC	-9.73867E-01
SXV1	8.34574E-02
SXV2	6.72683E-02
SXV3	3.76814E-02

Q SHIFT EFFECTS

	G22000	DQXDEX	DQX
-5.77230E+00	1.39254E+00	-1.15446E+01	-7.88496E-04
	G00220	DQYDEY	DQY
7.90935E+00	1.81900E+00	1.58187E+01	1.08042E-03
	G11110	DQXDEX	DQYDEX
-5.80613E+01	2.42533E+00	-5.80613E+01	-5.80613E+01
		DQX	DQY
		-3.96559E-03	-3.96559E-03

TABLE IIB  
-----

ALTERNATE AGS-BOOSTER LATTICE (1/4 AGS) WITH CHROMATICITY  
AND EDDY CURRENT SEXTUPOLES  
AT TUNES NEAR RESONANCE ( $Q_x = 4.01$ ,  $Q_y = 4.11$ )

hybrid BOOSTER LATTICE OF COMBINED FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH	=	201.780000	NSUP	=	12
QX	=	4.009997	QY	=	4.109997
QX'	=	1.290876	QY'	=	3.696270
ALFA	=	0.665890E-01	GAMMA(TR)	=	3.875241
BETAX	=	9.90474E+00	BETAY	=	9.63097E+00
			ETAX	=	2.05177E+00
BETAX(MAX)	=	13.715713	BETAY(MAX)	=	13.673790
DX(MAX)	=	2.937144	DY(MAX)	=	0.000000

NORMALIZED STRENGTHS  
ID            STRENGTH

SFC	-1.89019E-01
SDC	-9.73867E-01
SXV1	8.34574E-02
SXV2	6.72683E-02
SXV3	3.76814E-02

Q SHIFT EFFECTS

	G22000		DQXDEX		DQX
	-----		-----		-----
-8.26976E+02	1.48420E+00	-1.65395E+03		-1.12965E-01	
	G00220		DQYDEY		DQY
	-----		-----		-----
-2.45315E+02	1.99024E+00	-4.90630E+02		-3.35100E-02	
	G11110		DQXDEY		DQYDEX
	-----		-----		-----
-1.06011E+03	2.65366E+00	-1.06011E+03		-1.06011E+03	
			DQX		DQY
			-----		-----
			-7.24052E-02		-7.24052E-02

TABLE IIC

ALTERNATE hybrid BOOSTER LATTICE WITH [COMBINED FUNCTION  
ELEMENTS] WITH CHROMATICITY SEXTUPOLES AND EDDY CURRENTS  
DELTA(P)/P= 0.000000 [EX0 =6.8300E-05 EY0 =6.8300E-05]

THIRD ORDER EFFECTS OF SEXTUPOLES [ RESONANCE EFFECTS ]

3Qx = 12

COSINE	SINE	MODULUS	RANDOM	DE(S)
1.6394E+00	2.8678E-01	1.6643E+00	7.2788E-01	2.4758E-01
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
1.0828E-01	8.2526E-02	3.6093E-02	1.7055E-01	7.4592E-02

Qx+2Qy=12

COSINE	SINE	MODULUS	RANDOM	DE(S)
5.4062E+00	5.6395E+00	7.8122E+00	4.0555E+00	6.4563E-01
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
3.3516E-01	2.8874E-01	1.4989E-01	5.9672E-01	3.0977E-01



TABLE IIIA

ALTERNATE AGS-BOOSTER LATTICE (1/3 AGS ) WITH CHROMATICITY  
AND EDDY CURRENT SEXTUPOLES  
AT OPERATING TUNES (Qx = 6.82, Qy = 6.83)

BOOSTER LATTICE WITH SEP. FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH	=	269.040000	NSUP	=	8
QX	=	6.820000	QY	=	6.829999
QX'	=	0.000000	QY'	=	0.000000
ALFA	=	0.221530E-01	GAMMA (TR)	=	6.718681
BETAX	=	3.33250E+00	BETAY	=	1.36827E+01
			ETAX	=	6.38422E-01
BETAX (MAX)	=	14.132757	BETAY (MAX)	=	13.682687
DX (MAX)	=	1.937133	DY (MAX)	=	0.000000

NORMALIZED STRENGTHS  
ID            STRENGTH

SXV	6.69892E-02
SFCH	3.71406E-01
SDCH	-9.85768E-01

Q SHIFT EFFECTS

	G22000		DQXDEX		DQX
	-----		-----		-----
5.41842E+01	7.57342E-01	1.08368E+02		7.40156E-03	
	G00220		DQYDEY		DQY
	-----		-----		-----
1.19016E+02	9.16056E-01	2.38032E+02		1.62576E-02	
	G11110		DQXDEY		DQYDEX
	-----		-----		-----
-7.94214E+01	1.22141E+00	-7.94214E+01		-7.94214E+01	
			DQX		DQY
			-----		-----
		-5.42448E-03		-5.42448E-03	

TABLE IIIB  
-----

ALTERNATE AGS-BOOSTER LATTICE (1/3 AGS ) WITH CHROMATICITY  
AND EDDY CURRENT SEXTUPOLES  
AT ALTERNATE OPERATING TUNES (Qx = 5.82, Qy = 5.83)

BOOSTER LATTICE WITH SEP. FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH	=	269.040000	NSUP	=	8
QX	=	5.819837	QY	=	5.830096
QX'	=	0.000631	QY'	=	0.000067
ALFA	=	0.307175E-01	GAMMA(TR)	=	5.705678
BETAX	=	4.29045E+00	BETAY	=	1.39083E+01
			ETAX	=	1.02027E+00
BETAX(MAX)	=	14.173987	BETAY(MAX)	=	13.908285
DX(MAX)	=	2.015216	DY(MAX)	=	0.000000

NORMALIZED STRENGTHS	
ID	STRENGTH
-----	-----
SXV	6.69892E-02
SFCH	9.83404E-02
SDCH	-8.31124E-01

Q SHIFT EFFECTS

	G22000		DQXDEX		DQX
	-----		-----		-----
6.63888E+00	1.03313E+00	1.32778E+01		9.06870E-04	
	G00220		DQYDEY		DQY
	-----		-----		-----
5.07415E+01	1.31499E+00	1.01483E+02		6.93129E-03	
	G11110		DQXDEY		DQYDEX
	-----		-----		-----
-2.41117E+01	1.75332E+00	-2.41117E+01		-2.41117E+01	
			DQX		DQY
			-----		-----
		-1.64683E-03		-1.64683E-03	

TABLE IIIC

ALTERNATE AGS-BOOSTER LATTICE (1/3 AGS ) WITH CHROMATICITY  
AND EDDY CURRENT SEXTUPOLES  
AT TUNES NEAR RESONANCE ( $Q_x = 5.34$ ,  $Q_y = 5.35$ )

BOOSTER LATTICE WITH SEP. FUNCTION ELEMENTS  
DELTA(P)/P = 0.000000

TOTAL LENGTH	=	269.040000	NSUP	=	8
QX	=	5.339892	QY	=	5.349842
QX'	=	1.749702	QY'	=	1.046964
ALFA	=	0.364175E-01	BETAX(MAX)	=	14.497906
BETAX	=	4.89019E+00	BETAY	=	1.42118E+01
			ETAX	=	1.24746E+00
BETAY(MAX)	=	14.211846	GAMMA(TR)	=	5.240162
DX(MAX)	=	2.219571	DY(MAX)	=	0.000000

## NORMALIZED STRENGTHS

ID	STRENGTH
SXV	6.69892E-02
SFCH	9.83404E-02
SDCH	-8.31124E-01

## Q SHIFT EFFECTS

	G22000		DQXDEX		DQX
	-----		-----		-----
	2.53192E+01	1.63113E+00	5.06385E+01		3.45861E-03
	G11110		DQXDEY		DQYDEX
	-----		-----		-----
	-1.16873E+02	2.17484E+00	-1.16873E+02		-1.16873E+02
			DQX		DQY
			-----		-----
			-7.98239E-03		-7.98239E-03

TABLE IIIID  
-----

ALTERNATE BOOSTER LATTICE (1/3 AGS) WITH [SEP. FUNCTION  
ELEMENTS] CHROMATICITY SEXTUPOLES AND EDDY CURRENTS  
DELTA(P)/P= 0.000000 [EX0 =6.8300E-05 EY0 =6.8300E-05]

THIRD ORDER EFFECTS OF SEXTUPOLES [ RESONANCE EFFECTS ]

3Qx = 16

COSINE	SINE	MODULUS	RANDOM	DE(S)
1.5991E-01	-3.9287E-01	4.2417E-01	7.1560E-01	6.3099E-02
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
1.0645E-01	2.1033E-02	3.5484E-02	4.3468E-02	7.3333E-02

Qx+2Qy=16

COSINE	SINE	MODULUS	RANDOM	DE(S)
6.5096E-01	-6.5569E-01	9.2395E-01	4.1137E+00	7.6358E-02
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
3.3997E-01	3.4149E-02	1.5204E-01	7.0574E-02	3.1421E-01

TABLE IVA

AGS BOOSTER LATTICE WITH CHROMATICITY SEXTUPOLES  
AND EDDY CURRENTS  
AT TUNES NEAR RESONANCE ( $Q_x = 4.01$ ,  $Q_y = 4.11$ )

## SEPARATED FUNCTION AGS-BOOSTER LATTICE

DELTA(P)/P = 0.000000

TOTAL LENGTH	=	201.780000	NSUP	=	6
QX	=	4.010087	QY	=	4.110047
QX'	=	4.380868	QY'	=	0.364414
ALFA	=	0.631572E-01	GAMMA (TR)	=	3.979134
BETAX (MAX)	=	14.766930	BETAY (MAX)	=	14.019838
BETAX	=	4.62840E+00	BETAY	=	1.39299E+01
			ETAX	=	1.31580E+00
DX (MAX)	=	3.169083	DY (MAX)	=	0.000000

NORMALIZED STRENGTHS  
ID STRENGTH

SXF	5.90305E-02
SXD	-8.04138E-01
SXV	1.35000E-01

## Q SHIFT EFFECTS

G22000		DQXDEX	DQX
-----		-----	---
-4.56809E+00	6.12823E-16	-9.13617E+00	-6.24001E-04
G00220		DQYDEY	DQY
-----		-----	---
2.80864E+01	2.85772E-15	5.61727E+01	3.83660E-03
G11110		DQXDEY	DQYDEX
-----		-----	-----
-2.14197E+01	1.42251E-14	-2.14197E+01	-2.14197E+01
		DQX	DQY
		---	---
		-1.46296E-03	-1.46296E-03

TABLE IVB  
-----

STANDARD AGS - BOOSTER WITH CHROMATICITY SEXTUPOLES AND EDDY CURRENTS DELTA(P)/P= 0.000000 [EX0 =6.8300E-05 EY0 =6.8300E-05]				
-----				
THIRD ORDER EFFECTS OF SEXTUPOLES [ RESONANCE EFFECTS]				
3Qx = 12				
-----				
COSINE	SINE	MODULUS	RANDOM	DE(S)
1.3100E-01	-7.6377E-02	1.5164E-01	6.9477E-01	2.2558E-02
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
1.0335E-01	7.5193E-03	3.4451E-02	1.5540E-02	7.1199E-02
Qx + 2Qy = 12				
-----				
COSINE	SINE	MODULUS	RANDOM	DE(S)
9.2324E-01	1.0306E-02	9.2329E-01	3.5080E+00	7.6304E-02
DE(R)	DQ(S)	DQ(R)	DQ20(S)	DQ20(R)
2.8992E-01	3.4124E-02	1.2965E-01	7.0524E-02	2.6795E-01






Fig. 4a Standard AGS - Booster

- ↑ = FOCUSING QUADRUPOLE
- ↓ = DEFOCUSING QUADRUPOLE
- = BENDING MAGNET (DIPOLE)
- X = SEXTUPOLE







Fig. 4b The Standard AGS - Booster Lattice

0 5  
METERS

NOTE: ALL DIMENSIONS ARE IN METERS