

BNL-104615-2014-TECH

AGS/AD/Tech Note No. 187;BNL-104615-2014-IR

ACCUMULATOR BOOSTER FOR THE AGS AS A HEAVY ION BOOSTER

Y. Y. Lee

April 1983

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Accelerator Department BROOKHAVEN NATIONAL LABORATORY Associated Universities, Inc. Upton, NY 11973

AGS Division Technical Note No. <u>187</u>

ACCUMULATOR BOOSTER FOR THE AGS AS A HEAVY ION BOOSTER

Y.Y. Lee and L.W. Smith

April 15, 1983

Acceleration of heavy ions in the AGS by boosting the existing tandem van deGraaff facility energy with a cyclotron has recently¹ been proposed. Also, a study² by M.Q. Barton suggests that heavy ions up to A = 32 could be injected directly from the tandem into the AGS and, that with the proposed cyclotron, ions up to A = 130 could be injected. The study reports that the only difficult technical problem is the AGS rf system.

In this note we would like to point out the possibility of using the proposed³ AGS accumulator/booster ring to accomplish the acceleration between the tandem and the AGS. We briefly discuss three technical areas involved: 1) A/B ring rf system, 2) A/B ring magnetic field and 3) injection into the A/B ring. Injection into the AGS would be as described in Reference 3.

RF System

There are a large number of free 3-m straight sections in the A/B ring. If three new rf cavities were installed each with a frequency sweep of a factor of 2.5, the sytem could accept heavy ions with $\beta > 0.04$ corresponding to 0.75 MeV/amu. With the ion currents available, there would be no appreciable loading on the cavities.

Magnetic Field

The A/B bending magnetic field at injection depends on the charge-to-mass ratio, Z/A, and is approximately

217 A/Z Gauss

Because of the design of the A/B ring magnets, they could produce precise fields of very low value. The final energy of ions in the A/B ring is approximately

- 2 -

1000 Z/A MeV/amu,

making injection into the AGS much easier because of the higher magnetic field in the AGS. Injection into the AGS is feasible for even the heaviest ions. Acceleration of ions other than fully stripped must still await improvement of the AGS vacuum system.

Injection

Injection into the A/B ring could be multilturn into horiziontal phase space as is suggested for injection into the AGS in Reference 2. Another method is to use charge exchange (stripping) injection into the A/B ring. Since the emittance of the tandem beam is very small, one could arrange a very narrow stripping foil which would intercept a very small fraction of the recirculating beam in the A/B ring.⁴ We estimate the accumulated current, I_o , in terms of incoming (tandem) current, I_i , as

$$I_o/I_i = \frac{P}{f(1-P)}$$

where P is the probability of an ion of the desired charge state emerging from the stripping foil and f is the fraction of the circulating beam reintercepted by the stripping foil. Although the quantity, P, is not precisely known, it appears⁵ to be near to 20%. Extrapolating from some Monte Carlo injection studies in the design of the A/B ring, we estimate $f \approx 1\%$. We then get

$$I_0/I_i \cong 25$$

For example, a 20 n-amp ¹⁹⁷Au current from the tandem gives 2.3 x 10^7 ions in the A/B ring or \approx 1.4 x 10^8 ions in the AGS with six A/B loads fast extracted into the AGS.

Although the points made in this note are by no means a design, the concept looks sufficiently interesting and cost saving to merit a more detailed examination.

References

7 1 2 4

- 1. Proposal for a Heavy Ion Facility at Brookhaven, BNL Informal Report BNL-30630 (1982).
- 2. M.Q. Barton, Acceleration of Heavy Ions in the AGS, BNL-32713.
- 3. A/B Ring Proposal, BNL Informal Report BNL-32949 (1983).
- 4. Y.Y. Lee, AGS Division Technical Note 186 (1983).
- 5. H.D. Betz, Review of Modern Physics 44, 465 (1972).

mn

Distribution: Department Administration Department S&P Staff C. Chasman

- P. Reardon
- A. Schwarzschild
- P. Thieberger

- 3 -