

BNL-104544-2014-TECH

AGS/AD/Tech Note No. 113;BNL-104544-2014-IR

BEAM LOSS DUE TO G10 TARGETING

L. N. Blumberg

September 1974

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.AT(30-1)-16 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Accelerator Department BROOKHAVEN NATIONAL LABORATORY Associated Universities, Inc. Upton, N.Y. 11973

AGS DIVISION TECHNICAL NOTE No. 113

L. Blumberg, J.W. Glenn, F. Hornstra[†], J. Keane, Y.Y. Lee, M. Month, L.W. Smith, A. Soukas

September 17, 1974

BEAM LOSS DUE TO G10 TARGETING

Introduction

A measurement of the losses on the slow beam extraction equipment indicates a doubling of these losses, and thus induced residual radiation in this equipment when about 30% of the circulating beam is targeted on G10.

G10 Induced Losses on Extraction Equipment

A LM was also installed in the G10 area. The beam was increased on G10, and the loss in the slow extraction area (Σ LM)* was noted. The data are plotted on Fig. 1. Up to ~ 15% on G10 the loss on the extraction equipment is independent of beam on G10. For higher levels on G10 the losses increase dramatically. A doubling of the losses was measured at $\approx 30\%$ of the beam on G10. Unformately, the AGS failed before a calibration of the G10 LM was made. A calibration was inferred by comparing to the G10 telescope and an old calibration of ≈ 1100 counts on the telescope per 10^{12} protons on G10.

*Described in AGS Technical Note 112

Distribution

Dept. Admin. EP&S, AGS S&P

[†]Summer Visitor, 1974, from Fermi National Accelerator Laboratory Batavia, Illinois

