

# SUPPRESSION OF TRANSVERSE INSTABILITIES BY OCTUPOLE LENSES

E. C. Raka

January 1975

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.E(30-1)-16 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Accelerator Department  
BROOKHAVEN NATIONAL LABORATORY  
Associated Universities, Inc.  
Upton, New York

AGS DIVISION TECHNICAL NOTE

No. 115

E.C. Raka, M. Month  
January 28, 1975

SUPPRESSION OF TRANSVERSE INSTABILITIES BY OCTUPOLE LENSES

The Landau damping of transverse bunched beam instabilities can only be achieved with octupoles. [Sextupoles, which act on transverse coherences through the chromaticity, can provide no Landau damping because of the periodic synchrotron motion.<sup>(1)</sup>] As a rule of thumb, the betatron frequency spread produced by such a field should exceed the total frequency shift (real + imaginary parts added in quadrature) due to the impedance driving the instability.

To estimate the required frequency spread we calculate

$$\Delta Q_i = \frac{U}{\Omega_0} = \frac{NRr_p}{\pi^2 \gamma^3 Q_0 \beta_B^2} \left[ \frac{\xi_1}{h^2} - \frac{\epsilon_1}{h^2} - \frac{1}{2a^2} \right]$$

which is the real part of the frequency shift due to the beam itself and its interaction with a perfectly conductive vacuum chamber. For a beam .85cm height with  $h = 4\text{cm}$  at  $\gamma = 12$  ( $\beta = 1$ ) and a bunching factor of .1 we obtain a  $\Delta Q_i \approx .0014$  at  $5 \times 10^{12}$  protons. The corresponding value of  $U$  is  $3.2 \times 10^3/\text{sec}$ . A single bunch e-folding time of  $2.5 \times 10^{-3}$  sec has been observed. This gives a growth rate of  $400 \text{ sec}^{-1}$  which corresponds to the  $V$  term in the stability analysis and indicates that  $U \gg V$ .

Since the beam is larger in the horizontal plane we shall only calculate the frequency spread in the vertical plane due to the horizontal motion, i.e. ...

$$\Delta Q_0 = \frac{1}{16\pi} \epsilon_H \beta_H \beta_V \frac{b}{p}$$

where  $\epsilon_H = 13.5 \times 10^{-6}$  Meter radians is the normalized horizontal emittance, p the momentum and  $b = \int B'''' ds$  is the integrated octupole strength. For  $\gamma = 12$  and  $b = 5.5 \times 10^4$  K gauss/meter<sup>2</sup> we obtain a  $\Delta Q_0 = 9.2 \times 10^{-3}$  for the octupoles located at horizontal  $\beta_{min} = 10.8$  meters. Thus at  $10^{13}$  ppp if the beam emittances do not change there is a factor  $> 3$  available in the strength over the simple criterion.

Octupoles have been used to stabilize the vertical instability in the CERN P.S. for several years (2) and recent measurements (3) indicate that in fact less octupole strength is required than given by the above estimate. It is thus probable that the octupole strength given should be sufficient to provide effective Landau damping at the projected AGS intensity level of  $10^{13}$  ppp.

References

1. F. Sacherer, "Transverse Bunched Beam Instabilities - Theory." Proceedings of the 9th International Conf. on High Energy Accelerators, 1974.
2. Y. Baconnier, J. Gareyte, P. Lefèvre; High Intensity Phenomena Observed in the PS. 1971 Particle Accelerator Conference Proceedings IEEE Nuclear Science NS-18 #3 June 1971.
3. J. Gareyte, F. Sacherer; Head Tail Instabilities in the CERN PS. Proceedings of the 9th International Conf. on High Energy Accelerators.

jh  
 Distr. Dept. Admin.  
 AGS Division Physicists