

BNL-104550-2014-TECH

AGS/AD/Tech Note No. 120;BNL-104550-2014-IR

STOCHASTIC COOLING FOR PEDESTRIANS

L. W. Smith

June 1976

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.E(30-1)-16 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Accelerator Department BROOKHAVEN NATIONAL LABORATORY Associated Universities, Inc. Upton, New York

AGS DIVISION TECHNICAL NOTE

No. 120

STOCHASTIC COOLING FOR PEDESTRIANS

L.W. Smith June 21, 1976

I. Introduction

This is a summary of some notes I made a couple of years ago in an attempt to get an elementary understanding of stochastic cooling. Recent results from the ISR, of which I am only vaguely aware, are not included.

van der Meer¹ and Palmer² have examined the possibility of damping incoherent beam oscillations by electronic feedback (stochastic cooling). This circumvention of Liouville's theorem is achieved by observing the mean lateral position of a sample N of the N_T circulating particles. (N/N_T might typically be $\approx 10^{-5}$.) Approximately one quarter of a betatron wavelength downstream, a momentum kick is applied to move the mean transverse momentum toward zero. After any "real" coherent oscillations have been damped the system operates on the sensed statistical fluctuations of the mean beam position of the finite number, N, of the particles sampled.

In Section II I review the damping of coherent oscillations and in Section III the damping of incoherent oscillations. Section IV points out some possible problem areas which need further study.

Notation similar to Palmer's is used below.

II. Coherent Damping

At some machine azimuth the mean beam transverse position is sensed and at some angle θ (in betatron oscillation phase space) downstream a momentum kick, proportional to the sensed mean beam position, is applied to move (in the impulse approximation) the mean beam momentum toward zero. Transit time spread² between sensing and correcting stations is ignored here as are the technical problems of getting the correction signal to arrive at the

p, n

correcting station at the proper time with sufficient band-pass.

Figure 1 is a diagram of the motion of the mean beam position in betatron phase space. Units of y' are chosen so that the orbits are circles.





At point A a portion of the beam is within the sensing station and a mean lateral position, $\bar{r} \cos \alpha$, is detected. At an angle, $\theta = \alpha + \beta$, later a momentum kick, $\delta y'$, is given (point B). For the moment we ignore noise and image forces. Then

$$\delta y' = a r \cos \alpha$$

(at is the "gain" of the feedback system) and so

$$\bar{r'}^2 = \bar{r}^2 + \delta y'^2 - 2\bar{r}\delta y' \sin \beta$$
 .

Averaging over all initial phases, α ,

$$\langle \bar{\mathbf{r}}'^2 \rangle_{\mathbf{p}} = \frac{11}{2\pi} \int_{0}^{2\pi} \bar{\mathbf{r}}'^2(\alpha) d\alpha$$

weagetaa

$$\langle \tilde{\mathbf{r}}^{\prime 2} \rangle_{\mathbf{p}} = \tilde{\mathbf{r}}^{2} \left(1 - \frac{2 \operatorname{asin} \theta - a^{2}}{2} \right)^{2}$$
(1)

per pass through the system. From (1) we see that damping obtains for

$$0 < a < 2 \sin \theta$$

with optimum³ damping for $a = \sin \theta$.

Consider now the effect of noise in the system. Define $\langle b^2 \rangle^{\frac{1}{2}}$ as the sensing noise in units of the RMS position of one particle. Palmer² quotes the ISR value $\langle b^2 \rangle^{\frac{1}{2}} = 10^4$ cm, i.e. the RMS error in sensing the mean position of 10⁴ particles is 1 cm. The expectation value of the noise contribution to \tilde{r}'^2 is then $\frac{a^2 \langle b^2 \rangle}{N^2}$, and

$$\langle \bar{\mathbf{r}}^2 \rangle_{\mathbf{p}} = \bar{\mathbf{r}}^2 \left(1 - \frac{2 \operatorname{asin} \theta - a^2}{2} + \frac{a^2 \langle \mathbf{b}^2 \rangle}{N^2 \bar{\mathbf{r}}^2} \right)$$
(2)

and damping obtains for

$$0 < a < \frac{2 \sin \theta}{1 + \frac{\langle b^2 \rangle}{N^2 r^2}}$$

If $(\bar{r}_{-}\bar{r}')/\bar{r}_{-}$ per pass through the system is small Eq. (2) can be written (we drop the notation $\langle \cdot \rangle_{p}$ for simplicity).

$$\bar{\mathbf{r}}' \simeq \bar{\mathbf{r}} \left(1 - \frac{2 \operatorname{asin} \theta - a^2}{4} + \frac{a^2 \langle b^2 \rangle}{2 \operatorname{N}^2 \bar{\mathbf{r}}^2}\right) ,$$

and if a driving term due, e.g., to image forces exists

$$\mathbf{\tilde{r}'} \simeq \mathbf{\bar{r}} \left(1 - \frac{2 \operatorname{asin} - a^2}{4} + \frac{a^2 \langle b^2 \rangle}{2 \operatorname{N}^2 \mathbf{r}^2} + \lambda \operatorname{N} \right) \qquad (3)$$

 λ can be a function of the vacuum chamber parameters, azimuthal extent of the sensing station, setc. Ignored here is any possible beneficial Landau damping and due to momentum and γ spread. From Eq. (3) damping obtains for (sin $\theta \approx 1$)

$$\frac{2\lambda N}{1 + \frac{\langle b^2 \rangle}{N^2 \bar{r}^2}} \approx a \approx \frac{2(1 - \lambda N)}{1 + \frac{\langle b^2 \rangle}{N^2 \bar{r}^2}} \cdot \frac{1 + \frac{\langle b^2 \rangle}{N^2 \bar{r}^2}}{N^2 \bar{r}^2}$$

III: Incoherent Damping

We use van der Meer's method of averaging over all combinations of amplitude and phase. It is assumed that there is sufficient transit time spread per revolution so that the samples, N, are statistically independent from one revolution to the next. Consider Fig. 1; at position B, in phase space, a momentum kick, $\delta y'$, proportional to the y coordinate of the center of gravity at point A, is given to the beam center of gravity. The y,y' coordinates of the center of gravity at point B are,

$$\frac{1}{N}\left<\Sigma rcos\beta\right>$$
 , $\frac{1}{N}\left<\Sigma rsin\beta\right>$,

where the summations are oversall particles. After the kick the new coordinates of the i-th particle are ([y], [y']),

$$\left[r_{i}\cos\beta_{i}\right], \left[r_{i}\sin\beta_{i}-a\cos\theta\frac{1}{N}\left<\Sigma r\cos\beta\right>-a\sin\theta\frac{1}{N}\left<\Sigma r\sin\beta\right>\right]$$

We take the square of the new amplitude, sum over i and divide by N. 4 Then .

- 4 -

integrating over all possible combinations⁵ of r and β to evaluate $\langle \Sigma r \cos \beta \rangle^2$ and $\langle \Sigma r \sin \beta \rangle^2$ we get (cross terms do not contribute)

$$\sigma_{0}^{\prime 2} = \frac{2}{\sigma} \left(1 - \frac{2 \operatorname{asin} \theta - a^{2}}{2 \operatorname{N}} \right)^{2} \qquad (4)$$

The functional similarity to Eq. (1) suggests that this result could have been arrived at more simply. Indeed it appears that we can, for predicting average values, use the Ergodic theorem to obtain Eq. (4) directly from the Eq. (1).

If we include noise, set $\theta \cong \pi/2$, and assume a is small,

$$\sigma'^{2} = \sigma^{2} \left(1 - \frac{a}{N} + \frac{a^{2} \langle b^{2} \rangle}{N^{2} \sigma^{2}} \right)$$

and the system damps for

$$0 < a \approx \frac{N_{\sigma}^2}{\langle b^2 \rangle}$$

with an optimum gain

$$a_{o} = \frac{N_{o}}{2\langle b^{2} \rangle} .$$

The optimum damping per passage through the system is

$$\sigma'^{2} = \sigma^{2} \left(1 - \frac{2}{4\langle b^{2} \rangle}\right)$$

$$\sigma' \simeq \sigma_{\sigma} \left(1 - \frac{\sigma_{\sigma}^2}{8 \langle \mathbf{b}_{\sigma}^2 \rangle^2} \right)$$

 $\sigma(t) \cong \frac{\sigma(o)}{1 + \frac{\sigma_o^2 t}{8 \langle b^2 \rangle}}$

giving

with a 1/2 folding time (in revolutions) of $8\langle b^2 \rangle / \sigma^2$ (o). Note that this rate can be maintained only if a is programmed to decrease proportionally to $\sigma(t)$. Otherwise $\sigma(t)$ will asymtotically approach $\sigma(o)/\sqrt{2}$.

IV. <u>Possible Problems</u>

We have made a number of simplifying assumptions, among them:

a) A smooth distribution in phase space. If there is any "lumpiness" in the phase space density due, e.g., to nearness to a nonlinear resonance, and if the sensing and correcting stations have sufficient higher moment sensitivity, antidamping of beam envelope oscillations could occur. Also, if the distribution in phase space is non-smooth the evaluation of $\langle \Sigma r \cos \theta \rangle^2$ and $\langle \Sigma r \sin \beta \rangle^2$ is (to me) obscure. This also implies a non-stationary process, i.e. a dependence on the absolute origin in time so the Ergodic theorem could not be applied.

b) Only the mean square noise has been considered. In practice b would probably be distributed similarly to the Rayleigh distribution with a probabil-

$$P(b) \approx \frac{2b}{\langle b \rangle} \exp(-b^2/\langle b^2 \rangle) ,$$

and one might expect this to lead to a long tailed distribution in particle position. However since the noise contributions to y' (and y) are small and in large number, the Central Limit theorem would predict that the distribution in y' (and y) approaches the Normal distribution. If the beam has some initial y_{max} (due, e.g., to shaving) any noise will populate the region y_{max} .

c) Only damping in two dimensional phase space has been considered. If there is mixing in six dimensional phase space, complications could result.

d) Bandwidth considerations have been ignored. To damp incoherent oscillations a large bandwidth is required. Assuming that the bandwidth (and noise) of the system is set by the RC of the sensing station, deterioration of the signal to noise ratio will start⁷ at frequencies $f \ll (2_{\Pi}RC)^{-1}$, which is just the frequency needed to "instantaneously" sample the fluctuations of the N particles within the station. Since $C \approx N$ (i.e. the length of the station) one may reduce the bandwidth at the expense of damping time but the deterioration of the signal to noise ratio will remain.

- 6 - ...

V. <u>Conclusions</u>

For an "ideal" system stochastic cooling seems to work, but technical difficulties may limit its usefulness. Computer simulations of "non-ideal" systems could be informative if round-off errors, etc. can be kept smalled enough.

References

- 1. S. van der Meer, CERN/ISR-PO/72-13 (CRISP CERN/72-31).
- 2. R.B. Palmer, CRISP 73-19 (BNL 18395).
- 3. Physically, this is due to the fact that in the absence of any phase information the best bet is $\alpha = 0$ since the probability density distribution of y is $\frac{1}{\pi(r^2-y^2)^{\frac{1}{2}}}$.
- 4. Actually $\frac{1}{N} \sum r_i^2$ is a biased estimator of σ but since N>>1 the point is academic.
- 5. Using Parseval's theorem, see e.g., E.T. Whittaker and G.N. Watson, "Modern Analysis", Cambridge, N.Y.
- 6. Roughly, the Ergodic theorem states that averages performed in time on a typical member of a system give the same results as corresponding averages taken over the system at the same moment, i.e. time and ensemble averages are equivalent under suitable conditions. See e.g. J.L. Doob, "Stochastic Processes", Wiley, N.Y.
- 7. Another way of looking at this is that the impulse approximation breaks down when the particle transit time through the station approaches the system rise time.

LWS/1sk

Distr: Dept. Admin. AGS Division S&P Scientific Staff in EP&S, ISA, and PD