

# WALKERS GUIDE TO DECOHERENCE TIMES IN STOCHASTIC COOLING

L. W. Smith

September 1976

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**  
USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.E(30-1)-16 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Accelerator Department  
BROOKHAVEN NATIONAL LABORATORY  
Associated Universities, Inc.  
Upton, New York

AGS DIVISION TECHNICAL NOTE

No. 124

A<sup>1)</sup> WALKER'S GUIDE TO DECOHERENCE TIMES IN STOCHASTIC COOLING

L. W. Smith

September 1, 1976

I. Introduction

Since there seems to be some confusion about the statistical processes inherent in stochastic cooling I give derivations of two probability density distributions that may serve to illuminate the situation.

In the ideal case the statistical fluctuation sensed survives to the correction station essentially intact but is smeared out to present a statistically independent sample by the time of the next pass through the sensing station. Any physically realizable system must necessarily be a compromise of these ideal conditions.

I consider two problems: 1) What is the time dependent probability density distribution of the number of particles in a sample of beam, and 2) What is the time dependent probability density distribution of the difference of the number of particles on either side of a sample of beam; with given initial conditions?

II. Unsplit Sample

Consider a sample of a beam containing, on the average,  $\langle n \rangle$ <sup>2)</sup> particles. Let  $\lambda$  be the probability per unit time that any one particle leaves the sample (due e.g. to momentum spread). We seek the probability,  $P_m(t)$ , that  $m$  particles are present at time,  $t$ , if there are exactly  $m_0$  present at time zero. From elementary considerations,

---

1) Random.

2) The notation  $\langle \rangle$  signifies expectation value.

$$P_m(t + \delta t) = P_{m-1}(t) \lambda \langle n \rangle \delta t + P_{m+1}(t) \lambda (m+1) \delta t + P_m(t) (1 - \lambda (\langle n \rangle + m) \delta t) ,$$

and letting  $\delta t \rightarrow 0$  ,

$$\frac{dP_m}{dt} = P_{m-1} \lambda \langle n \rangle + P_{m+1} \lambda (m+1) - P_m \lambda (\langle n \rangle + m) \quad (1)$$

This can be solved exactly by introducing the generating function

$$\varphi = \sum_m P_m x^m$$

and multiplying Eq. 1 by  $x^m$  and summing we get

$$\frac{\partial \varphi}{\partial t} = (1 - x) \left\{ -\lambda \langle n \rangle \varphi + \lambda \frac{\partial \varphi}{\partial x} \right\} .$$

Since  $\langle n \rangle$ ,  $m_0$ ,  $m \gg 1$  the Central Limit Theorem of statistics says that  $P_m$  will be very well approximated by the normal distribution (with mean and variance a function of time) so it is sufficient for our purposes to determine only the mean ( $\langle m \rangle$ ) and variance ( $\langle m^2 \rangle - \langle m \rangle^2$ ) of  $m$ .

Multiplying Eq. 1 by  $m$  and summing over  $m$ ,

$$\frac{d\langle m \rangle}{dt} = \lambda (\langle n \rangle - \langle m \rangle)$$

or

$$\langle m \rangle = \langle n \rangle - e^{-\lambda t} (\langle n \rangle - m_0) .$$

Multiplying Eq. 1 by  $m^2$  and summing over  $m$ ,

$$\frac{d\langle m^2 \rangle}{dt} = \lambda (\langle n \rangle + \langle m \rangle (2\langle n \rangle + 1) - 2\langle m^2 \rangle) ,$$

whose solution is

$$\langle m^2 \rangle = \langle m \rangle + \langle m \rangle^2 - m_0 e^{-2\lambda t} ,$$

or

$$\text{var } (m) = \langle m \rangle - m_0 e^{-2\lambda t} = \sigma_m^2 .$$

The time dependent probability density distribution of  $m$  is

$$P_m(t) = (2\pi \sigma_m^2)^{-\frac{1}{2}} \exp \left\{ -\frac{(m - \langle m \rangle)^2}{2\sigma_m^2} \right\} .$$

### III. Split Sample

Consider a sample of beam containing  $N$  particles. If there are  $m$  and  $n$  particles (initially  $m_0$  and  $n_0$ ) on either side of the beam the sensed transverse beam position is

$$\bar{y} = \frac{m-n}{m+n} \sigma_{\text{beam}} = \frac{m-n}{N} \sigma_{\text{beam}} .$$

Let  $\bar{v}$  equal the probability per unit time (due e.g. to betatron oscillations) that a particle will change sides. The joint probability distribution  $P_{m,n}(t)$  satisfies

$$\frac{dP_{m,n}}{dt} = P_{m-1, n+1} \bar{v}(n+1) + P_{m+1, n-1} \bar{v}(m+1) - P_{m,n} \bar{v}(m+n) . \quad (2)$$

As in II, we calculate only the mean and variance of  $m-n$ . Since  $m$  and  $n$  are not statistically independent we have

$$\text{var}(m-n) = \text{var}(m) + \text{var}(n) - 2 \text{cov}(m,n)$$

or

$$\text{var}(m-n) = \langle m^2 \rangle + \langle n^2 \rangle - (\langle m \rangle - \langle n \rangle)^2 - 2\langle mn \rangle .$$

Multiplying Eq. 2 by  $n$ ,  $m$ ,  $n^2$ ,  $m^2$ , and  $mn$  and each time summing over  $m$ ,  $n$ , we get

$$\langle m - n \rangle = (m_0 - n_0) e^{-2\bar{v}t} ,$$

and, after a moderate amount of algebra

$$\text{var}(m-n) = N(1 - e^{-4\bar{v}t}) = \sigma_{m-n}^2 .$$

The time dependent probability density distribution of m-n is

$$P(t) = (2\pi \sigma_{m-n}^2)^{-\frac{1}{2}} \exp \left\{ -\frac{[(m-n) - \langle m-n \rangle]^2}{2\sigma_{m-n}^2} \right\} .$$

#### IV. Conclusion

Using the above, and the assumption that the initial (t=0) conditions are normally distributed, the interested reader, with a little arithmetic (or computer time), may calculate the odds<sup>3</sup> on the success of a particular stochastic cooling system.

- 3) "For most men ('till by losing rendered sager) will bet with a wager" George Gordon, Lord Byron.  
will back their own opinions with a wager"

Dist: AD Sci Staff  
J. Sanford

George Gordon, Lord Byron.

Dist: AD Sci. Staff  
J. Sanford