



BNL-105071-2014-TECH

Booster Technical Note No. 24;BNL-105071-2014-IR

APERTURE LIMITATIONS DUE TO NON-LINEAR COUPLING

G. Parzen

April 1986

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

APERTURE LIMITATIONS DUE TO NON-LINEAR COUPLING

Booster Technical Note

No. 24

G. PARZEN

APRIL 2, 1986

*HIGH ENERGY FACILITIES
Brookhaven National Laboratory
Upton, N.Y. 11973*

The sextupoles for correcting chromaticity and the eddy current sextupole fields can drive the $2\nu_x - 2\nu_y = 0$ coupling resonance.

y - motions that start at $y_0 = 27\text{mm}$ at QD, corresponding to $\epsilon_y = 50 \times 10^{-6}$, will grow as large as 38mm , depending on the choice of ν_x, ν_y .

The next two graphs show the implications of this non-linear growth on the acceptable emittance.

Depending on one's assumptions, the acceptable emittance could be as low as $\epsilon_y = 38 \times 10^{-6}$.


This non-linear effect could be avoided by moving ν_x, ν_y about .05 from the $\nu_x = \nu_y$ resonance. However this may not be desirable as this puts ν_x, ν_y in a region where there are many higher order resonance lines.

It is possible that this non-linear effect could be corrected by a system of octupole correctors, since octupoles can also excite the $2\nu_y - 2\nu_x = 0$ resonance.

The first figure plots the maximum value of y at QD, y_{\max} , against the initial value of χ at QF, χ_0 . If one assumes that the largest safe value of y_{\max} is 30mm, then the largest allowed χ_0 is 23mm corresponding to $\epsilon = 38 \times 10^{-6}$.

The second figure shows the effect of the ν_x, ν_y choice on the non-linear coupling. ν_x is held constant at $\nu_x = 4.82$ while ν_y is varied from 4.75 to 4.89. The largest coupling occurs at $\nu_x \approx \nu_y$ and the coupling is removed by going .05 away from the $\nu_x = \nu_y$ resonance. The coupling depends on the choice of the initial phase of χ , and two curves are shown for two different choices of χ_0 and χ'_0 .

Non-Linear γ -growth



Non-Linear Coupling

