



BNL-105831-2014-TECH

EP&S No. 116;BNL-105831-2014-IR

The dependence of low momentum particle production on target material and thickness

J. W. Glenn

July 1985

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Alternating Gradient Synchrotron Department
BROOKHAVEN NATIONAL LABORATORY
Associated Universities, Inc.
Upton, New York 11973

EP&S Division Technical Note
No. 116

THE DEPENDENCE OF LOW MOMENTUM PARTICLE PRODUCTION
ON
TARGET MATERIAL AND THICKNESS

J.W. Glenn, D.M. Lazarus, P.H. Pile, J. Sculli, J.C. Walker

19 July 1985

Motivated by a need to simultaneously optimize antiproton production in Beam C1 and negative kaon flux in the LESBI and to determine the best target for neutrino (pion) production, pion and antiproton fluxes produced by 28.4 GeV/c proton incident on Al, Cu and Pt targets of various lengths at 0° and 1.4 GeV/c were made in the C1 beam while 10.5° kaon production was simultaneously measured at .870 GeV/c in the LESBI. The measurements were normalized to the secondary emission chamber (SEC) immediately upstream of the "C" target but no attempt was made to correct the measurements for beam acceptance to obtain an absolute normalization. The counter telescope which monitors targeting is of course sensitive to target material and length and is therefore not useful for normalization.

The targets used and the data obtained are given in Table I. The interaction length for Pt was obtained by interpolation using a plot (Fig. 1) of interaction lengths vs atomic number for a variety of materials taken from the "Blue Book."¹

The 0° data obtained in C1 indicates both the pion and antiproton production saturates at less than half an interaction length whereas the 10.5° kaon data shows no clear saturation at one interaction length. This is easily understood as the result of absorption of secondary particles produced at small angles with a relatively large average amount of material between the point of production and the downstream surface of the target. Secondaries produced at 10.5° pass through comparatively little material before exiting from the side of the target in the direction of the LESBI.

1. Particle Properties Data Booklet - Particle Data Group LBL (1984).

TABLE I

l l/l_0	1.88" Pt	3.5" Pt	.74" Cu	2.96" Cu	5.25" Cu	5.25" Al
	.522	.972	.125	.499	.886	.338
0° Data						
π^- /SEC	67.3	60.7	55.4	103.5	156.4	124.8
\bar{p} /SEC	.120	.120	.0532	.159	.156	.094
10.5° Data						
K^- /SEC	14.7	21.0/18.7	5.65	14.5	15.9	8.0
π^-/K^-	5.2	4.9/5.9	5.3	6.2	11.1	12.2

Figures 2 and 3 indicate that at 0° the Cu targets yield a greater number of both pions and antiprotons than the equivalent length of Pt. The negative pion to antiproton ratio is lower for the Pt target and is preferable for Exp. 789 whose usable antiproton flux is limited by accidentals resulting from beam pions. The 10.5° negative kaon data is shown in Fig. 4 and the π^-/K^- ratio in Fig. 5. A somewhat higher yield of K^- is obtained from Pt than from Cu but the primary benefit for kaon users arises from the shorter interaction length of Pt which allows good kaon production from a target of less than 3 inches reducing the depth of focus at the mass slit. Figure 5 indicates a marked degradation of the π^-/K^- ratio for lengths greater than 4 inches.

Conclusions

A Pt target of 2-3 inches would provide the optimum compromise for simultaneous operation. Both the \bar{p}/π^- ratio in C1 and the π^-/K^- ratio in LESBI would benefit from a shorter target and more protons on target.

Small angle negative pion production is enhanced by the use of Cu target. The acceptance of Beam C1 is ± 20 mr vertically and ± 40 mr horizontally. The narrow band neutrino horn acceptance includes the angular range of 10 mr to 140 mr and therefore a target length between $.5 < l/l_0 < 1$ is called for. Cu appears to be the preferred target material to simultaneously optimize flux and depth of focus for the narrow band ν beam.




Figure 1. The dependence of interaction length l_0 on atomic weight.

1.4 GeV/c π^- 0°




Figure 2. The yield or number of π^- per SEC count at 0° and 1.4 GeV/c vs. target length/interaction length.

1.4 GeV/c \bar{p} 0°




Figure 3. The yield or number of \bar{p} per SEC count at 0° and 1.4 GeV/c vs. target length/interaction length.




Figure 4. The yield or number of K^- per SEC count at 10.5° and .87 GeV/c per SEC count vs. target length/interaction length.

.87 GeV/c 10.5°




Figure 5. The π^-/K^- ratio vs. target length in the LESBI at 10.5° and .87 GeV/c.