

## Possibility of obtaining secondary alpha beams at AGS

C. L. Wang

March 1969

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.AT-30-2-GEN-16 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.



Accelerator Department  
BROOKHAVEN NATIONAL LABORATORY  
Associated Universities, Inc.  
Upton, L.I., N.Y.

EP & S DIVISION TECHNICAL NOTE

No. 26

C.L. Wang

March 11, 1969

~~POSSIBILITY~~ POSSIBILITY OF OBTAINING SECONDARY ALPHA BEAMS AT AGS

No alpha particles were detected both in the early experiments<sup>1,2</sup> at the CERN Proton Synchrotron and in a recent measurement<sup>3</sup> at the Argonne ZGS. The measurements reported here show, however, that there are sizable fluxes of alphas produced at the Brookhaven AGS. Details of the experiment will appear in a later publication,<sup>4</sup> and only a brief description is presented here. This short note is intended primarily for a quick communication to potential users,<sup>5</sup> regarding the possibility of obtaining useful secondary alpha beams at the AGS for various experimentation.

The measurement was carried out at the AGS test beam. Secondary particles produced at 18°48' from the 28.5 BeV internal protons incident on the BeO target at G10 were momentum analyzed by a bending magnet, and then classified according to their time-of-flight (TOF) and their energy loss ( $dE/dx$ ) in a scintillator. The  $dE/dx$  analysis is essential for the alpha identification, because  $He^{4++}$  has twice the momentum of the deuteron, and hence the same TOF. An example of the TOF spectrum at 1.4 BeV/c is shown in Fig. 1(a). Here the detection efficiencies of pions and protons were reduced purposely in order to display the peaks better. In Fig. 1 the data plotted in (a') is the same as (a) except that the vertical scale is expanded 8 times. Any portion of the TOF spectrum can be analyzed in terms of the energy loss of the particles involved. The  $dE/dx$  spectrum of the deuteron peak in Fig. 1(a) is shown in Fig. 1(b) and (b'). The data in (b) is the same as in (b') except that the vertical scale is expanded 128 times. Clearly, the deuteron peak in the TOF spectrum is decomposed into deuteron and alpha peaks in the  $dE/dx$  analysis. The scattered dots under the deuteron peak in Fig. 1(b) is purely instrumental; namely when (b') is expanded 128 times in the vertical direction, many channels in the deuteron peak go beyond the range of the oscilloscope display and somehow show up as the scattered dots. To confirm that the small peak is really due to alphas, the  $dE/dx$  system was calibrated by the protons and  $He^{3++}$ . Here,  $He^3$  has twice the momentum of the proton. The resulting  $dE/dx$  spectrum is shown in Fig. 1(c) and (c'). The data



in (c) is the same as in (c') except that the vertical scale is expanded 104 times. Thus, the relative abscissas of the proton, deuteron,  $\text{He}^{3++}$  and  $\text{He}^{4++}$  peaks, and hence their relative energy losses establish the first unambiguous identification of the alpha production at the high energy accelerators.

The ratio of alpha to deuteron production is given in Fig. 2. The alpha production is of the order of 1% of the deuteron production in this momentum range. Preliminary data from our recent measurement<sup>6</sup> on secondary production at the AGS test beam show that the deuteron to pion ratio is about 1%. Thus, the ratio of alpha to pion production is of the order of  $10^{-4}$  at the momentum range covered here, and this can be used as a "rule of thumb" for a rough estimate of the available alpha fluxes in the existing experimental beams at the AGS. From the consideration of transverse momentum distribution, it is probable that at larger production angles, the alpha to pion ratio can be even higher than  $10^{-4}$  in the pion momentum range above 1 BeV/c. The absolute yield of alphas would of course be smaller.

An interesting application of the alpha beam is the measurement of the cross sections for  $\alpha + \alpha \rightarrow \text{D}$  (or  $\text{He}^3$ ) +--- below 1000 MeV/nucleon. Such cross sections are critical for the interpretation of the D,  $\text{He}^3$  and  $\text{He}^4$  composition of the cosmic rays, and are very valuable for the understanding of the origin and propagation of the primary cosmic rays in the interstellar space.<sup>7</sup> Such experiment now becomes feasible with the possible secondary alpha beams and the helium bubble chambers in operation at Argonne. A cloud chamber triggerable with electronic counter logics might even be preferable. A detailed discussion of this subject will be given in a separate paper.<sup>8</sup>

I am grateful to Dr. James R. Sanford for his interest in the measurement of alpha production, and his careful reading of the manuscript. Thanks are due to Mr. Michael Koson for his technical assistance and to Mr. Joseph Lypecky for his various assistance on the AGS floor.

Distr: E. Engels - Harvard  
 D. Frisch - MIT  
 L. Lederman - Columbia  
 A. Mann - Pennsylvania  
 S. Munday - CERN  
 J. Sandweiss - Yale  
 A. Tollestrup - Cal Tech  
 W. Wenzel - LRL  
 A. Wattenberg - Illinois  
 D.H. White - Cornell  
 M. White - Princeton

EP & S Div. Physicists  
 Dept. Administration  
 R. Cool  
 F. Huson  
 T. Kycia  
 S. Ozaki  
 R. Rau  
 A. Thorndike  
 F. Turkot



## Figure Captions

- Fig. 1 (a) Time-of-flight spectrum of 1.4 BeV/c secondary particles produced at 18°48' from 28.5 BeV protons incident on Be. The abscissa is the flight time and the ordinate is the relative abundance of the particles; both in arbitrary scales. Here, the detection efficiencies of pions and protons were reduced purposely in order to give a better balance in the height of the peaks.
- (a') Data in (a') is the same as in (a) except that the vertical scale is expanded 8 times.
- (b) The dE/dx spectrum the deuteron peak in (a) is decomposed into deuteron and alpha peaks. The abscissa is the energy loss and the ordinate is the relative abundance corresponding to particular energy losses; both in arbitrary scale. The dots under the deuteron peak are due to overflows of corresponding channels in an expanded display of the pulse height analyzer. (See text)
- (b') Data in (b') is the same as in (b) except that the vertical scale is reduced 128 times. Note the alpha peak does not show up even as a kink.
- (c) The dE/dx spectrum of protons and  $\text{He}^3$ , used to calibrate the dE/dx system.
- (c') Data in (c') is the same as in (c) except that the vertical scale is reduced 128 times. Note the relative abscissas of proton, deuteron,  $\text{He}^3$  and  $\text{He}^4$  peaks. Also note that the  $\text{He}^3$  peak does not show up even as a kink. Here the detection efficiency of protons was not reduced as in (a).
- Fig. 2 Ratio of alpha to deuteron production as a function of secondary momentum per nucleon. The solid line is drawn to guide the eyes.



## References

1. V.T. Cocconi, T. Fazzini, G. Fidecaro, M. Legros, N.H. Lipman, and A.W. Merrison Phys Rev Letters 5 19 (1960).
2. L. Gilly, B. Leontic, A. Lundby, R. Meurier, J.P. Stroot and M. Szeptycka, Proceedings of the 1960 Annual International Conference on High-Energy Physics at Rochester (Interscience Publishers, Inc. N.Y. 1960) P. 808.
3. Gary James Marmer, Ph.D. Thesis 1968, The Ohio State University (Unpublished).
4. Alpha Production by 28.5 BeV/Protons Incident on BeO ———To be published.
5. There was an inquiry about an alpha beam at the AGS———J.R. Sanford, Private Communication.
6. G.W. Bennett, E.J. Bleser, I.H. Chiang, L. Ettlinger, J.D. Fox, A.L. Read, J. Sculli, T.E. Toohig, C.L. Wang, T.O. White, R.R. Wilson, T. Yamanouchi, Preliminary experiment of BNL/Johns Hopkins/NAL/Rochester collaboration.
7. R.E. Lingenfelter and R. Ramaty, Private communication; Also see e.g. 10th International Conference on Cosmic Rays, Canadian Journal of Physics 46 S627 (1968).
8. Alpha Production, Alpha Beam and Cosmology———In preparation.






Fig. 1




KEE SEMI-LOGARITHMIC 46 4970  
2 CYCLES X 70 DIVISIONS MADE IN U.S.A.  
KEUFFEL & ESSER CO.

$\sigma/\rho$

0.01

0.001



0.5

1.0

1.5 BeV/c/nucleon

Fig. 2

Momentum