

Results for intrabeam scattering growth rates for a bi-gaussian distribution

G. Parzen

August 2005

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.


DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

C-A/AP/#144
August 2005

Results for the intrabeam scattering growth Rates for a bi-gaussian distribution

George Parzen



**Collider-Accelerator Department
Brookhaven National Laboratory
Upton, NY 11973**

Results for the intrabeam scattering growth rates for a bi-gaussian distribution.

George Parzen

March 2004
BNL Report C-A/AP No.144

Abstract

This note lists results for the intrabeam scattering growth rates for a bi-gaussian distribution. The derivation of these results will be given in a future note.

Introduction

This note finds results for the intrabeam scattering growth rates for a bi-gaussian distribution.

The bi-gaussian distribution is interesting for studying the possibility of using electron cooling in RHIC. Studies done using the SIMCOOL program [1] indicate that in the presence of electron cooling, the beam distribution changes so that it develops a strong core and a long tail which is not described well by a gaussian, but may be better described by a bi-gaussian. Being able to compute the effects of intrabeam scattering for a bi-gaussian distribution would be useful in computing the effects of electron cooling, which depend critically on the details of the intrabeam scattering.

Gaussian distribution

Before defining the bi-gaussian distribution, the gaussian distribution will be reviewed.

$Nf(x, p)$ gives the number of particles in $d^3x d^3p$, where N is the number of particles in a bunch. For a gaussian distribution, $f(x, p)$ is given by

$$f(x, p) = \frac{1}{\Gamma} \exp[-S(x, p)] \quad (1)$$

$$\begin{aligned}
S &= S_x + S_y + S_s \\
S_x &= \frac{1}{\bar{\epsilon}_x} \epsilon_x(x_\beta, p_{x\beta}/p_0) \\
x_\beta &= x - D(p - p_0)/p_0 \\
p_{x\beta}/p_0 &= p_x/p_0 - D'(p - p_0)/p_0 \\
\epsilon_x(x, x') &= \gamma_x x^2 + 2\alpha_x x x' + \beta_x x'^2 \\
S_y &= \frac{1}{\bar{\epsilon}_y} \epsilon_y(y, p_y/p_0) \\
\epsilon_y(y, y') &= \gamma_y y^2 + 2\alpha_y y y' + \beta_y y'^2 \\
S_s &= \frac{1}{2\sigma_s^2} (s - s_c)^2 + \frac{1}{2\sigma_p^2} ((p - p_0)/p_0)^2 \\
S_s &= \frac{1}{\bar{\epsilon}_s} \left(\frac{1}{\beta_s} (s - s_c)^2 + \beta_s ((p - p_0)/p_0)^2 \right) \\
\beta_s &= \sigma_s / \sigma_p \\
\bar{\epsilon}_s &= 2\sigma_s \sigma_p \\
S_s &= \frac{1}{\bar{\epsilon}_s} \epsilon_s(s - s_c, (p - p_0)/p_0) \\
\Gamma &= \int d^3x d^3p \exp[-S(x, p)] \\
\Gamma &= \pi^3 \bar{\epsilon}_x \bar{\epsilon}_y \bar{\epsilon}_s p_0^3 \\
\bar{\epsilon}_i &= \langle \epsilon_i(x, p) \rangle \quad i = x, y, s
\end{aligned} \quad (2)$$

D is the horizontal dispersion. $D' = dD/ds$. $\langle \rangle$ indicates an average over all the particles in a bunch.

Growth rates for a Gaussian distribution

In the following, the growth rates are given in the Rest Coordinate System, which is the coordinate system moving along with the bunch. Growth rates

are given for $\langle p_i p_j \rangle$. From these one can compute the growth rates for $\langle \epsilon_i \rangle$ using the relations given at the end of this note.

$$\begin{aligned}
\frac{1}{p_0^2} \frac{d}{dt} \langle p_i p_j \rangle &= \frac{N}{\Gamma} \int d^3 \Delta \exp[-R] C_{ij} \\
C_{ij} &= \frac{2\pi}{p_0^2} (r_0/2\bar{\beta}^2)^2 (|\Delta|^2 \delta_{ij} - 3\Delta_i \Delta_j) 2\bar{\beta}c \ln[1 + (2\bar{\beta}^2 b_{max}/r_0)^2] \\
\bar{\beta} &= \beta_0 \gamma_0 |\Delta/p_0| \\
r_0 &= Z^2 e^2 / M c^2 \\
R &= R_x + R_y + R_s \\
R_x &= \frac{2}{\beta_x \bar{\epsilon}_x} [\gamma^2 D^2 \Delta_s^2 + (\beta_x \Delta_x - \gamma \tilde{D} \Delta_s)^2] / p_0^2 \\
\tilde{D} &= \beta_x D' + \alpha_x D \\
R_y &= \frac{2}{\beta_y \bar{\epsilon}_y} \beta_y^2 \Delta_y^2 / p_0^2 \\
R_s &= \frac{2}{\beta_s \bar{\epsilon}_s} \beta_s^2 \gamma^2 \Delta_s^2 / p_0^2 \tag{3}
\end{aligned}$$

The integral over $d^3 \Delta$ is an integral over all possible values of the relative momentum for any two particles in a bunch. β_0, γ_0 are the beta and gamma corresponding to p_0 , the central momentum of the bunch in the Laboratory Coordinate System. $\gamma = \gamma_0$

The above 3-dimensional integral can be reduced to a 2-dimensional integral by integrating over $|\Delta|$ and using $d^3 \Delta = |\Delta|^2 d|\Delta| \sin\theta d\theta d\phi$.

$$\begin{aligned}
\frac{1}{p_0^2} \frac{d}{dt} \langle p_i p_j \rangle &= \frac{N}{\Gamma} 2\pi p_0^3 \left(\frac{r_0}{2\gamma_0^2 \beta_0^2} \right)^2 2\beta_0 \gamma_0 c \int \sin\theta d\theta d\phi (\delta_{ij} - 3g_i g_j) \\
&\quad \frac{1}{F} \ln \left[\frac{\hat{C}}{F} \right] \\
g_3 &= \cos\theta = g_s \\
g_1 &= \sin\theta \cos\phi = g_x \\
g_2 &= \sin\theta \sin\phi = g_y \\
\hat{C} &= 2\gamma_0^2 \beta_0^2 b_{max}/r_0 \\
F &= R/(|\Delta|/p_0)^2
\end{aligned}$$

$$\begin{aligned}
F &= F_x + F_y + F_s \\
F_x &= \frac{2}{\beta_x \bar{\epsilon}_x} [\gamma^2 D^2 g_s^2 + (\beta_x g_x - \gamma \tilde{D} g_s)^2] \\
F_y &= \frac{2}{\beta_y \bar{\epsilon}_y} \beta_y^2 g_y^2 \\
F_s &= \frac{2}{\beta_s \bar{\epsilon}_s} \beta_s^2 \gamma^2 g_s^2
\end{aligned} \tag{4}$$

Bi-Gaussian distribution

The bi-gaussian distribution will be assumed to have the form given by the following.

$Nf(x, p)$ gives the number of particles in $d^3x d^3p$, where N is the number of particles in a bunch. For a bi-gaussian distribution, $f(x, p)$ is given by

$$f(x, p) = \frac{N_a}{N} \frac{1}{\Gamma_a} \exp[-S_a(x, p)] + \frac{N_b}{N} \frac{1}{\Gamma_b} \exp[-S_b(x, p)] \tag{5}$$

In the first gaussian, to find Γ_a, S_a then in the expressions for Γ, S , given above for the gaussian distribution, replace $\bar{\epsilon}_x, \bar{\epsilon}_y, \bar{\epsilon}_s$ by $\epsilon_{xa}, \epsilon_{ya}, \epsilon_{sa}$. In the second gaussian, in the expressions for Γ, S , replace $\bar{\epsilon}_x, \bar{\epsilon}_y, \bar{\epsilon}_s$ by $\epsilon_{xb}, \epsilon_{yb}, \epsilon_{sb}$. In addition. $N_a + N_b = N$. This bi-gaussian has 7 parameters instead of the three parameters of a gaussian.

Growth rates for a Bi- Gaussian distribution

In the following, the growth rates are given in the Rest Coordinate System, which is the coordinate system moving along with the bunch. Growth rates are given for $\langle p_i p_j \rangle$. From these one can compute the growth rates for $\langle \epsilon_i \rangle$ using the relations given at the end of this note.

$$\begin{aligned}
\frac{1}{p_0^2} \frac{d}{dt} \langle p_i p_j \rangle &= N \int d^3 \Delta C_{ij} \left[\left(\frac{N_a}{N} \right)^2 \frac{\exp(-R_a)}{\Gamma_a} + \left(\frac{N_b}{N} \right)^2 \frac{\exp(-R_b)}{\Gamma_b} \right. \\
&\quad \left. + 2 \frac{N_a N_b}{N^2} \frac{\Gamma_c}{\Gamma_a \Gamma_b} \exp(-T) \right] \\
C_{ij} &= \frac{2\pi}{p_0^2} (r_0 / 2\bar{\beta}^2)^2 (|\Delta|^2 \delta_{ij} - 3\Delta_i \Delta_j) 2\bar{\beta} c \ln[1 + (2\bar{\beta}^2 b_{max} / r_0)^2]
\end{aligned}$$

$$\begin{aligned}\bar{\beta} &= \beta_0 \gamma_0 |\Delta/p_0| \\ r_0 &= Z^2 e^2 / M c^2\end{aligned}$$

$$\begin{aligned}\frac{1}{\bar{\epsilon}_{ic}} &= \frac{1}{2} \left(\frac{1}{\bar{\epsilon}_{ia}} + \frac{1}{\bar{\epsilon}_{ib}} \right) \quad i = x, y, s \\ \frac{1}{\bar{\epsilon}_{id}} &= \frac{1}{2} \left(\frac{1}{\bar{\epsilon}_{ia}} - \frac{1}{\bar{\epsilon}_{ib}} \right)\end{aligned}$$

$$\begin{aligned}r_0 &= Z^2 e^2 / M c^2 \\ \Gamma_a &= \pi^3 \bar{\epsilon}_{sa} \bar{\epsilon}_{xa} \bar{\epsilon}_{ya} p_0^3\end{aligned}$$

$$\begin{aligned}R_a &= R_{xa} + R_{ya} + R_{sa} \\ R_{xa} &= \frac{2}{\beta_x \bar{\epsilon}_x} [\gamma^2 D^2 \Delta_s^2 + (\beta_x \Delta_x - \gamma \tilde{D} \Delta_s)^2] / p_0^2 \\ \tilde{D} &= \beta_x D' + \alpha_x D \\ R_{ya} &= \frac{2}{\beta_y \bar{\epsilon}_y} \beta_y^2 \Delta_y^2 / p_0^2 \\ R_{sa} &= \frac{2}{\beta_s \bar{\epsilon}_s} \beta_s^2 \gamma^2 \Delta_s^2 / p_0^2\end{aligned}$$

$$\begin{aligned}T &= T_x + T_y + T_s \\ T_x &= R_{xc} - R_{xd} \\ T_y &= R_{yc} - R_{yd} \\ T_s &= R_{sc} - R_{sd}\end{aligned}$$

$$\begin{aligned}R_{xd} &= 2\{[-\gamma D \bar{\Delta}_s]^2 \\ &\quad + [(\beta_x \bar{\Delta}_x - \gamma \bar{D} \bar{\Delta}_s)]^2\} \\ &\quad / (\beta_x \bar{\epsilon}_{xd}^2 / \bar{\epsilon}_{xc}) \\ R_{yd} &= \frac{2\beta_y}{\bar{\epsilon}_{yd}^2 / \bar{\epsilon}_{yc}} \bar{\Delta}_y^2 \\ R_{sd} &= \frac{2\beta_s}{\bar{\epsilon}_{sd}^2 / \bar{\epsilon}_{sc}} \bar{\Delta}_s^2 \\ \bar{\Delta}_i &= \Delta_i / p_0 \quad i = x, y, s\end{aligned}\tag{6}$$

R_a, R_b, R_c are each the same as R_a given above except that $\bar{\epsilon}_{ia}$ are replaced

by $\bar{\epsilon}_{ia}, \bar{\epsilon}_{ib}, \bar{\epsilon}_{ic}$ respectively.

The above 3-dimensional integral can be reduced to a 2-dimensional integral by integrating over $|\Delta|$.

$$\begin{aligned}
\frac{1}{p_0^2} \frac{d}{dt} \langle p_i p_j \rangle &= 2\pi p_0^3 \left(\frac{r_0}{2\gamma_0^2 \beta_0^2} \right)^2 2\beta_0 \gamma_0 c \int \sin\theta d\theta d\phi (\delta_{ij} - 3g_i g_j) \\
&\quad N \left[\left(\frac{N_a}{N} \right)^2 \frac{1}{\Gamma_a F_a} \ln \left[\frac{\hat{C}}{F_a} \right] + \left(\frac{N_b}{N} \right)^2 \frac{1}{\Gamma_b F_b} \ln \left[\frac{\hat{C}}{F_b} \right] \right. \\
&\quad \left. + 2 \frac{N_a N_b}{N^2} \frac{\Gamma_c}{\Gamma_a \Gamma_b} \frac{1}{G} \ln \left[\frac{\hat{C}}{G} \right] \right] \\
g_3 &= \cos\theta = g_s \\
g_1 &= \sin\theta \cos\phi = g_x \\
g_2 &= \sin\theta \sin\phi = g_y \\
\hat{C} &= 2\gamma_0^2 \beta_0^2 b_{max}/r_0 \\
F_i &= R_i / (|\Delta|/p_0)^2 \quad i = a, b, c \\
G &= T / (|\Delta|/p_0)^2
\end{aligned} \tag{7}$$

F_a, F_b, F_c are each the same F that was defined for the Gaussian distribution except that the $\bar{\epsilon}_i$ are replaced by $\bar{\epsilon}_{ia}, \bar{\epsilon}_{ib}, \bar{\epsilon}_{ic}$ respectively.

The above results for the growth rates for a bi-gaussian distribution are expressed as an integral which contains 3 terms, each of which is similar to the one term in the results for the gaussian distribution. These three terms may be given a simple interpretation. The first term represents the contribution to the growth rates due to the scattering of the N_a particles of the first gaussian from themselves, the second term the contribution due to the scattering of the N_b particles of the second gaussian from themselves, and the third term the contribution due to the scattering of the N_a particles of the first gaussian from the N_b particles of the second gaussian.

Emittance growth rates

One can compute growth rates for the average emittances, $\langle \epsilon_i \rangle$ in the Laboratory Coordinate System, from the growth rates for $\langle p_i p_j \rangle$ in the Rest Coordinate System. In the following, dt is the time interval in the Laboratory System and $d\tilde{t}$ is the time interval in the Rest System. $dt = \gamma d\tilde{t}$

$$\begin{aligned}
 \frac{d}{dt}\epsilon_x &= \frac{\beta_x}{\gamma} \frac{d}{d\tilde{t}} \langle p_x^2/p_0^2 \rangle + \frac{D^2 + \tilde{D}^2}{\beta_x} \gamma \frac{d}{d\tilde{t}} \langle p_s^2/p_0^2 \rangle - 2\tilde{D} \frac{d}{d\tilde{t}} \langle p_x p_s/p_0^2 \rangle \\
 \frac{d}{dt}\epsilon_y &= \frac{\beta_y}{\gamma} \frac{d}{d\tilde{t}} \langle p_y^2/p_0^2 \rangle \\
 \frac{d}{dt}\epsilon_s &= \beta_s \gamma \frac{d}{d\tilde{t}} \langle p_s^2/p_0^2 \rangle
 \end{aligned} \tag{8}$$

I thank I. Ben-Zvi for his comments and encouragement. The results given above were found using the results given in references [2,3,4]. The derivation of the results is given in Ref.[5]

References

1. V.Parhomchuk and I. Ben-Zvi, BNL report C-A/AP/47, April 2001; A. Fedotov, Y. Eidelman (Private Communication 2004)
2. G.Parzen, BNL report C-A/AP/No.150 (2004)
3. A. Piwinski Proc. 9th Int. Conf. on High Energy Accelerators (1974) 405, M. Martini CERN PS/84-9 (1984), A. Piwinski Cern 87-03 (1987) 402, A. Piwinski CERN 92-01 (1992) 226
4. J.D. Bjorken and S.K. Mtingwa, Part. Accel.13 (1983) 115, K. Kubo and K. Oide Phys. Rev. S.T.A.B., 4, (2001) 124401
5. G. Parzen, BNL report C-A/AP/No.169 (2004)