



BNL-105877-2014-TECH

EP&S No. 162;BNL-105877-2014-IR

## Soil activation in various AGS/RHIC sites by Monte Carlo simulation

K. Yip

November 2005

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

*For Internal Distribution Only*

*Experimental Support and Facilities Division  
Collider-Accelerator Department  
BROOKHAVEN NATIONAL LABORATORY  
Upton, New York 11973*

*ES&F Division Technical Note # 162*

**Soil Activation in various AGS/RHIC sites by Monte Carlo simulation**

*Kin Yip*

*November 14, 2005*

**Soil Activation in various AGS/RHIC sites by Monte Carlo  
simulation**

*Nov. 14, 2005*

*Kin Yip*

*Experimental Support and Facilities Division*

*Collider-Accelerator Department*

As part of the Brookhaven National Laboratory Land Use and Institutional Controls, information on each of the BNL contaminated sites and facilities is needed. For the Collider-Accelerator Department, E. Lessard and the author are responsible for providing historical soil activation information on various AGS and RHIC sites which have had substantial running records and proton irradiation. This is an ongoing and continuous process. The author describes here what has been achieved for various AGS beam lines so far.

The author has used the simulation package MCNPX (<http://mcnpx.lanl.gov>) maintained by Los Alamos National Laboratory to compute the neutron fluxes (essentially the only concern for soil activation) in the soil deep under various sites. The simulation computational process is very CPU intensive. The author has installed and configured the MCNPX simulation to run in the *parallel* mode using MPICH (<http://www-unix.mcs.anl.gov/mpi/mpich>) for a few years. This mainly works in the godzilla cluster (in C-AD) but occasionally also in other computers such as rlnxsp.rcf.bnl.gov in RCF. The author has also maintained and updated the software for other people such as D. Beavis to make use of.

The soil activation (in the units of pCi/L – pico-Curie per liter) is given by:

$$\frac{\phi}{\lambda} \times N_I \times N_p \quad \text{[Equation 1]}$$

where  $N_I$  is the number of atoms per interaction (which is 0.075 for  $^3\text{H}$  and 0.02 for  $^{22}\text{Na}$ ),  $N_p$  the number of protons which have been recorded by the Main Control Room and  $\lambda$  the interaction length, typically 40 cm.


The author is mostly responsible for producing the neutron flux contours. Therefore, the plots below show just these flux contours at various depths for those sites which we have completed the MCNPX simulation. The flux contours are always in the unit of  $\text{cm}^{-2}$  per incident proton.

Recently (since version 2.5.f), MCNPX has finally provided their stand-alone mesh tally plotting capability without additional external software. This new feature has made it more convenient for the author to produce detailed contour plots such as those shown in Figure 7 and Figure 8. The geometry of the latter two plots were mainly set up by the author's 2005 summer student N. Sapanski.

The original B Target has seldom been used. There is only one beam dump for the entire B line and this B Beam Dump is just behind the B' Target. Therefore, only one simulation which has included both B' Target and Beam Dump is needed.



**Figure 1: Soil Activation (in fluxes  $\text{cm}^2/\text{proton}$ ) at East Side of the g-2 Target Blockhouse.**



**Figure 2: Soil Activation (in fluxes  $\text{cm}^2/\text{proton}$ ) below the g-2 Target Blockhouse.**




Figure 3: Cross section view of soil activation area below the A Dump.




Figure 4: Plan view of soil activation area beneath the A Dump.




Figure 5: Cross section view of soil activation area below the B' Target and Dump.




Figure 6: Plan view of soil activation area beneath the B' Target and Dump.






Figure 7: Cross section view of soil activation area below the C Target.




Figure 8: Plan view of soil activation area beneath the C Target.