

# BNL-99187-2013-TECH C-A/AP/33;BNL-99187-2013-IR

# Coupling Matrices in AGS with different types of partial snakes

E. D. Courant

November 2000

Collider Accelerator Department Brookhaven National Laboratory

## **U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

C-A/AP/33 December 2000

# **Coupling Matrices in AGS with different types of partial snakes**

E. D. Courant



Collider-Accelerator Department Brookhaven National Laboratory Upton, NY 11973

# Coupling Matrices in AGS with different types of partial snakes

#### E D Courant December 5, 2000

The solenoid partial snake in the AGS produces an irreducible amount of coupling between the horizontal and vertical betatron oscillation modes. Therefore alternate partial-snake schemes have been proposed: (a) a helix (with correction dipoles at its ends)<sup>1</sup>; (b) a sequence of four deflecting magnets with field orientations at various angles<sup>2</sup>.

Matrices for the AGS with a partial snake have been computed using the MATHEMATICA computer program. The helical matrices are evaluated using an algorithm previously formulated<sup>3</sup>, which takes into account the solenoidal fields and nonlinearities dictated by Maxwell's equations in a helical magnet, and the end effects due to the orbit excursions away from the axis of the helix. The partial snake is assumed to be in the C10 straight section of the AGS, and the matrices, with or without snakes, are evaluated at the upstream end of C10.We neglect sextupole and other multipole fields in the dipoles and in the helical magnet.

In these calculations we assume the following parameters:

AGS without snakes:  $v_x = 8.698$ ,  $v_y = 8.749$  (about .05 units away from coupling resonance).

Strength of partial snakes (solenoidal, helical, or dipole) adjusted to make 5% snakes, e.g. 9° spin rotators.

Solenoid: length 2.5 m; field strength ramping proportional to main field: 0.167 T at injection (1.5 GeV) to 1.825 T at  $\gamma G = 46.5$  (transfer to RHIC).

Helix as in ref. 1: Helix length 1.5 m, compensating magnets 0.3 m each. Helix field ramping from 1.25T at injection to 1.61 T at transfer.

4-magnet snake: magnet lengths 0.41, 0.66, 0.66, 0.41 m; field direction (from vertical)  $-60^{\circ}$ , 147°, 33°, -120°; field ramping from 1.5T (injection) to 1.94T (transfer)

In each case the coupling strength is measured by two quantities: the "minimum tune separation", and the "coupling angle". Both of these are derived from the 4x4 submatrix of the 6x6 orbit matrix given below. If the (symplectic) 4x4 matrix is of the form

М	п
т	N

where M, N, m, n are 2x2 submatrices, then the coupling arises from the off-diagonal submatrices m and n. The normal modes <u>u</u> and <u>v</u> are related to the geometric coordinates <u>x</u> and <u>y</u> by

 $\underline{\mathbf{x}} = \underline{\mathbf{u}} \cos \varphi + \mathbf{D} \, \underline{\mathbf{v}} \sin \varphi; \ \underline{\mathbf{y}} = -\mathbf{D'} \, \underline{\mathbf{u}} \sin \varphi + \underline{\mathbf{v}} \cos \varphi$ 

where  $\varphi$  may be called the "coupling angle" and D is a certain symplectic 2x2 matrix. The angle  $\varphi$  and the matrix D can be obtained from the matrices *M*, *m*, *n*, *N*. Full coupling corresponds to  $\varphi = 45$  degrees

The "minimum tune separation", i.e. the tune splitting if the uncoupled tunes are exactly at resonance, is

$$\Delta v = (Det(n+m'))^{1/2} / (2\pi).$$

4

<sup>&</sup>lt;sup>1</sup> T Roser et al, Spin Note No. 72, 1998

<sup>&</sup>lt;sup>2</sup> E D Courant, Spin Note No. 83, 1999

<sup>&</sup>lt;sup>3</sup> E D Courant, Spin Note No. 4 (1996)

where m' is the symplectic conjugate of m.

This quantity  $\Delta v$  is a better measure of the coupling strength than the coupling angle, since it is independent of how close one is to resonance, while the coupling angle will always be 45° exactly at resonance no matter how weak the coupling is, as long as it is not zero.

Below we tabulate the matrices at C10 for AGS injection energy (1.5 GeV), for the energies of the important intrinsic resonances  $0+v_y$  and  $36-v_y$ , and for the transfer energy  $\gamma G = 46.5$ . We do this for the 5% solenoid snake, the helical snake, and the 4-dipole snake described above. We also show the orbit excursions inside the snake for the helical and 4-dipole case. (in the solenoid case the orbit displacement is zero). The matrices are similar to those previously obtained by N. Tsoupas and M. Okamura.

Note that the coupling parameters with a solenoid are independent of energy. The coupling effects in the helix and 4-dipole cases decrease approximately with the square of the energy (not quite, because the helical or dipole fields have to increase a bit with energy because  $(1+\gamma G)$  increases a little faster than  $\beta\gamma$ ). At injection energy the coupling in the 4-dipole case is distinctly stronger than for the solenoid (this is due to the focusing effect at the ends of the dipoles, where the excursions are quite large – up to 5 cm). The helix has an order of magnitude weaker coupling, and exhibits less coupling than the solenoid even at injection energy. At the higher energies -  $\gamma G = 27$  to 46 – the coupling is very small in both the helix and dipole cases.

We see that the helix is definitely preferable – if it is practicable.

-1.31324	-11.2107	0	0	0	2.08681
0.167845	0.671359	0	0	0	-0.318092
0	0	1.57424	-19.7498	0	0
0	0	0.177118	-1.58682	0	0
-0.06747	-2.16502	0	0	1.	-11.5419
0	0	0	0	0	1.
T					

AGS without snakes

Tunes {8.698, 8.749}

Matrix with solenoid snake

-1.31093	-11.1934	0.0442375	-0.554929	0	2.08299
0.168355	0.675285	0.00497101	-0.0444856	0	-0.318932
0.0368755	0.31486	1.57265	-19.7279	0	-0.0585927
-0.0047357	-0.0189952	0.176721	-1.58147	0	0.0089713
-0.06747	-2.16502	0	0	1.	-11.5419
0	0	0	0	0	1.

Minimum tune split = 0.0144566; coupling angle 8.08129 Tunes {8.69744, 8.75032}

Injection: 
$$\gamma G = 4.65904$$

:

Helical Snake:

Matrix with helical snake

-1.23141	-10.5775	-0.0130737	0.16348	0	1.8941
0.220509	1.08209	-0.00114092	0.00848222	0	-0.403393
-0.0106962	-0.0918811	1.50175	-18.7783	0	0.0163952
0.00194014	0.0095958	0.128475	-0.940651	0	-0.00375087
-0.0790867	-2.21748	0.000304394	-0.00384412	1.	-11.5189
0	0	0	0	0	1.

Minimum tune split = 0.00448604; coupling angle 2.26851 tunes {8.73802, 8.79535}



#### 4-Dipole Snake Matrix with 4-dipole snake

Matrix with + dipole shake						
-1.16851	-10.0816	-0.0159327	0.18963	0	1.84956	
0.243827	1.2475	-0.0351435	0.44673	0	-0.443225	
0.00648536	0.0692911	1.51749	-19.0392	0	-0.00780593	
0.0301393	0.252109	0.12114	-0.860595	0	-0.0469726	
-0.0668694	-2.1598	-0.00172722	0.0227037	1.	-11.5333	
0	0	0	0	0	1.	

Minimum tune split = 0.0539474; Coupling angle = 24.7769 tunes {8.74383, 8.81666}



## $\gamma G$ at resonance $0+\nu = 8.749$ :

Helical Snake

Matrix with helical snake,

		,				
	-1.2856	-10.9968	-0.00454085	0.0569063	0	2.00705
	0.185674	0.810388	-0.000473248	0.00404146	0	-0.347115
	-0.00376408	-0.0321978	1.54975	-19.4215	0	0.00586524
	0.000546455	0.00239514	0.160639	-1.36787	0	-0.00106141
	-0.0735955	-2.1907	0.0000610821	-0.000772655	1.	-11.5299
Γ	0	0	0	0	0	1.

Minimum tune split = 0.00151209; coupling angle 0.828196 tunes {8.71181, 8.7645}



#### 4- dipole snake

Matrix with 4-dipole snake

-1.27088	-10.8553	-0.00534599	0.0635777	0	2.01906
0.193523	0.866087	-0.0118967	0.151276	0	-0.360449
0.00252978	0.0259674	1.54818	-19.4481	0	-0.00342083
0.0106925	0.0890585	0.158157	-1.34081	0	-0.0168911
-0.0673455	-2.16395	-0.000337626	0.00443806	1.	-11.5389
0	0	0	0	0	1.

Minimum tune split = 0.0186456; Coupling angle = 10.4723 tunes {8.7158, 8.76827}



 $\gamma G$  at resonance 36- $\nu = 27.251$  :

Helical snake

Matrix with helical snake,

-1.30994	-11.1852	-0.000547999	0.00687408	0	2.06946
0.169973	0.687949	-0.00006112	0.000544763	0	-0.321571
-0.000456803	-0.00390049	1.57132	-19.7106	0	0.000721203
0.0000593128	0.000240219	0.175151	-1.56068	0	-0.000113869
-0.0694881	-2.17314	0.0000025	-0.000032135	1.	-11.538
0	0	0	0	0	1.

Minimum tune split = 0.000179397; coupling angle 0.102084 tunes {8.69967, 8.75085}



#### 4-dipole snake

Matrix with 4-dipole snake

-1.31693	-11.2033	-0.000635388	0.00755368	0	2.09529
0.170907	0.694581	-0.0014196	0.018054	0	-0.32315
0.000319777	0.00323311	1.56192	-19.6312	0	-0.000461704
0.00130226	0.0108269	0.174855	-1.55745	0	-0.00207864
-0.0674648	-2.16498	-0.00001387	0.000182324	1.	-11.5415
0	0	0	0	0	1.

Minimum tune split = 0.00224526; Coupling angle = 1.28866 tunes {8.69961, 8.75038}



### $\gamma$ G at transfer to RHIC = 46.5 :

Helical snake

Matrix with helical snake,

-1.31207	-11.2017	-0.000194072	0.00243463	0	2.07779
0.168598	0.677229	-0.000021768	0.000194669	0	-0.319324
-0.000161853	-0.0013818	1.57321	-19.7359	0	0.000256213
0.0000208027	0.0000835805	0.176422	-1.57757	0	-0.0000397
-0.0686653	-2.16981	0.0000005	-0.00000677	1.	-11.5396
0	0	0	0	0	1.

Minimum tune split = 0.0000634312; coupling angle 0.0362184 tunes {8.69859, 8.74965}



4-dipole snake

.

#### Matrix with 4-dipole snake

	r · · · · ·				
-1.32096	-11.2337	-0.000224722	0.00267147	0	2.10196
0.168928	0.679575	-0.000502253	0.00638759	0	-0.319882
0.000113691	0.00114803	1.56312	-19.6472	0	-0.00016606
0.000461554	0.00383676	0.176317	-1.57643	0	-0.000738246
-0.0674689	-2.16501	-0.0000029	0.0000383578	1.	-11.5417
0	0	0	0	0	1.

Minimum tune split = 0.000794997; Coupling angle = 0.455641 tunes {8.69804, 8.74894}

