

## Resonance search with pp

D. C. Peaslee

April 1984

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

RESONANCE SEARCH WITH  $\bar{p}p$

D. C. Peaslee

Department of Physics and Astronomy  
University of Maryland  
College Park, Maryland 20742

Physics Publication 84-141

April, 1984

Narrow resonances in  $\bar{p}p$  reactions have been reported for some time. They have a curious, now-you-see-it-now-you-don't character: the one at 2.02 GeV, for example, being alternately discovered, disproved and rediscovered several times (see Appendix). Here we pursue the hypothesis that such resonances are real and reflect underlying mesonic states of the main sequence, that their narrowness is peculiar to the  $\bar{p}p$  channel and may vary with the conditions of observation, and that this selective narrowness may make  $\bar{p}p$  interaction useful to explore meson sequences up to  $m \sim 4$  GeV: that is, up to incident  $\bar{p}$  momenta of 7-8 GeV/c.

## 1. Review of $\bar{p}p$ resonance information.

It is perhaps not generally appreciated how substantial a contribution  $\bar{N}N$  studies have already made to meson spectroscopy. In the region  $m > 2$  GeV information about meson sequences comes mainly from two sources: two-body decays with special signatures like  $\pi^0\pi^0$  or  $\bar{K}K$ , and reactions involving  $\bar{N}N$ .

The first category has provided evidence on the leading trajectory with  $J = L + 1$  on the  $\bar{q}q$  model:  $J^P = 4^{+(1,2)}$ ,  $5^{-(3)}$ ,  $6^{+(4)}$ ,  $7^{-(5)}$ . These states define a good straight line of  $m^2$  vs.  $J$ , with  $dm^2/dJ \sim 1.14$  GeV<sup>2</sup> and an intercept of  $J \sim 1/2$  at  $m^2 = 0$ .

Although one of these states, the  $h(2040)$ , has been seen in the  $\bar{N}N$  channel from  $\pi^-p$  at 18 GeV<sup>(6)</sup>, the great majority of resonances inferred from  $\bar{N}N$  studies seem to lie on other trajectories, most notably the fundamental  $J = L - 1$  sequence. If we define this sequence as starting with the  $\rho^-(1600)$  and the  $X(1850)$  for  $J^P = 1^-, 2^+$ , the states in Table I define a good straight line with  $dm^2/dJ = 0.97$  GeV<sup>2</sup> and an  $m^2 = 0$  intercept around  $J = -3/2$ .

Table I. Candidates for  $J = L - 1$  Trajectory

<u><math>J^P</math></u>	<u><math>M(\text{GeV})</math></u>	<u>Reference</u>
$1^-$	1.55	2
$2^+$	1.83	7
$3^-$	2.11	6,8,9
$4^+$	2.36	6,8,9,10
$5^-$	2.48	6,8
$6^+$	2.71	4

The values in Table I are plotted in Fig. 1 along with the leading  $J = L + 1$  trajectory. If the linearity persists to higher values, the trajectories will extrapolate to cross at  $J \sim 12$ . If the difference in slope is ascribed to spin-orbit coupling  $\delta m^2 = C_1(\underline{\Sigma} \cdot \underline{L})$ , then  $C_1 \sim 8.5 \times 10^{-2} \text{ GeV}^2$ .

## 2. Possible higher resonances.

Such evidence as exists regarding higher resonances comes entirely from involvement of  $\bar{N}N$  channels. The reaction  $\bar{p}p \rightarrow 6$  prongs at 6.4 GeV<sup>(11)</sup> indicates a meson at 3.04 GeV and  $\Gamma \sim 200$  MeV, and another candidate at 3.42 GeV that might be narrower. The reaction  $\bar{p}n \rightarrow X\pi^-$  from  $\bar{p}d$  at 5.5 GeV<sup>(12)</sup> indicated narrow resonances at  $m_X = 2.85$  and 3.05 GeV. And in  $\pi^-p \rightarrow \bar{p}p\pi^-X$  at 16 GeV<sup>(13)</sup> a resonance was perceived at 2.95 GeV at a width compatible with the experimental resolution of order 20 MeV. Less outstanding but also visible in the same data summary graph, reproduced as Fig. 2, are peaks at 3.16 and 3.45 GeV. As with



the 2.02 GeV example we assume the later obliteration of these effects<sup>(14)</sup> to result from changes in the experimental setup.

A schematic summary of this information is as follows: possible states at  $(2.90)^2$ ,  $(3.04)^2$ , 3.16 and  $(3.44)^2$  GeV, where  $( )^2$  indicates presence in two separate experiments. Comparison with Fig. 1 suggests possible correspondences as in Table II.

Table II. Trajectory Assignments for Higher States

<u>m(GeV)</u>	<u>J=L+L</u>	<u>J=L-1</u>
2.90	$8^+$	$7^-$
3.04	---	$8^+$
3.16	$9^-$	$9^-$
3.44	$11^-$	$11^-$

Of course this is highly speculative but indicates that mesonic states of very high spin may already have been detected.

### 3. Narrowness of $\bar{N}N$ States.

Some understanding of the narrowness of  $\bar{N}N$  resonances is desirable if they are to be relied on when searching for meson states. Presumably this has to do with the existence of  $\bar{N}N$  systems as definable entities not the same as  $\bar{q}q$  systems but strongly related to them. A  $\bar{q}q$  state of given  $J^{PC}$  may be

connected with a number of different  $\bar{N}N$  configurations, which serve as doorway states. Although reactions proceed mainly through the  $\bar{q}q$  core, they also have direct contributions from the  $\bar{N}N$  states. The interference effects between these two reaction routes may depend sensitively on the particular channel being observed - or in some cases even the angular range of the detector because of different  $t$  dependence of the interfering amplitudes.

If the level density of related  $\bar{N}N$  states were  $10^2-10^3$  times that of the  $\bar{q}q$  state in question, the situation would resemble  $n\gamma$  capture on heavy nuclei. The particle widths of the many individual states form an envelope, the strength function, which is characteristic of a single particle in an optical potential: the present analogy nominating  $\bar{N}N$  as the individual states and  $\bar{q}q$  as the single-particle system. The difference is that there the level density of  $\bar{N}N$  states is probably only  $\sim 10\times$  that for  $\bar{q}q$ , so that the "envelope" exhibits granularity.

Qualitatively, this picture can explain the  $\bar{N}N$  observations. In an occasional experimental setup the reaction through a mesonic state will be dominated by one or a few  $\bar{N}N$  channels, producing a peak with characteristic width of order (a few)  $\times 10$  MeV. Because this effect depends sensitively on interference, it will not be seen equally under different conditions; it may sometimes be entirely washed out, or in an extreme case might shift position slightly by moving to a different constituent  $\bar{N}N$  state.

In this way we can perhaps understand why a narrow state at 2.02 GeV should appear<sup>(15)</sup> in  $\pi^-p \rightarrow p\bar{p}p\pi^-$  but not<sup>(16)</sup> in  $\pi^-p \rightarrow p\bar{n}p$  nor<sup>(17)</sup> in  $\pi^+p \rightarrow \Delta^{++}p^-p$  and re-appear<sup>(18)</sup> in  $\bar{p}p \rightarrow \bar{p}n \pi^+\pi^+\pi^-$  and  $\bar{n}p\pi^+\pi^-\pi^-$  and also<sup>(19)</sup> in  $\gamma p \rightarrow p\bar{p}p$ . All of these observations had selection criteria based on fast particles in the final state. Of particular interest by comparison was  $\pi^+n \rightarrow \bar{p}pp$  at 10 GeV<sup>(20)</sup> where a sufficiently comprehensive measurement was made to determine  $J^P = 4^+$ . In such a complete observation one would expect a more or less unselected sum over all  $\bar{N}N$  channels; and indeed the resonance appears to have  $\Gamma \gtrsim 100$  MeV, appropriate to the h(2040).

Parenthetically one may remark that in all cases where a narrow state at 2.02 GeV was seen, the data with sufficient statistics<sup>(15,18)</sup> showed the presence of similar states at  $\sim 1.93$  and  $\sim 2.18$  GeV, although they were not always claimed. According to the general arguments given here, these also represent mesonic states.

Of course the model of  $\bar{N}N$  interference with  $\bar{q}q$  states should be articulated further for specific instances. Even in this qualitative form, however, it may serve to encourage reliance on narrow  $\bar{N}N$  states as a search mechanism for locating mesonic states of high mass. Candidates so surveyed could then be examined more thoroughly, perhaps with other channels, to arrive at  $J^{PC}$  assignments.



## Appendix

The chronology of the narrow 2.02 GeV state shows an amusing oscillation, displayed below:

Pub. Date	Result	Reaction	Ref.
July '77	+	$\pi^- p \rightarrow p\bar{p}p\pi^- X$	15
Feb. '79	-	$\pi^- p \rightarrow p\bar{p}n$	16
June '79	+	$ep \rightarrow ep\bar{p}p$	21
Apr. '80	-	$\pi^+ p \rightarrow \Delta^{++}\bar{p}p$	17
Nov. '80	-	$\pi^- p \rightarrow p\bar{p}p\pi^-$	22
Mar. '83	+	$\bar{p}p \rightarrow (\bar{p}n\pi^+/p\bar{n}\pi^-)\pi^+\pi^-$	18
Dec. '83	+	$\gamma p \rightarrow p\bar{p}p$	19

In this account we have omitted two measurements that were ambiguous: one that could claim with low statistics to see no effect<sup>(23)</sup>, but which had a suggestive accumulation of events at 2.02 GeV; and one that saw the peak<sup>(20)</sup> in  $\bar{p}p$  at 2.02 GeV and measured  $J^P = 4^+$  but did not find it narrow. Also omitted are the negative experiments that were sufficiently "dirty" to permit the objection that they might have washed out any narrow resonances - one because of its beam<sup>(24)</sup> and one because of its target<sup>(25)</sup>.

### References

1. W. D. Apel et al., Phys. Lett. 57B, 398 (1975); W. Blum et al., ibid., 403.
2. Particle Data Group, Phys. Lett. 111B, 1 (1982).
3. B. Alper et al., Phys. Lett. 94B, 422 (1980).
4. W. Cleland et al., Nucl. Phys. B208, 228 (1982); F. Binon et al., CERN EP/83-98 (1983).
5. D. L. Denney, Phys. Rev. D 28, 2726 (1983).
6. M. Rozanska et al., Nucl. Phys. B162, 505 (1980).
7. G. Costa et al., Nucl. Phys. B175, 402 (1980); N. M. Cason et al., Phys. Rev. Lett. 48, 1316 (1982).
8. A. A. Carter et al., Phys. Lett. 67B, 117 (1977).
9. C. Evangelista et al., Nucl. Phys. B153, 253 (1979).
10. R. S. Dulude et al., Phys. Lett. 79B, 335 (1978).
11. G. Alexander et al., Phys. Rev. Lett. 25, 63 (1970).
12. H. Braun et al., Phys. Lett. 60B, 481 (1976).
13. C. Evangelista et al., Phys. Lett. 72B, 139 (1977).
14. T. A. Armstrong et al., Phys. Lett. 85B, 304 (1979).
15. P. Benkhieri et al., Phys. Lett. 68B, 483 (1977).
16. P. Benkhieri et al., Phys. Lett. 81B, 380 (1979).
17. R. M. Bionta et al., Phys. Rev. Lett. 44, 909 (1980).
18. F. Azooz et al., Phys. Lett. 122B, 471 (1983).
19. J. Bodenkamp et al., Phys. Lett. 133B, 275 (1983).
20. J. W. Lamsa et al., Phys. Rev. D 26, 1769 (1982).
21. B. G. Gibbard et al., Phys. Rev. Lett. 42, 1593 (1979).
22. S. U. Chung et al., Phys. Rev. Lett. 45, 1611 (1980).

23. S. Kooijman et al., Phys. Rev. Lett. 45, 316 (1980).
24. A. D. J. Banks et al., Phys. Lett. 100B, 191 (1981).
25. A. S. Carroll et al., Phys. Rev. Lett. 44, 1572 (1980).

Figure Captions

Figure 1 - Plots of  $m^2$  vs.  $J$  for postulated leading trajectories

$J = L + L$  and  $J = L - 1$ .

Figure 2 - Reproduction of Fig. 3 from Ref. 13 on  $\pi^-p$  at 16

GeV/c.






Fig. 1




Fig. 2