

Luminosity Formulae

A. G. Ruggiero

November 1983

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

LUMINOSITY FORMULAE

A.G. Ruggiero

(BNL, November 20, 1983)

~~LUMINOSITY CALCULATIONS~~

~~Ring~~

A. Both Colliding Beams are bunched

$$L = N_1 N_2 f_{\text{encounter}} F$$

$$F = \frac{2}{(2\pi)^{3/2} (\sigma_{l_1}^2 + \sigma_{l_2}^2)^{1/2}} \int_{-\infty}^{+\infty} \frac{ds}{(\sigma_{x_1}^2 + \sigma_{x_2}^2)^{1/2} (\sigma_{z_1}^2 + \sigma_{z_2}^2)^{1/2}} \times$$

$$\times \exp \left\{ -2s^2 \left(\frac{1}{\sigma_{l_1}^2 + \sigma_{l_2}^2} + \frac{\alpha^2/4}{\sigma_{x_1}^2 + \sigma_{x_2}^2} \right) \right\} \quad (1)$$

$$= \int_{-\infty}^{+\infty} G(s) ds$$

$$\sigma^2 = \frac{\epsilon \beta(s)}{6\pi} \quad \text{H and V, zero dispersion}$$

ϵ emittance for 95% of beam with bi-gaussian distribution

$$\beta(s) = \beta^* + \frac{s^2}{\beta^*}$$

N_1, N_2 , number of particles per bunch

σ_e , rms bunch length

α , total crossing angle assumed in the x -plane.

If both beams are made of B bunches each

$$f_{\text{encounter}} = B \cdot \text{revolution}$$

B. One Bunched Beams colliding with an Unbunched Beams

$$L = N_U N_B \text{ encounter } F'$$

$$F' = \frac{1}{2\pi^2 R} \int_{-\infty}^{+\infty} \frac{ds}{(\sigma_{xU}^2 + \sigma_{xB}^2)^{1/2} (\sigma_{zU}^2 + \sigma_{zB}^2)^{1/2}} \times \exp \left\{ - \frac{\alpha^2 s^2}{2(\sigma_{xU}^2 + \sigma_{xB}^2)} \right\} \quad (2)$$

$$= \int_{-\infty}^{+\infty} G(s) ds$$

- $2\pi R$, circumference of reference orbit ~~assumed to~~
 be assumed to be the same for both beams
- N_U , total number of particles in the unbunched beam
- N_B , no. of particles per bunch in the bunched beam
- B , no. of bunches in the bunched beam

$$\text{Encounter} = B \text{ revolution}$$

Extreme Cases

A. Both Colliding Beams are bunched

Long Bundles : $\sigma_e \Rightarrow 2\sigma_x / \alpha$

σ_x and σ_z are constant ~~is~~ over interaction region

Define effective cross-sections and length

$$\frac{1}{\sigma_x^2} = \frac{\sigma_{x1}^2 + \sigma_{x2}^2}{2}$$

$$\frac{1}{\sigma_z^2} = \frac{\sigma_{z1}^2 + \sigma_{z2}^2}{2}$$

$$\frac{1}{\sigma_e^2} = \frac{\sigma_{e1}^2 + \sigma_{e2}^2}{2}$$

We have

$$L = \frac{N_1 N_2 B \text{ revolution}}{2\pi \alpha \bar{\sigma}_e \bar{\sigma}_z} \tag{3}$$

having assumed that both beams have the same number of bunches B -

B. One Bunched Beam colliding with an Unbunched Beam

σ_x and σ_z are constant over interaction region

Define effective cross-sections

$$\overline{\sigma}_x^2 = \frac{\sigma_{xU}^2 + \sigma_{xB}^2}{2}$$

$$\overline{\sigma}_z^2 = \frac{\sigma_{zU}^2 + \sigma_{zB}^2}{2}$$

We have

$$L = \frac{N_U N_B B \text{ revolutions}}{2\pi \alpha (\sqrt{\pi} R) \overline{\sigma}_z} \quad (4)$$

Comment

For crossing at a large angle and for long bunches or coasting beams the equations shown above are correct as long

$$\sqrt{2} \frac{\overline{\sigma_x}}{\alpha} \ll l \quad (5)$$

where l is half of the length of the vacuum chamber shared by both beams unshielded from each other.

Eq. (3) and (4) are derived assuming eq. (5) holds.

If relation (5) does not hold the integration limits in eq. (1) and (2) are to be replaced by $-l$ and $+l$ respectively instead of $-\infty$ and $+\infty$.

(6)

C. Short Bundles Colliding Head-on ~~Case~~

Extreme Case from eq. (1) with $\alpha = 0$ -

$\bar{\sigma}_x$ and $\bar{\sigma}_z$ are constant ~~of~~ over interaction region -

Assume $\bar{\sigma}_e \ll l$

Then we have with the usual notation

$$L = \frac{N_1 N_2 B \text{ revolution}}{4\pi \bar{\sigma}_x \bar{\sigma}_z}$$

having again assumed that both beams have the same number of bundles B -

References

Lloyd Smith
A.G. Ruggiero

, PEP - Note - 20 , April 27, 1972
, FN - 271 , December 1974