



Brookhaven
National Laboratory

BNL-101579-2014-TECH

RHIC/PG/36;BNL-101579-2013-IR

Multiple Coulomb Scattering a) Tandem Exit b) Booster Exit

G. R. Young

February 1984

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MULTIPLE COULOMB SCATTERING AND EMITTANCE
GROWTH IN STRIPPER FOILS

- a) Tandem Exit
- b) Booster Exit

Glenn Young

February 23, 1984

Brookhaven National Laboratory

is small,

Rewriting the equation for $\langle \Theta^2 \rangle$, we have

$$\sqrt{\langle \Theta^2 \rangle} \equiv \Theta_{\text{plane}}^{\text{rms}} = \sqrt{2\pi \frac{\rho N_A}{A} \pm \frac{Z^2}{2} \ln(210 Z^{-1/3})} \quad \frac{2Z}{\rho} \frac{e^2}{hc} \frac{hc}{\beta c}$$

$(\beta = v/c)$

We can approximate $\frac{X_0^{-1}}{4 \alpha r_0^2 N_A} \sim \frac{Z^2 \ln(210 Z^{-1/3})}{A}$

and err by only 10% or less for our stripping foils, except for carbon where the neglect of the second coulomb log gives ~16% error for $(\Theta_{\text{plane}}^{\text{rms}})^2$. This gives a convenient formula in terms of X_0 , which is tabulated; the correction is included by multiplying by $(1+\delta)$ where δ is evaluated for each material.

For carbon, $\delta = -0.079$
 copper, $\delta = 0.00193$
 tantalum, $\delta = 0.0478$

This then works out to

$$\Theta_{\text{plane}}^{\text{rms}} = (1+\delta) Z, \frac{14.98 \text{ MeV}/c}{\rho \beta} \sqrt{\frac{\Delta}{X_0}}$$

where Δ is the scatterer thickness in g/cm^2 and X_0 is in g/cm^2 . X_0 is tabulated in Tsai's article, or see the Particle Data Properties booklet, $X_0 = 42.70 \text{ g}/\text{cm}^2$ carbon; $6.82 \text{ g}/\text{cm}^2$ tantalum
 $12.86 \text{ g}/\text{cm}^2$ copper

2/23/84

In the following graphs, values of $\theta_{\text{plane}}^{\text{rms}}$ are given for C, S, Cu, I and Au ions for the following cases.

- (a) Tandem exit energy/nucleon expected for 2 stage operation
5-20 $\mu\text{g}/\text{cm}^2$ carbon stripper
- (b) booster exit 300-500 MeV/nucleon i.e. matching to
performance expected for gold
10-100 mg/cm^2 copper stripper

Emitance growth can be found from

$\varepsilon = \beta (\theta_0^2 + \theta_{\text{plane}}^{\text{rms}^2})$, where $\varepsilon = \beta \theta_0^2$ is the emittance prior to passing through the stripper. The last graphs then give values of the β function required to limit the emittance growth at the tandem exit to $0.5 \pi \text{ mm mrad}$, and at the booster exit to $1.0 \pi \text{ mm mrad}$. Carbon is fully stripped after the tandem exit stripper. The curves for ^{32}S and ^{63}Cu at the booster exit are included for completeness only. The stripper thicknesses listed are overkill by at least a factor of 100 or more.

As usual, gold yields the most stringent requirements.

At the tandem exit, 20 $\mu\text{g}/\text{cm}^2$ carbon @ 1 MeV/A requires $\beta^* = 14 \text{ cm}$. At the booster exit, 367 MeV/A thru a 70 mg/cm^2 Copper stripper requires $\beta^* = 2.02 \text{ m}$. This line can handle all lighter beams, as they use thinner stripping foils.

NOTE ADDED

Another formula for multiple scattering of heavy ions is given by Green (U. Washington, Seattle, Annual Report 1975). It gives somewhat more optimistic numbers than the one from Jackson.

The formula is

$$\Theta_{1/2} = K \frac{Z Z_1 Z_2}{E_1} \tau^n \quad \left(\text{Divide by } 1.175 \text{ to get } \sigma_{\text{plane}}^{\text{rms}} \right)$$

where $K = 1.92 \cdot 10^{-2}$ mrad, $Z_1 = Z$ of ion, $Z_2 = Z$ of foil,
 $Z = (Z_1^{2/3} + Z_2^{2/3})^{3/2}$ (this appears from calculating an effective screening radius),
 E_1 is the total kinetic energy of the ion, and τ and n are given by

$$\tau = \frac{41.5 \delta}{Z^2 A}$$

$\delta =$ foil thickness in $\mu\text{g}/\text{cm}^2$, $A =$ foil Atomic weight

and $n = N^{-.115} - .115$ where $N = \ln(1.03 + \tau)$. Needless to

say, this is a phenomenological formula.

PAST EXPERIENCE AT OAKRIDGE IS THAT THIS GIVES TOO SMALL VALUES BY ~ FACTOR OF TWO

To give some values to compare, consider ¹⁹⁷Au at the tandem and booster exits, passing thru 20 $\mu\text{g}/\text{cm}^2$ C and 70 $\mu\text{g}/\text{cm}^2$ Cu Stippers, respectively. A comparison of the two formulae results is below.

	Tandem				Booster			
	.5 MeV/A	1 MeV/A	1.5 MeV/A	300	350	400	450	MeV/A
Jackson's	3.782	1.894	1.263	1.40	1.86	2.34	2.87	
Green's	.978	.489	.326	3.77	5.14	6.71	8.50	
Booster	300	350	400	450	300	350	400	450
Jackson's	.844	.733	.654	.590	1.40	1.86	2.34	2.87
Green's	.515	.441	.386	.343	3.77	5.14	6.71	8.50

$\Theta_{\text{plane}}^{\text{rms}}$ (mrad)

β^* value needed

46
3/1/84

CONCLUSION

- (1) Some attention has to be paid to the transport between the tandem high energy end and the stripper foil. A quad doublet has to be used to focus x and y to a small spot. β^* values somewhere in the range of 0.14 m (Jackson formula) to 2 m (Greene) formula are needed.

A measurement of emittance after the high energy stripper for the heavy beams is needed. We want a growth of less than 0.5π mm mrad so the injection into the booster works.

- (2) Stripping at the booster exit will require β^* of 2 m or larger, but again in both x and y . Again a quad doublet before the stripper is needed.

- (3) Better (smaller) values of scattering angles due to multiple Coulomb scattering could be had by using gold foils. Less thickness is needed to strip to the equilibrium charge state distribution due to the density effect in stripping.

For the tandem exit, this probably does not work as the thickness required ($10 \mu\text{g}/\text{cm}^2$) is mechanically quite fragile.

For the booster exit, foils of $20-30 \text{ mg}/\text{cm}^2$ are needed, which are quite rugged (0.02 mm thick). As the β^* required varies as the thickness (inversely) the β^* needed after the booster could be relaxed by a factor of two in both planes.

$\theta_{\text{plane}}^{\text{rms}}$ Mult Scatt5Tandem ExitCarbon Stripper (5, 10, 20 $\mu\text{g}/\text{cm}^2$)

(MeV/A)

 $\theta_{\text{plane}}^{\text{rms}}$ (mrad)

Ion	E/A	P/A	P	β	5 $\mu\text{g}/\text{cm}^2$	10 $\mu\text{g}/\text{cm}^2$	20 $\mu\text{g}/\text{cm}^2$
^{12}C (Z=6)	6.5	110.2	1322.8	.1175	0.182	0.258	0.364
	7	114.4	1372.9	.1219	0.169	0.239	0.339
	7.5	118.4	1421.3	.1261	0.158	0.224	0.316
	8	122.3	1468.1	.1302	0.148	0.210	0.296
	8.5	126.1	1513.5	.1342	0.139	0.197	0.279
^{32}S (Z=16)	3.5	80.8	2586.4	.0864	0.338	0.478	0.676
	4	86.4	2765.4	.0924	0.296	0.418	0.591
	4.5	91.7	2933.5	.0979	0.263	0.372	0.526
	5	96.6	3092.6	.1032	0.237	0.335	0.473
	5.5	101.4	3244.0	.1082	0.215	0.304	0.430
^{63}Cu (Z=29)	2	61.1	3847.6	.0654	0.544	0.769	1.088
	2.5	68.3	4302.4	.0731	0.435	0.616	0.871
	3	74.8	4713.6	.0804	0.363	0.513	0.725
	3.5	80.8	5092.0	.0864	0.311	0.440	0.622
	4	86.4	5444.3	.0924	0.272	0.385	0.544
^{127}I (Z=53)	.5	30.5	3876.6	.0328	1.968	2.783	3.936
	1	43.2	5483.1	.0463	0.986	1.394	1.971
	1.5	52.9	6716.3	.0567	0.657	0.929	1.314
	2	61.1	7752.4	.0654	0.493	0.698	0.986
^{197}Au (Z=79)	.5	30.5	6013.3	.0328	1.891	2.674	3.782
	1	43.2	8505.3	.0463	0.947	1.339	1.894
	1.5	52.9	10418.2	.0567	0.631	0.893	1.263

$\theta_{\text{plane}}^{\text{rms}}$ Mult Scatt6Booster ExitCopper Strippes (10, 40, 70, 100 mg/cm^2)

Ion	E/A	P/A ($\frac{\text{MeV}}{c \cdot A}$)	p ($\frac{\text{GeV}}{c}$)	β	10 mg/cm^2	40 mg/cm^2	70 mg/cm^2	100 $\frac{\text{mg}}{\text{cm}^2}$
^{32}S Z=16	300	805.5	25.78	0.6541	0.397	0.794	1.051	1.255
	350	880.1	28.2	0.6868	0.346	0.692	0.915	1.094
	400	951.4	30.44	0.7145	0.308	0.616	0.815	0.974
	450	1020.2	32.65	0.7385	0.278	0.554	0.736	0.879
	500	1086.97	34.78	0.7593	0.254	0.508	0.672	0.803
^{63}Cu Z=29	300	805.5	50.75	0.6541	0.366	0.732	0.968	1.157
	350	880.1	55.4	0.6868	0.318	0.636	0.841	1.006
	400	951.4	59.94	0.7145	0.283	0.566	0.749	0.895
	450	1020.2	64.27	0.7385	0.256	0.502	0.677	0.810
	500	1087.0	68.48	0.7593	0.233	0.466	0.616	0.737
^{127}I Z=53	300	805.5	102.3	0.6541	0.331	0.662	0.876	1.047
	350	880.1	111.8	0.6868	0.289	0.578	0.765	0.914
	400	951.4	120.8	0.7145	0.257	0.514	0.680	0.813
	450	1020.2	129.6	0.7385	0.232	0.464	0.614	0.734
	500	1087.0	138.0	0.7593	0.212	0.424	0.561	0.670
^{197}Au Z=79	300	805.5	158.7	0.6541	0.319	0.638	0.844	1.009
	350	880.1	173.4	0.6868	0.277	0.554	0.733	0.876
	400	951.4	187.4	0.7145	0.247	0.494	0.654	0.781
	450	1020.2	200.9	0.7385	0.223	0.446	0.590	0.705
	500	1087.0	214.1	0.7593	0.203	0.406	0.537	0.642

^{12}C Tandem Exit

$\theta_{\text{plane}}^{\text{rms}}$ Mult Scatt

GRAPHIC CORPORATION Buffalo, New York
Friedman J. S. A.
SEMI-LOGARITHMIC
1 CYCLE X 70 DIVISIONS
AD-0830-00

$\theta_{\text{plane}}^{\text{rms}}$ (mrad)

0.4⁴
0.3³
0.2²
0.1¹


6 6.5 7 7.5 8 8.5

MeV/A

20 $\mu\text{g}/\text{cm}^2 \text{ C}$

10 $\mu\text{g}/\text{cm}^2 \text{ C}$

5 $\mu\text{g}/\text{cm}^2 \text{ C}$



32S

Tandem Exit

θ_{plane}^{rms}

Mult Scatt

GRAPHIC CORPORATION
Farmingdale, U.S.A.
Barbosa West York
SEMI-LOGARITHMIC
1 CYCLE X 70 DIVISIONS
AD-0030-00

θ_{plane}^{rms}
(mrad)


0.1
0.2
0.3
0.4
0.5
0.6
0.7
10

3 3.5 4 4.5 5 5.5
MeV/A

20 $\mu\text{g}/\text{cm}^2$ C

10 $\mu\text{g}/\text{cm}^2$ C

5 $\mu\text{g}/\text{cm}^2$ C




^{63}Cu

Tandem Exit

$\theta_{\text{plane}}^{\text{rms}}$

Mult Scatt



AD-0030-00
1 CYCLE X 70 DIVISIONS
SEMI-LOGARITHMIC
GRAPHIC CONTROLS CORPORATION
Buffalo, New York
Printed in U.S.A.

$\theta_{\text{plane}}^{\text{rms}}$
(mrad)

20 $\mu\text{g}/\text{cm}^2$ C

10 $\mu\text{g}/\text{cm}^2$ C

5 $\mu\text{g}/\text{cm}^2$ C

(MeV/A)

12/7

Tandem Exit

θ_{plane}^{rms}

Multiscatt


10

5

1

0.5

0.1



SEMI-LOGARITHMIC 2 CYCLES X 70 DIVISIONS AO-0039-60

GRAPHIC PAPERS GRAPHIC CONTROLS CORPORATION Buffalo, New York, U.S.A.

197 Au

Tandem Exit

θ_{plane}^{rms}

Mult Scatt


10

5

1

0.5

0.1



SEMI-LOGARITHMIC
2 CYCLES X 70 DIVISIONS
AD-8000-80

GRAPH PAPER GRAPHIC CONTROLS CORPORATION
Butte, Nev. U.S.A.
Printed in U.S.A.


θ_{plane}^{rms}
(mrad)

MeV/A

32S Boosters Exit θ_{plane}^{rms} Mult Scatt

SEMI-LOGARITHMIC 2 CYCLES X 70 DIVISIONS AB-0033-60

GRAPHIC PAPERS GRAPHIC CONTROLS CORPORATION B. 1100 N. 17th Ave. PRINCETON, N. J. U. S. A.




1

0.5

0.1

250 300 350 400 450 500 (MeV/A)

^{63}Cu Booster Exit $\theta_{\text{plane}}^{\text{rms}}$ Mult Scatt



GRAPHIC COPY FILM CORPORATION Buffalo, New York
PR 4025-111 I A

0.1 1 2 3 4 5 6 7 8 9 10

127 I

Booster Exit

θ_{rms}
plane

Mult Scatt

10

5

AP-0830-60

2 CYCLES X 70 DIVISIONS

SEMI-LOGARITHMIC

θ_{rms}
plane
(mrad)

1

GRAPHIC CONTROLS CORPORATION

Bohler, H. # 7001
F. 1000 U.S.A.

0.5

GRAPH PAPER

0.1

250

300

350

400

450

500

(ANAL)

100 mg/cm² Cu

70 mg/cm² Cu

40 mg/cm² Cu

10 mg/cm² Cu



197Au

Booster Exit

θ_{plane}^{rms}

Mult Scatt

10

5


SEMI-LOGARITHMIC
2 CYCLES X 70 DIVISIONS
00-9033-80

θ_{plane}^{rms}
(mrad)

1

GRAPHIC CONTROLS CORPORATION
Buffalo, N.Y. 14240
Printed in U.S.A.

0.5



100 mg/cm² Cu

70 mg/cm² Cu

40 mg/cm² Cu

10 mg/cm² Cu

250 300 350 400 450 500
(MeV/A)

0.1

Tandem $241T$, Carbon, β_{max}^* values for $\Delta E = 0.577 \text{ mm mrad}$

Ion	MeV/A	$5 \mu\text{g/cm}^2$	$10 \mu\text{g/cm}^2$	$20 \mu\text{g/cm}^2$
^{12}C	6.5	15.1	7.51	3.77
	7	17.5	8.75	4.35
	7.5	20.0	10.0	5.01
	8	22.8	11.4	5.71
	8.5	25.9	12.5	6.42
^{32}S	3.5	4.38	2.19	1.09
	4	5.71	2.85	1.42
	4.5	7.23	3.61	1.81
	5	8.90	4.45	2.23
	5.5	10.8	5.41	2.71
^{63}Cu	2	1.69	0.85	0.42
	2.5	2.64	1.32	0.66
	3	3.79	1.89	0.95
	3.5	5.17	2.59	1.29
	4	6.76	3.38	1.69
^{127}I	.5	0.13	0.065	0.032
	1	0.51	0.257	0.128
	1.5	1.16	0.579	0.29
	2	2.06	1.03	0.51
^{197}Au	.5	0.14	0.070	0.035
	1	0.56	0.28	0.14
	1.5	1.26	0.63	0.31

Booster Exit, Copper, β^+ max values for $\Delta E = 1 \pi$ mm mrad

Ion	MeV/A	10 mg/cm ²	40 mg/cm ²	70 mg/cm ²	100 mg/cm ²
³² S	300	6.34	1.59	0.91	0.63
	350	8.35	2.09	1.19	0.84
	400	10.5	2.63	1.5	1.05
	450	12.9	3.23	1.84	1.29
	500	15.5	3.88	2.21	1.55

⁶³ Cu	300	7.47	1.87	1.07	0.75
	350	9.89	2.47	1.41	0.99
	400	12.5	3.12	1.79	1.25
	450	15.3	3.83	2.19	1.53
	500	18.4	4.6	2.63	1.84

¹²⁷ I	300	9.13	2.28	1.30	0.91
	350	11.97	2.99	1.71	1.20
	400	15.1	3.78	2.16	1.51
	450	18.6	4.65	2.66	1.86
	500	22.2	5.55	3.17	2.22

¹⁹⁷ Au	300	9.83	2.46	1.40	0.98
	350	13.0	3.25	1.86	1.30
	400	16.4	4.1	2.34	1.64
	450	20.1	5.03	2.87	2.01
	500	24.3	6.08	3.47	2.43


Tandem Exit

^{12}C

β_{mix}^{40} values (m)

MuHScott

GRAPH PAPER GRAPHIC CONTROLS CORPORATION Buffalo, New York, U.S.A.
SEMI-LOGARITHMIC 2 CYCLES X 70 DIVISIONS AD-00000-00



Tandem Exit

^{32}S

β^+ max values (m)

Mult Scatt

AG-0085-30
2 CYCLES X 70 DIVISIONS
SEMI-LOGGING
GRAPHIC CONTROLS CORPORATION
Burlingame, New York
Printed in U.S.A.

20^2

$\beta^*(m)$

10

5

2

1

3

3.5

4

4.5


5.5

(MeV/A)⁵

5 $\mu\text{g}/\text{cm}^2$ C

10 $\mu\text{g}/\text{cm}^2$ C

20 $\mu\text{g}/\text{cm}^2$ C



Tandem Exit

^{63}Cu

β^* max values (m)

Mult Scatt

10

5


2

1

0.5

0.2²

0.1




GRAPHIC CONTROLS CORPORATION Buffalo, New York
 Printed in U.S.A.
 SEMI-LOGARITHMIC
 2 CYCLES X 70 DIVISIONS AD-0033-80

Tandem Epit 127I β^+ max. values (m) Mult Scatt


SEMI-LOGARITHMIC 2 CYCLES X 70 DIVISIONS AB-0683-80
 GRAPHIC CONTROLS CORPORATION Printed in U.S.A. Cambridge, Mass. Vert.

β^+ (m)
 0.1
 0.05
 0.02
 0.01

5 $\mu\text{g}/\text{cm}^2 \text{ C}$
 10 $\mu\text{g}/\text{cm}^2 \text{ C}$
 20 $\mu\text{g}/\text{cm}^2 \text{ C}$



1 Tandem Exit ^{197}Au β^* max values Mult Scatt



GRAPH PAPER GRAPHIC CONTROLS CORPORATION Buffalo, New York Printed in U.S.A.

SEMI-LOGARITHMIC

2 CYCLES X 70 DIVISIONS

AB-0033-80

0.5

0.2

0.1


0.05

0.02

0.01


10 Booster Exit ^{32}S β^+ max values Mult Scatt

AD-6893-60
2 CYCLES X 70 DIVISIONS
 β^+ SEMI-LOGARITHMIC
GRAPHIC CONTROLS CORPORATION
Burlington, N.J. 07078
Printed in U.S.A.



Booster Epi+ ⁶³Cu β^{max} values Mult Seatt

GRAPH PAPER GRAPHIC CONTROLS CORPORATION Buffalo, New York
 2 CYCLES X 70 DIVISIONS 70-0033-68
 SEMI-LOGARITHMIC



50

20

10

5

2

1

10 mg/cm² Cu

40 mg/cm² Cu

70 mg/cm² Cu

100 mg/cm² Cu

250 300 350 400 450 500 (I₀V/A)

Booster Exit 127 I β^* max values Mult Scatt

SEMI-LOGARITHMIC 2 CYCLES X 70 DIVISIONS AD-8033-60 GRAPHIC CONTROLS CORPORATION Buffalo, New York Printed in U.S.A.

β^* (m)

50


20

10

5

2

1



100 Booster Exit ^{197}Au β^* max values Mult Scatt

100
50
20
10
5
2
1

β^* (m)

GRAPHIC CONTROLS CORPORATION
PHOTO, NEW YORK
U.S.A.
SEMI-LOGARITHMIC
2 CYCLES X 70 DIVISIONS
AO-6533-80

