



Brookhaven
National Laboratory

BNL-101595-2014-TECH

RHIC/PG/52;BNL-101595-2013-IR

Electron Trappings In RHIC From A Debunched Proton Beam

A. G. Ruggiero

April 1984

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Electron Trappings
in RHIC
from a Debunched Proton Beam

A. G. Ruggiero

BNL, April 27, 1984

(1)

Ion Trapping / Electron Trapping

Proton Beam Completely Debundled

Gas Composition as derived in RHIC-PG-51

Densities:

Molecules	<u>Z</u>	<u>A</u>	<u>warm</u>	<u>cold</u>
			25%	75%
H ₂	2	2	$2.1 \times 10^7 \text{ cm}^{-3}$	$2.3 \times 10^7 \text{ cm}^{-3}$
He	2	4	-	$2.3 \times 10^7 \text{ cm}^{-3}$
CO	14	28	$2.1 \times 10^7 \text{ cm}^{-3}$	-

Ionization cross-section σ_i

$$\sigma_i = 1.8 \times 10^{-19} Z_i^2 \text{ cm}^2$$

Ionization Rate for the i th specie per proton

$$\frac{dn_i}{dt} = c \sigma_i n_i \quad (\beta = 1)$$

(2)

<u>ions</u>	ionization rate
H_2^+	0.243 s^{-1} / proton
He^+	0.186
CO^+	0.397
electrons	0.826 s^{-1} / proton

The ions (positively charged) are rejected by the potential barrier of the proton beam -

The electrons could be trapped

Average energy of the electrons at production

$$\frac{1}{40} \text{ eV}$$

which corresponds to a velocity

$$v_e \approx 10^7 \text{ cm/sec}$$

(3)

To "clear" the electrons one can create one or several gaps in the beam - these gaps though are to be no less than ~~300 nsec~~ nsec if the electrons have to reach a wall 3 cm away with the velocity given above -

Assume the probe beam has a uniform density distribution with elliptical cross-section of semi-axis σ_x, σ_z -

$$I, \text{ beam current} = Ne f_{\text{rev}} = 57 \times 10^{12} \times 1.6 \times 10^{-19} \times 78.2 \times 10^3 \\ = 0.71 \text{ Amperes}$$

Electric Field produced by the beam

$$E_{x,z} = \frac{(120 \pi \text{hm}) I_{\text{avg}}}{\sigma_x + \sigma_z} \frac{(x,z)}{\sigma_{x,z}} \quad \text{Volt/cm}$$

The voltage distribution is

$$V = \frac{43 \text{ Volt}}{\sigma_x + \sigma_z} \left(\frac{x^2}{\sigma_x} + \frac{z^2}{\sigma_z} \right)$$

(4)

The potential barrier depth is calculated by setting $x = \sigma_x$ and $z = \sigma_z$, and that is

$$\Delta V = 43 \text{ Volts}$$

Oscillation Frequency of the Trapped Electrons

$$\omega_{x,z}^2 = \frac{(120 \text{ ohm}) I e c^2}{m_e c^2 \sigma_{x,z} (\sigma_x + \sigma_z)}$$

where $m_e c^2 = 0.5 \text{ MeV}$ and $\sigma_{x,z}$ is in meter.

$$\langle \sigma_{x,z} (\sigma_x + \sigma_z) \rangle \approx 20^2, \quad \sigma \approx 4 \text{ mm}$$

As a result

$$\omega_{x,z} \approx 700 \text{ MHz}$$

Betatron Tune Depression

$$\Delta \nu_{x,z} = \frac{e}{4\pi m \beta^2 \gamma c^2} \oint ds \beta_{x,z}(s) \frac{\partial E_{x,z}^{electron}(s)}{\partial x,z}$$

$E^{electron}$ is the electric field generated by the electrons in the background

$$\frac{\partial E_{x,z}^{electron}}{\partial x,z} = \eta \frac{(120 \text{ statV}) I}{\sigma_{x,z} (\sigma_x + \sigma_z)} = \frac{86 \eta}{\sigma_{x,z} (\sigma_x + \sigma_z)}$$

η , charge neutralization coefficient caused by the electrons

We have

$$\Delta \nu_{x,z} = \frac{120 e I}{4\pi m \beta^2 \gamma c^2} \oint \frac{\eta(s) \beta_{x,z}(s)}{\sigma_{x,z} (\sigma_x + \sigma_z)} ds$$

$$= \frac{120 I e}{8\pi m c^2 \beta^2 \gamma} \bar{\eta} \int \frac{\beta(s)}{\sigma^2(s)} ds$$

or

(6)

$$\Delta v = \frac{120 I_e}{4\pi mc^2 \beta^2 \gamma} \frac{\bar{\eta}}{\varepsilon} 2\pi R$$

with

$$\varepsilon = \frac{20^2}{\beta} = 0.05 \text{ @ mm mrad}$$

$$R = 610.2 \text{ m}$$

$$\gamma = 100$$

$$mc^2 = 0.9383 \text{ GeV}$$

We get

$$\Delta v = 4.64 \bar{\eta}$$

It is safe to keep
therefore we requires

$$\Delta v \leq 0.0025$$

$$\bar{\eta} \leq 0.0005$$

7

Removal of the Electrons

Electrons have a low individual drift
After a time t they have travelled a
distance

$$l = v_e t$$

where $v_e \approx 1 \times 10^7$ cm/sec

We assume that after this distance they are
removed. We take

$$t = \eta \tau_e$$

$\tau_e = 1.21$ sec is the production time per
proton

With $\eta = 0.0005$

$$l = \underline{\underline{6,050 \text{ m}}}$$

(8)

Drift in a Dipole (B_z)

This occurs at velocity v_D

$$v_D = \frac{E_x}{B_z} \frac{\omega_L^2}{\omega_L^2 + \omega_x^2} = \alpha \frac{\omega_L \omega_x^2}{\omega_L^2 + \omega_x^2}$$

$$\omega_L = \frac{e B_z}{m_e} \quad , \quad 2\pi \times \text{Larmor frequency}$$

$$= 5.6 \times 10^{11} \text{ Hz}$$

Therefore $\omega_L \gg \omega_x$, and

$$v_D = \alpha \frac{\omega_x^2}{\omega_L} = 3.5 \times 10^5 \text{ cm/sec}$$

Drift in a Quadrupole with gradient B'

$$v_Q = x \frac{\omega_Q \omega_x^2}{\omega_Q^2 + \omega_x^2}$$

where

$$\omega_Q^2 = \frac{1}{2} \left(\frac{eB'}{mc} \right)^2 \sigma^2$$

with $B' = 550 \text{ kG/m}$ and $\sigma = 4 \text{ mm}$

$$\omega_Q = 2.9 \times 10^{10} \text{ Hz}$$

Therefore $\omega_Q \gg \omega_x$ and

$$v_Q = \sigma \frac{\omega_x^2}{\omega_Q} = 6.8 \times 10^6 \text{ cm/sec}$$

In ~~the~~ conclusion the drift velocity is quite smaller in the dipoles - The calculations of page 7 to clear the electrons have to be repeated by taking

$$v_e = v_D \approx 3.5 \times 10^5 \text{ cm/sec}$$

We obtain then

$$l = 212 \text{ m}$$

This corresponds to the length traversed in the dipoles: it is about 10 dipoles, that is 5 cells.

Solution: install pairs of clearing electrodes in the warm sections at the beginning and at the end of each long straight section -


Install also pairs of clearing electrodes in the arcs, equally spaced, 4 full cells apart - So there are in total 24 pairs, 12 in cold sections and 12 in warm sections (see figure next page) -

Take a gap separation d flush with the vacuum chamber and ~~to~~ a length $L = d$ -

To sweep the electrons away the following voltage is required

$$V_c = m_e c^2 \left(\frac{dV_e}{cL} \right)^2$$

which is quite negligible compared to $\Delta V = 43 \text{ Volt}$
It is quite safe to take $V_c \approx 100 \text{ Volt}$



- Cleaning Electrods in cold section
- x " " in warm section