

## A Chromatic Correction Scheme for the RHIC3 Lattice

A. Antillon

January 1985

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

High Energy Facilities  
Advanced Projects

RHIC-8

BROOKHAVEN NATIONAL LABORATORY  
Associated Universities, Inc.  
Upton, New York 11973

RHIC Technical Note No. 8

A Chromatic Correction Scheme  
for the  
RHIC3 Lattice

Armando Antillón

January 2, 1985

ABSTRACT

The RHIC3 lattice is the current design for the Relativistic Heavy Ion Collider. In this note we use special families of sextupoles in the arcs to reduce, in insertions, momentum dependence of machine functions.


## 1. Introduction

A sextupole scheme has been proposed<sup>1,2</sup> for the chromaticity correction of the symmetric RHIC3 lattice. In such schemes, 3 families of sextupoles have been placed very close to the arcs. An attempt to improve the chromatic behavior has been made manually by decoupling the 3 families in 6. There was improvement, but they were not very significant.

In order to avoid the small dispersion and the lack of symmetry in the insertions, in x and y directions, we analyse the placement of the sextupoles in the arcs. The method we follow is a mix of mathematics and intuition. The linear part of  $\Delta\beta/\beta$  ( $\phi, \Delta p/p$ ) help us to understand that we must have symmetry around the center of the arcs,<sup>3</sup> and intuition the rest.

## 2. Description of the RHIC3 Lattice

The RHIC3 lattice has 6 arcs and 6 insertions with supersymmetry 6. Each typical cell has the structure



where X<sub>1</sub> and X<sub>2</sub> are available sextupoles. Each arc has 12 cells. One of the insertions is as follows:

I = Q8I Q7I Q6I BS Q5I BS Q4I Q3I Q2I Q1I Q1\* BC2I QC BC1I  
CR BC10 QC BC20 Q1\* Q10 Q20 Q30 Q40 BS Q50 BS Q60 Q70 Q80

Each superperiod is made as follows:

$$6\{C_{\text{Inner}}\} + I + 12\{C_{\text{out}}\} + I^{-1} + 6\{C_{\text{Inner}}\} .$$

In the arcs,

	$\beta_x$	$\beta_y$	$\eta_x$
QF	51.6	7.6	1.3
QD	7.6	51.6	.66

The natural chromaticity is

$$\xi_x = -73.9, \quad \xi_y = -64.8$$

and the nominal tunes are

$$\nu_x = 34.41847, \quad \nu_y = 34.41628 \quad .$$

For a chromaticity of

$$\xi_x = 1 = \xi_y$$

the two families of sextupoles have values ( $B''\ell_{\text{sex}}/B\rho \text{ m}^{-2}$ )

$$\begin{aligned} \text{SF} &= -.19460 \\ \text{SD} &= +.41529 \quad . \end{aligned}$$

### 3. Method for Choosing Special Families of Sextupoles

The linear expression<sup>4</sup> of  $\Delta\beta/\beta$  ( $\phi, \Delta\rho/\rho$ ) has been used for obtaining the contribution of each sextupole to the  $\beta$ 's max in the insertions. We can see that the contribution is symmetric around the center of each arc. However, as our lattice is symmetric, we must expect the same distribution of the families in all the arcs. This idea is enhanced by Figure I, as we now explain. Figure I.a. shows the behavior of  $\beta_x, \beta_y$  in one superperiod, for a deviation in the momentum of 0%. Figures I.b,c have the same meaning for  $\Delta p/p = +1.0\%$  and  $-1.0\%$  respectively. In both cases  $\beta_y$  has been shifted up in one arc in order to have clarity in the drawing. With the help of Patricia, we have observed some general properties for RHIC.

1. The sextupoles in the center of the arcs (Z) are necessary to maintain the symmetry of the machine functions.

2. The sextupoles placed where  $\beta_x$  has locally a maximum (S8, B, D, F, H, J, Z) change practically only the x direction ( $\beta_x, v_x, \dots$ ), and the same is applicable to  $\beta_y$  and the y direction for A, C, E, G, I, K.
3. The sextupoles connected by the (---) line have the same sign and opposite to the sign of those connected by the (-.-.) line which also have the same sign.






Fig. I.b.  $\beta_x$  and  $\beta_y$  for momentum deviation of +1%.  $\beta_y$  has been shifted up in the central arc to avoid overlapping with  $\beta_x$ . The sextupoles S8, B, D, F, H, J, Z correct practically only x direction, and A, C, E, G, I, K correct practically only y direction.






Fig. 1.c.  $\beta_x$  and  $\beta_y$  for momentum deviation of -1%.

Only a few runs of Patricia have shown that good values for this sextupoles are

A = -0.21803	C = 0.22142
E = -0.18914	G = 0.12302
I = -0.	K = 0.
S8 = 0.	
D = 0.	B = -0.
H = 0.0535	F = -0.01837
Z = 0.21057	J = -0.02316

In order to keep  $\xi_x = \xi_y = 1$ , SF and SD now have the values

SD = 0.42513  
SF = -0.21240 .

In Figure II, we compare the machine functions for the case of two families, SF and SD (continuous line) and the latter case of 8 extra families of sextupoles (dashed line). The chromatic behavior has been improved even without an optimization of the sextupoles.




Fig. IIa

Fig. II. In this figure the continuous lines give the momentum dependence of machine functions for two standard sextupole families, SF and SD. The dashed lines represent the same with special families of sextupoles (not optimized) a) change in the tune, b) fractional change of the  $\beta$ 's in the insertions (QC, Q1), c) fractional change of the  $\beta$ 's at the crossing points, and d) dispersion function at the crossing point.




Fig. IIb





Fig. IIc




Fig. IIId

## Summary

We have introduced a simple sextupole scheme for the correction of chromatic effects in RHIC which give good improvement even for nonoptimized special sextupoles. However, the nature and simplicity of the method make us to have the conjecture that it can be extended to other symmetric machines with symmetric sextupole distribution, and to antisymmetric machines with antisymmetric sextupole distribution.

## References

1. J. Claus, private communication.
2. RHIC Proposal, BNL 51801.
3. A. Antillon, RHIC-AP-1, BNL.
4. E.D. Courant and S. Snyder, Ann. Phys. 3, 1, (1958).