

BNL-101925-2014-TECH AD/RHIC/13;BNL-101925-2013-IR

Beam Life Time in the Presence of Beam Blow Up

S. Y. Lee

November 1985

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

High Energy Facilities Advanced Projects

BROOKHAVEN NATIONAL LABORATORY Associated Universities, Inc. Upton, New York 11973

RHIC Technical Note No. 13

Beam Life Time in the Presence of Beam Blow Up

S. Y. Lee

11/12/85

Ŧ.

1. Introduction

The life time of the beam particles in the accelerators is determined by (1) the beam dynamics in the presence of nonlinear magnetic elements, (2) the interaction between the beam particles and the gas molecules in the vacuum tube, (3) the interaction between the particles in the two colliding beams and (4) the beam-blow up rate in the presence of collective effect, intrabeam scattering^{1,2,3)} etc.

This note is to evaluate the beam life time of the relativistic heavy ion collider, where the intrabeam Coulomb scattering is an important factor in the design study.

2. Beam life time due to reaction between two colliding beams

When the beam life time is determined mainly by beam-beam reaction, the reactin rate can be expressed as

$$\lambda = 6 \, \mathscr{L}\sigma_{\rm bb}^{\rm /BN} \tag{1}$$

where \mathscr{L}, σ_{bb} , B and N_B are the luminosity, the reaction cross section, the number of bunches in the accelerator and the number of particles per bunch. The factor 6 in Eq. (1) comes from 6 interaction points in RHIC. The total cross section σ_{bb} consists of nuclear reaction, Coulomb reaction and Bremsstrahlung with pair production and subsequent e capture (4) causing the beam to be lost. Table 1 shows the beam life time of RHIC due to various reaction mechanism.

3. Effect of beam blow up on the life time

The reaction rate λ defined in Eq. (1) depends on the luminosity and the number of particles in the beam. After some simple algebra, we obtain

$$\lambda(t) = \frac{9\gamma N_B \sigma_{bb} f_{rev}}{\beta \epsilon_N}$$
(2)

J N

· 1

	Beam-gas nuclear reaction λ 1	Beam-beam nuclear reaction $\frac{\lambda}{2}$		Beam-beam Coulomb dissociation λ_3	(1,2,2,2,2,3,3,3,3,3,3,3,3,3,3,3,3,3,3,3,	Initial Half Life
Beam	@ 10 ⁻¹⁰ Torr	A on A	p on A	A on A	A on A	A on A
P	×10 ⁻³ /h 0.15	×10 ⁻³ /h 1.6	×10 ⁻³ /h 1.6	×10 ⁻³ /h	×10 ⁻³ /h	h 396
đ	0.19	8.7	3.0	 ·		78
С	0.36	5.2	16.9	 /		125
S	0.55	4.6	27.9			135
Cu	0.76	4.6	38.8	0.52	0.12	116
I	1.08	3.9	55.1	13.2	4.6	30
Au	1.37	2.1	69.3	16.0	31.6	14

Table l.	Initial	Reaction	Rate λ =	-I dI/	dt and Total	Half
	Life of	Particle	Beams for	Head-on	Collisions	

where γ , N_B , ε_N , f_{rev} , σ_{bb} and β^* are the Lorentz relativistic factor, number of particles per bunch, the normalized emittance, the revolution frequency, total cross section between particles in two beams and the betatron amplitude function at the interaction point respectively. Since N_B and ε_N are both function of time, we obtain

$$\frac{1}{\lambda} \frac{d\lambda}{dt} = \frac{1}{N_{\rm B}} \frac{dN_{\rm B}}{dt} - \frac{1}{\varepsilon_{\rm N}} \frac{d\varepsilon_{\rm N}}{dt}$$

$$= -\lambda - \mu(t) \qquad (3)$$

0 2

The equation for the reaction rate is indeed nonlinear. When the emittance blow up rate $\mu(t)$ is taken to be a CONSTANT, Eq. (3) can easily be solved to give,

$$\lambda = \lambda_{o} e^{-\mu t} / \left(1 + \frac{\lambda_{o}}{\mu} \left(1 - e^{-\mu t}\right)\right)$$
(4)

where λ_0 is the initial reaction rate. When the beam blow-up rate is faster then the reaction rate, i.e., $\mu >> \lambda_0$, we obtain

$$\lambda \simeq \lambda_{o} e^{-\mu t}$$
 (5)

On the other hand, when the beam does not blow up, i.e., $\mu = 0$, we have $\lambda \approx \lambda_0$. In the heavy ion collider, we encounter $\mu >> \lambda_0$ (3).

From Eq. (4), we obtain then

$$I(t)/I_{o} = (1 + \frac{\lambda_{o}}{\mu} (1 - e^{-\mu t}))^{-1}$$

$$\approx \exp\left(\frac{\lambda_{o}}{\mu} (1 - e^{-\mu t})\right).$$
(6)

Figure 1 shows $\lambda(t)$ and N(t) as a function of time for Au ion (eqs. (4) and (6)) with parameters $\mu = 0.1/hr$, $\lambda_0 = .0495/hr$. The number of particles N(t) is to be compared with the constant λ_0 decay law of $e^{-\lambda_0 t}$.

4. Luminosity

The luminosity depends on the beam size as well as the intensity of the beam,

$$\mathscr{L}(t) = 6 \frac{\gamma N_B^2 B f_{rev}}{4\varepsilon_N \beta^*}$$
 (7)

Thus,

$$\frac{1}{\mathscr{L}}\frac{\mathrm{d}\mathscr{L}(t)}{\mathrm{d}t} = -2\lambda - \mu \tag{8}$$

or,

$$\mathscr{L}(t)/\mathscr{L}_{o} = e^{-\mu t}/(1 + \frac{\lambda_{o}}{\mu} (1 - e^{-\mu t}))^{2}$$
 (9)

Figure 1 shows the luminosity as a function of time for Au ion. The average luminosity for a T hours run becomes

$$\langle \mathscr{L} \rangle / \mathscr{L}_{o} = \frac{1}{\mu T} (1 - e^{-\mu T}) / (1 + \frac{\lambda_{o}}{\mu} (1 - e^{-\mu T})) .$$
 (10)

The average luminosity for T = 10 hours operation of Au on Au becomes

$$<\mathcal{L}>/\mathcal{L}_{o}$$
 = 0.48 (μ = 0.1/hr, λ_{o} = 0.0495/hr).

For the most other ions, the Coulomb dissociation and pair production cross sections are rather small⁴⁾. The reaction rate is irrelevant in determining the average luminosity. The average luminosity is given by (for $\mu = 0.1/hr$)

$$\langle \mathscr{L} \rangle / \mathscr{Q}_{o} = (1 - e^{-\mu T}) / \mu T \simeq .63$$



Fig. 1. The beam depletion rate, $\lambda(t)$, the number of particles, $N(t)/N_0$, and the luminosity $\mathscr{L}/\mathscr{L}_0$ are plotted as a function of time t. $\mu = 0.1/hr$, $\lambda_0 = .0495/hr$ for Au ion is used in this calculation.

APPENDIX

In general, the beam emittance blow up rate $\mu(t)$ is not a constant. Intrabeam scattering calculation shows that⁵⁾

$$\varepsilon_{\rm N}(t) \simeq \varepsilon_{\rm N}^{\rm o} + \infty/t$$
 (A.1)

We obtain therefore

$$\mu(t) = \frac{1}{\varepsilon_{N}} \frac{d\varepsilon_{N}}{dt} = \frac{\frac{1}{2}\alpha t^{-\frac{1}{2}}}{\varepsilon_{N}^{0} + \alpha \sqrt{t}} . \qquad (A.2)$$

The solution of Eq. (3) becomes

$$\lambda(t) = e^{-\int_{0}^{t} \mu(t)dt} \left(\frac{1}{\lambda_{o}} + \int_{0}^{t} e^{-\int_{0}^{t'} \mu dt''} dt'\right)^{-1}$$

$$= \lambda_{o} / (1 + x) \left\{1 + 2\lambda_{o} \left(\frac{\varepsilon_{N}^{o}}{\alpha}\right)^{2} \left(x - \ln(1 + x)\right)\right\}$$
(A.3)

where

$$x(t) = \frac{\alpha \sqrt{t}}{\varepsilon_{N}^{o}} . \qquad (A.4)$$

Whence the intensity and the luminosity become

$$-\int^{t} \lambda(t') dt'$$

I = I_o e^o (A.5)

$$\mathscr{L} = \mathscr{L}_{o} e^{o} \left| \frac{-2\int^{t} \lambda(t') dt'}{(1 + x(t))} \right|.$$
 (A.6)

$$\partial = \mathcal{X}$$

References

- 1. A. Piwinski, Proc. 9th Int. Conf. on High Energy Accelerators, p 405 (1974).
- 2. J.D. Bjorken and S.K. Mtingwa, Particle Accelerator 13, 115 (1983).
- G. Parzen, Strong Intrabeam Scattering in Heavy Ion and Proton Beams; IEEE Trans. Nucl. Sci. <u>NS-32</u>, No. 5, Part II, 3466 (1985).
- 4. S.Y. Lee and J. Weneser, "The mutual dressing of bare relativistic heavy ions during transit in collider orbit," to be published.
- 5. S.Y. Lee, Beam Loss Due to Aperture Limitation Resulting from Intrabeam Scattering, BNL-35438.