

RHIC Dynamic Aperture for Lattices With Some $\beta^* = 3$ Insertions

G. Parzen

May 1987

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

RHIC Dynamic Aperture for Lattices With Some $\beta^* = 3$ Insertions

G. Parzen

May 12, 1987

1. Introduction

This note points out that it is possible to operate RHIC at $\gamma = 30$ with some insertions having $\beta^* = 3$, and still have enough dynamic aperture to allow the beam to grow because of intrabeam scattering. At the $\beta^* = 3$ insertions the luminosity is higher by a factor of 2.

Tracking studies show that starting with a lattice that has six $\beta^* = 6$ insertions, the dynamic aperture decreases as one introduces $\beta^* = 3$ insertions, and the decrease is roughly monotonic with the number of $\beta^* = 3$ insertions. The results show that RHIC can operate with a beam of gold ions at $\gamma = 30$ for ten hours with up to three $\beta^* = 3$ insertions.

2. Tracking Results

Figure 1 plots the stability limit, A_{SL} or dynamic aperture found as the number of $\beta^* = 3$ insertions is increased from 0 to 6. Note that $\beta^* = 6$ at the insertions where $\beta^* \neq 3$. The tracking runs start with $\epsilon_x = \epsilon_y$, and the A_{SL} plotted is the lowest A_{SL} found for 10 different sets of random field errors at $\Delta p/p = 0$.

Figure 2 adds the dynamic aperture, A_{SL} , required by the intrabeam scattering for a beam of gold ions with $N_b = 1.1 \times 10^9$ ions/bunch at $\gamma = 30$, for ions that start out with $\epsilon_x = \epsilon_y$.

One sees from Figure 2, that one can have up to three $\beta^* = 3$ insertions for a beam of gold ions at $\gamma = 30$.

Figure 1 also seems to indicate that the periodicity of the $\beta^* = 3$ insertions does not seem to play a large role. The lattice with three $\beta^* = 3$ insertions has a periodicity = 3, the lattice with six $\beta^* = 3$ insertions has a periodicity = 6, and the lattice with one $\beta^* = 3$ and five $\beta^* = 6$ insertions have a periodicity = 1. The result in Figure 1 seems to indicate a roughly monotonic dependence of A_{SL} on the number of $\beta^* = 3$ insertions with little dependence on the periodicity of the insertions.

I am much indebted to S.Y. Lee for providing the input data for the various lattices that were studied.




Fig. 1.




Fig. 2