

BNL-101938-2014-TECH AD/RHIC/26;BNL-101938-2013-IR

The Beam-Beam Interaction in Lattices with Non-Identical Insertions

G. Parzen

April 1987

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Accelerator Development Department

۴ Þ

Brookhaven National Laboratory Associated Universities, Inc. Upton, New York 11973

RHIC Technical Note No. 26

The Beam-Beam Interaction in Lattices with Non-Identical Insertions

G. Parzen

April 29, 1987

The Beam-Beam Interaction in Lattices with Non-Identical Insertions

ł

G. Parzen 4/29/87

l

1. INTRODUCTION

If one starts out with a lattice with 6 identical insertions, then the beam-beam interaction has the periodicity 6, and the stop bands, due to the non-linear beam-beam forces, are at the resonance lines $nv_x + mv_y = q$ where q is a multiple of 6. If now some of these insertions are replaced with insertions having different β_x , β_y at the interaction point, then the beam-beam interaction effects could be worsened because the periodicity of the beam-beam interaction may be lowered. This paper shows that if the different insertions introduced obey certain conditions listed below, then the beam-beam stop-bands and the beam-beam v-shift are not changed by the introduction of the different insertions. These conditions on the insertions are the following:

- 1) β_y / β_x is the same for all insertions.
- 2) $\Delta \psi_x \Delta \psi_y$ the change in the betatron phase between crossing points is not changed by the different insertions.
- 3) $\beta_{x'}\beta_{y}$ and the $\sigma_{x'}\sigma_{y}$ of the beam may be considered to be constant along the path where the beams interact.

4) $\alpha / \beta_x^{1/2}$ is the same for all insertions, where α is the crossing angle of the beams. Usually, the above conditions are easy to fulfill, as they appear to occur naturally. It would seem desirable to require that, if possible, all insertions introduced into the lattice should obey the above 4 conditions.

2. Derivation of the Conditions

The beam-beam stop bands are given by;

$$\Delta \upsilon_{nm} \sim \int ds \ \beta_x^{n/2} \ \beta_y^{n/2} \ D_x^n \ D_y^m \ V \ exp\left[i \ (n\psi_x \ + \ m\psi_y)\right], \qquad (2.1)$$

2

where we have dropped factors which are the same at all the insertions in the lattice $D_x = \partial/\partial x$ and $D_y = \partial/\partial y$. V = V(x,y) is the potential function that gives the field due to the beam.

The factor $\beta_x^{n/2} \beta_y^{n/2}$ in Eq. (2.1) indicates that insertions with larger β_x, β_y contribute more to the stop bands. However at larger β_x, β_y the beam becomes larger and $D_x^n D_y^m V$ becomes smaller by the same factor $1/(\beta_x^{n/1} \beta_y^{m/2})$ provided β_y/β_x and $\alpha/\beta_x^{1/2}$ are the same for all insertions.

This can be verified by starting with the expression for V(x,y)

$$V = -\frac{\lambda}{4\pi\varepsilon_{o}} \int_{0}^{\infty} dt \, \frac{1 - \exp\left[-\left(x^{2}/(2\sigma_{x}^{2} + t) + y^{2}/(2\sigma_{y}^{2} + t)\right)\right]}{(2\sigma_{x}^{2} + t)^{1/2} (2\sigma_{y}^{2} + t)^{1/2}}, \qquad (2.2)$$

and differentiating under the integral sign to find $D_x^n, D_y^m V$ and evaluating the result at $x = \alpha s$, y = 0. This involves some manipulation of the integrand.

The condition (2) on $\Delta \psi_x$ and $\Delta \psi_y$ between crossing points guarantees that the contributions due to the 6 interaction regions will add up with the same phases.

The v-shifts are given by Eq. (2.1) without the phase factor and by putting n = 2, m = 0 for Δv_x and n = 0, m = 2 for Δv_y . Thus the above arguments for the stop bands work for the v-shifts above.