



Brookhaven
National Laboratory

BNL-102112-2014-TECH

RHIC/AP/1;BNL-102112-2013-IR

Gold Ion Diffusion to Pair Production

S. Peggs

June 1993

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Gold ion diffusion due to pair production

Pair production is NOT a potent transverse or longitudinal diffusion mechanism. This can be seen as follows.

The cross section for pair production from gold ions at full energy ($T = 100 \text{ GeV/u}$) is [1]

$$\sigma_{\text{pair}} = 33,000 \text{ barns} = 3.3 \times 10^{-20} \text{ cm}^2 \quad 1$$

so that the rate of such collisions for 2 IRs at the design luminosity of $2 \times 10^{26} \text{ cm}^{-2}\text{sec}^{-1}$ is

$$\frac{dN}{dt} = 2 L \sigma_{\text{pair}} = 1.32 \times 10^7 \text{ pairs per second} \quad 2$$

Since there are nominally $N_B = 57$ bunches each containing $N_I = 10^9$ gold ions, the mean number of pair producing events that a single test ion undergoes during a $T_{\text{store}} = 36,000$ second store (assuming all parameters are constant) is

$$\langle n \rangle = \frac{\frac{dN}{dt} T_{\text{store}}}{N_B N_I} = 8.3 \quad 3$$

that is, a small number of order 1. It only remains to show that the typical longitudinal and transverse kicks on a gold ion during pair production are small compared to the natural bunch distributions.

The rms transverse momentum kick and rms instantaneous energy loss during pair production are expected to be of order 1 MeV/c and 1 MeV , respectively[1,2]. In contrast, the rms transverse beam momentum is typically

$$\langle x'^2 \rangle^{1/2} = \sqrt{\frac{\epsilon_N}{6\pi \beta_{\text{typ}} \beta \gamma}} m_u W \beta \gamma c = 0.45 \text{ GeV/c} \quad 4$$

where the normalized emittance is at least 10π microns, a typical beta value $\beta_{\text{typ}} = 30$ meters, the relativistic factor $\beta\gamma = 108.4$, the atomic weight for gold is $W = 197.0$, and an atomic mass is $m_u = 0.931$ GeV. The rms energy spread in the beam is typically approximately

$$\sigma_E \approx \frac{\sigma_p}{p_0} W (T + m_u) = 17.7 \text{ GeV} \quad 5$$

where the relative momentum spread $\sigma_p/p = 8.9 \times 10^{-4}$ at collision differs from the relative energy spread by the relativistic factor of $\beta \approx 1$.

Since $\langle x'^2 \rangle^{1/2}$ and σ_E are both very much greater than their pair production kick counterparts, and since there are so few such events per gold ion per store, there is no problem with beam diffusion from this source.

Acknowledgements

Thanks to Sebastian White and Mike Harrison for their comments.

References

- 1 J-S Wu, M.J. Rhoades-Brown, C. Bottcher and M.R. Strayer, "Detailed analysis of the background e^+ , e^- spectra generated by relativistic heavy-ion beam crossing", Nuclear Instruments and Methods in Physics Research A311 (1992) 249-257
- 2 Private communications with Sebastian White.