

## Rest Mass of the Fully Stripped Gold Ions

D. Trbojevic

January 1994

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

## Rest Mass of the Fully Stripped Gold Ions

Dejan Trbojevic

January 13, 1994

The mass of an atom is usually expressed as:

$$M(N, Z) = Z * (m_p + m_e) + N * m_n - (BE)_{\text{nucleus}} - (BE)_{\text{electrons}} \quad (1)$$

where N is the number of neutrons,  $m_p$ ,  $m_e$ ,  $m_n$  are the proton, electron, and neutron rest masses, respectively; and  $(BE)_{\text{nucleus}}$  and  $(BE)_{\text{electrons}}$  represent the total binding energies of the nuclear particles and of the atomic electrons, respectively. Atomic mass of the 197 gold is(1):

$${}_{79}\text{Au}^{197} \rightarrow M_{\text{Au}} = 196.966548 \pm .000006 \text{atmu} = 183.47322 \text{GeV}$$

The rest mass of the fully stripped gold ions  $M^*(N, Z)$  is:

$$M^*(N, Z) = M(N, Z) + (BE)_{\text{electrons}} - Z * m_e \quad (2)$$

Values of the fundamental physical constants as: speed of light  $c$ , the electron rest mass  $m_e$ , the atomic mass in GeV are(2):

$$c = 299\,792\,458 \text{m/s}$$

$$m_e = 9.109\,389\,7(54) \pm 0.59 * 10^{-31} \text{kg} \Rightarrow 0.51099906 \text{MeV}$$

$$e = 1.60217733(49) \pm 0.310^{-19} \text{C}$$

$$\text{atmu} = 0.931\,494\,32(28) \pm 0.3 \text{GeV}$$

$$Z * m_e = 79 * 0.51099906 \text{MeV} = 0.0403689257 \text{GeV}$$

The binding energy of the electrons from the K, L, M, and other shells could be estimated from the X-ray emission data (2 electrons in the K shell with 8.7 KeV, 8 electrons in the L shell with 13.0 KeV etc.) as:

$$(BE)_{\text{electrons}} = 0.327\text{MeV} = 0.000327\text{GeV}$$

$$M^*(N,Z) = 183.43318 \text{ GeV} = 196.92356 \text{ amu}$$

The magnetic rigidity  $B\rho$  for the fully stripped gold ions could be calculated if the energy or momentum are known:

$$B\rho = p/Zc = m_{\text{rest}} * \beta\gamma/79c = \left( m_{\text{rest}} \sqrt{\gamma^2 - 1} \right) / (79 * c) \text{ Tm}$$

$$B\rho_{\text{Au}^{79}} = 7.7451547 * \sqrt{\gamma^2 - 1} (\text{Tm})$$

## References

1. A.H. Wapstra and N.B. Cover, Nuclear Data Table 9, 265 (1971).
2. E. Richard Cohen and Barry N. Taylor, Physics Today, August 1992, part 2, pp. 9–14.