



Brookhaven
National Laboratory

BNL-101657-2014-TECH

RHIC/AP/1;BNL-101657-2013-IR

Preliminary Chromatic Correction

A. Antillon

July 1984

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy
USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Preliminary Chromatic Correction

for the RHIC2

Lattice Design


Armando Antillon
BNL, July 3, 1984

ABSTRACT

A lattice with the name of RHIC2 has been considered as a candidate for the Relativistic Heavy Ion Collider. In this note we study its chromatic properties in two cases: 1) when two families of sextupoles are included, 2) when special families of sextupoles are placed in the arcs of the ring in order to diminish higher order resonance effects and chromatic effects.

1.- Description of the RHIC2 lattice.

This lattice consists of 6 arcs and 6 insertions with supersymmetry 3. Each arc is made of 12 cells with the next structure



where 1, 2 and 3 mean three different drift spaces.

One of the insertions has the next arrangement

Insertion = $I = Q7I \ Q6^* \ Q6I \ BS \ Q5I \ BS \ Q4I \ Q3I \ Q2I \ Q1I \ BC2I \ BC1I$
 $CR \ BC10 \ BC20 \ Q10 \ Q20 \ Q30 \ Q40 \ BS \ Q50 \ BS \ Q60 \ Q60^* \ Q70$

where BS is dispersion supresor dipole and the dipoles BC1I, BC10 allow head-on colisions.

Each superperiod is made in the following way

$$6 \times C_I + I_1 + 12 \times C_o + I_1^{-1} + 6 \times C_I$$

where I_1^{-1} means mirror of I_1 and the cells C_I differ from C_o in the drift spaces between bending magnets and quadrupoles.

Some of the machine functions are

in the arcs

	β_x	β_y	η_x
QF	51.5	7.4	1.38
QD	7.4	51.5	0.69

in the insertions

$$\begin{aligned}\beta_x^* &= 17.7m \\ \beta_y^* &= 2.99m \\ \beta_{x\max} &= 267.3m \\ \beta_{y\max} &= 667.4m\end{aligned}$$

The natural chromaticity has values

$$\xi_x = -51.0 \quad , \quad \xi_y = -69.9$$

and the nominal tune is $\nu_x = 31.61521 \quad , \quad \nu_y = 31.61399$

2.- Chromatic Properties with two families of sextupoles.


In order to avoid the main resonances near the nominal tune, the chromaticity has been chosen to be 1, even without knowing if this value is compatible with head-tail instabilities. Next tables show the momentum dependence of some machine functions. The integrated strengths of the sextupoles ($\frac{B'' \ell_{\text{sex}}}{B_0 f} m^{-2}$) are in this case:

$$SF = -.14352$$

$$SD = +.42412$$

All calculations have been done for $\left| \frac{\Delta p}{p} \right| \leq 0.5\% .$

v_x



v_y




Table I.- Tune variations with momentum. Two families of sextupoles are considered. Straight line is chromaticity 1. $v_x = 31.61521$, $v_y = 31.61399$.





Table II.- Variation of max in insertions. $\beta_{ox} = 267.322\text{m}$ and
 $\beta_{oy} = 667.463\text{m}$.






Table III.—Behavior of β_x , β_y , η in the crossing point.




Table IV.- Variation of momentum with amplitud with resolution of 100 turns. Points a, b, c represent emittances $0, 1.5\pi \times 10^{-6} \text{ m-rad}$, $3\pi \times 10^{-6} \text{ m-rad}$ respectively. We kept chromaticity 1 in order to avoid 8th and 10th order resonances.

With the previous characteristic, the phase space shows the next structure for 100 turns. When the number of turns is smaller it is possible to identify high order resonances.




Table Va.

ϵ		X DIRECTION	Y DIRECTION
1.5 π $\times 10^6$ m-rad			
3 π $\times 10^6$ m-rad			

Table Vb.




Table Vc.

Table V.- a) x and y phase spaces for 100 turns. Momentum 0%. b) Momentum 0.5%. The diffusion of the states in phase spaces are consequence of higher order resonances. c) Momentum -0.5%. Direction of phase space is also affected by higher order resonances.

3.- Chromatic Properties with special families of sextupoles.


The phase spaces of table V and the large tune dependence with amplitude, shown in table IV, have been suggested that the higher order resonance effects appear as consequence of aberration phenomena². In order to correct phase spaces without changing the chromatic behavior of part 3, we have tried to put some special sextupoles in insertions, in places where dispersion is practically zero. The attempt was no successful.

Another attempt to diminish the higher order resonance effects has been made by improving the chromatic behavior. This has been done as we now describe.

We know^{3,4} that $\Delta\beta/\beta$ change with $\Delta p/p$ as

$$\frac{\Delta\beta}{\beta}(\phi, \frac{\Delta p}{p}) = 2\zeta^1(\phi) \frac{\Delta p}{p} + (\zeta^1(\phi)^2 + 2\zeta^2(\phi)) \left(\frac{\Delta p}{p}\right)^2 + \dots$$

We will assume that $2\zeta^2 \ll \zeta^1^2$. With this, $(\Delta\beta/\beta)_{\max}$ will decrease if ζ^1_{\max} is minimized. This has been done, in both x and y direction, by changing the SF, SD sextupoles, to values Z, A in places indicated.



The new values for SF, SD are slightly different from part 3.

$$SF = -.14109 , SD = +.42894 \text{ and}$$

$$Z1 = -.34524 , Z2 = .00007$$

$$A1 = .2 , A2 = .6$$

With this special sextupoles, the machine functiones are




Table VI.- Behavior of tune with momentum when special sextupoles are placed in the arcs. SF and SD have to change lightly in order to keep chromaticity 1.




Table VII.- β_{\max} in insertions is better with special sextupoles.

$$\beta_x = 267.322 \text{ and } \beta_y = 667.463.$$




Table VIII.- In the crossing points β and η remain basically the same with and without special sextupoles.

The tune diagram still shows large tune dependence with amplitude, but smaller than in the previous case.




Table IX.- Tune variation with amplitude is a little bit better. Points a, b, c represent emittances $0, 1.5\pi \times 10^{-6}$ m-rad, $3\pi \times 10^{-6}$ m-rad. respectively.

Finally, phase space is much better with special sextupoles than only with SF, SD.





EMITTANCE	X DIRECTION	Y DIRECTION
1.5π $\times 10^{-6}$ m-rad		
3π $\times 10^{-6}$ m-rad		

Table Xa.





EMITTANCE	X DIRECTION	Y DIRECTION
$1.5 \pi \times 10^{-6}$ m-rad		
$3 \pi \times 10^{-6}$ m-rad		

Table Xb.

EMITTANCE	X DIRECTION	Y DIRECTION
1.5π $\times 10^{-6}$ m-rad		
3π $\times 10^{-6}$ m-rad		

Table Xc.

Table X.- a) x and y phase spaces for 100 turns and special sextupoles.
 Momentum 0%. b) Momentum 0.5%. Diffusion of points of phase
 space have been partially corrected. c) Momentum -.05%. Direc-
 tion of phase space have been also partially corrected with
 special sextupoles.

4.- Summary.

We have used special families of sextupoles in the arcs of RHIC2 in order to reduce discomfort in phase space of higher order resonances. In general, chromatic behavior is better than with two families of sextupoles.

Bibliography.

- 1.- J. Claus, Private communication (1984)
- 2.- A. Ruggiero, Private Communication (1984)
- 3.- E.D. Courant and H.S.Snyder, Ann. Phys. 3, 1, (1958)
- 4.- A. Antillon, BNL-34615 (1984)

911 A. Antillon

902 D. Brown

J. Claus

E. Courant

F. Dell

C. Goodzeit

H. Hahn

H. Halama

J. Kats

S. Y. Lee

R. Louttit

G. Morgan

G. Parzen

L. Roberts

R. Shutt

P. Thompson

E. Willen