

Aperture Requirements and Proton Performance in RHIC

G. Parzen

October 1984

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

APERTURE REQUIREMENTS
AND
PROTON PERFORMANCE IN RHIC

G. Parzen

10/3/84

and
Proton Performance in RHIC

G. Parzen, 10/3/84

Important Limitations

- 1) Intra beam scattering
- 2) Beam-Beam interaction
- 3) Instabilities, $Z_{||}/n$, Z_{\perp}/n
- 4) Aperture (closed orbit errors, random $\Delta B/B$, chromatic $\Delta B/B$)

Intra beam Scattering, $N_b = 10^8$ / bunch

- 1) Longitudinal growth
- 2) Transvers growth
- 3) aperture required $\begin{cases} \sigma = 30, \pm 26 \text{ mm} \\ \sigma = 320, \pm 320 \text{ mm} \end{cases}$
- 4) $t = 50$ hr operation
- 5) Luminosity

Protons


2.5 δ vs γ :

$N_b = 1 \times 10^{11}$

$A_{rf} = .3 \text{ ev-sec}$

$V = 1.2 \text{ MV}$

$h = 6 \times 57$




$\Delta p/p \text{ bucket}$

$t=50$

$t=10$

$t=0$

Protons
 E_N vs γ
 $N_b = 1 \times 10^{11}$




Half - Aperture vs δ ($6\sigma_H + 2.5\lambda_p\delta$ vs δ)

Protons, $N_b = 1 \times 10^{10}$


Required Aperture

- 1) $\delta = 30, \pm 26 \text{ mm}$
- 2) $\delta = 320, \pm 8 \text{ mm}$


} for $t = 10 \text{ hrs}$,
 includes closed orbit
 and $\Delta B/\beta$ effects.



Average Luminosity vs γ
Protons, $N_b = 1 \times 10^{11}$



σ_R vs γ
Protons, $N_b = 1 \times 10^{11}$



Beam Beam Interaction

$$N_b = 1 \times 10^{11}, \quad \epsilon = 20 \times 10^{-6}$$

$$\Delta V_{BB} = .0052, \quad \text{head on collisions, } t=0$$

$$\Delta V \sim N_b / \epsilon N \quad \text{independent of energy}$$

Instabilities, $N_b = 10^{11}$

$$\gamma = 320, \quad \sigma_{\text{rel}} = 15.8 \text{ cms}, \quad \delta_0 = .102 \times 10^{-3}$$

$$A_{\text{eff}} = .3 \text{ ev-sec}, \quad I_{\text{pk}} = 12 \text{ A}, \quad I_{\text{av}} = .43 \text{ A}$$

$$Z/n = 10 \text{ ohms}$$

$$N_{b,T} = .5 \times 10^{11} \quad (\text{Threshold})$$

(AGR says 1×10^{11})

$$N_{b,T} \sim \delta^3$$

$$\left(\begin{array}{l} I_p < \frac{4\pi E}{e} |\eta| \frac{\delta^2}{(Z/n)} \\ I_p = N_b e \beta c / \sqrt{2\pi} \sigma_r \end{array} \right)$$

$$\delta = .1 \times 10^{-3} \rightarrow .13 \times 10^{-3}$$

takes 1/3 hour due to FBS

(8)

$$N_{b,T} (\text{Transverse}) = 2.9 \times 10^{11} \quad \text{for } \delta = .13 \times 10^{-3}$$

Problem if $N_b > N_{b,T} (\text{Transverse})$

$$\text{Transverse } I_p < \frac{10 \epsilon}{\rho} \ln \left| \frac{\delta}{B} \right| \frac{1}{(Z_1/n)}$$

$$N_{b,T} \sim \delta^2$$

May require δ increase by other means.

$N_b = 10^{12}$

Intrabeam Scattering

$\int \rightarrow \cdot 4.76 \times 10^{-3}$ (t = 10 hrs) , $\delta = 320$
 $2.5 \delta = 1.4 \times 10^{-3}$, $(\frac{\Delta p}{p})_{\text{bucket}} = 2.3 \times 10^{-3}$

Aperture required = $7.8 \text{ mm} + 2.5 \text{ mm}$
= 10.3 mm

Beam-Beam Interaction

$\Delta V \sim \frac{N_b}{E_N}$ $L_0 \sim \frac{N_b^2 \delta}{E_N}$

To keep , $\Delta V = .0052$, $E_N \rightarrow 10 \times 20 = 200 \times 10^{-6}$

$\sigma_H = 2.3 \text{ mm}$

$\delta = .48 \times 10^{-3}$, $X_p \delta = .78 \text{ mm}$

Aperture Required = $6 \sigma_H + 2.5 X_p \delta = 16 \text{ mm}$

With closed orb, t and P/B effect

Aperture Required = $\pm 20.5 \text{ mm}$

(7/6)

$N_b = 10^{12}$, Instabilities

$$z/h = 10, \quad N_{b,T} = .5 \times 10^{11}$$

$$\delta = .11 \times 10^{-3} \rightarrow .27 \times 10^{-3}$$

(would take $\frac{1}{2}$ hr due to IBS)

$$N_{b,T} \text{ (transverse)} = 1.3 \times 10^{12} \text{ for } \delta = .27 \times 10^{-3}$$

Luminosity, $N_b = 10^{12}$, $\delta = 320$

$$L_0 = 13 \times 10^{32}, \quad L = 1.3 \times 10^{32} \text{ at end.}$$

$$L_{AV} \approx 2 \times 10^{32}$$

Added Notes

- 1) The result used for the threshold current for the longitudinal instability is in error by a factor of 2. This was pointed out by Harald Hehn. The result should read

$$I_p < \frac{2\pi E}{e} \left| \eta \right| \frac{\delta^2}{(Z_{||}/n)}$$

and the numerical results should be modified by this factor of 2.

- 2) One possibly important effect has not yet been looked at in regard to operating RHIC with $N_b = 1 \times 10^{12}$ protons/bunch. This is the incoherent space charge effect, which is the ν -shift caused by the charge and current of the beam.