

## Beam-Beam Limited Luminosity

H. Hahn

February 1985

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Beam-Beam Limited Luminosity

H. Hahn

BNL-2/11/85

BEAM-BEAM LIMITED LUMINOSITY

Assumption and definitions:

$$\alpha = 0$$

$$\epsilon_H = \epsilon_V = \epsilon_N$$

$$\langle \beta \rangle^2 = \beta_H \beta_V$$

$$\sigma_{H,V} = \left( \frac{\epsilon \beta_{H,V}}{6\pi \gamma} \right)^{1/2}$$

$$r = \frac{\sigma_H}{\sigma_V} \quad (0 \leq r \leq 1)$$

Luminosity

$$L = \frac{3}{2} f_o h_B N_B^2 \frac{\gamma - 1}{\epsilon \langle \beta \rangle}$$

with  $f_o$  = rotation frequency  
 $h_B$  = number of bunches  
 $N_B$  = ions/bunch

Tuneshift (RHIC-PG-26)

$$\Delta v_V = 3 r_p \frac{Q^2 N_B}{A \epsilon_N} \frac{1}{1+r}$$

$$\Delta v_H = 3 r_V \frac{Q^2 N_B}{V A \epsilon_N} \frac{1}{1+r^{-1}}$$

Luminosity limit:  $(\sigma_H \leq \sigma_V)$

$$\frac{L}{\Delta v_V} = \frac{f_o \cdot h_B}{2 \cdot r_p} \cdot \frac{N_B \cdot Y}{\langle \beta \rangle} \cdot (1 + r)$$

RHIC lattice

-- symmetric:  $r = \left( \frac{1.44 \text{ m}}{6.50 \text{ m}} \right)^{1/2} = 0.47$

-- asymmetric:  $r = 1$

$$\left( \frac{L_{\text{asym}}}{L_{\text{sym}}} \right)_{\text{Beam-Beam}} = 1.36$$