

## BNL-101677-2014-TECH RHIC/AP/21;BNL-101677-2013-IR

A Chromatic Correction Scheme for the Antisymmetric RHIC Lattice. The First Approximation.

A. Antillon

March 1985

Collider Accelerator Department Brookhaven National Laboratory

## **U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

# A Chromatic Correction Scheme for the Antisymmetric RHIC Lattice. The First Approximation.

Armando Antillon

March 7, 1985

#### Abstract

We use special families of sextupoles in the arcs, with antisymmetric distribution. The chromatic behavior of the machine functions are , in general, better than with only two families.

ł

#### I. Introduction

In reference 1 we used a symmetric sextupole scheme for correcting the symmetric RHIC3 lattice. There, we suggested that an antisymmetric scheme can correct an antisymmetric lattice. In this work we explore such idea for the current RHIC lattice<sup>2</sup>, and without optimizing the scheme we see that, in general, chromatic effects are reduced.

#### II. Description of the RHIC Lattice

Each typical cell has the structure



where  $X_1, X_2, \dots$  will be special sextupoles that we can use. SF and SD adjust the chromaticity to the desired value.

The insertions are made as follows.

I = Q8I Q7I Q6I BS2 Q5I BS1 Q4I QXI Q3I Q2I Q1I BC2I BC1I CR BC10 BC20 Q10 Q20 Q30 QX0 Q40 BS1 Q50 BS2 Q60 Q70 Q80

In the arcs, for  $\Delta p/p = 0$ ,  $\beta$  and  $\eta$  have values

	β <sub>x</sub>	<sup>β</sup> y	n x
QF	49.6	8.8	1.5
QD	8.8	49.6	.7

Natural Chromaticity: 
$$\xi_x = -56.6$$
,  $\xi_y = -56.5$   
tunes:  $v_x = 28.40867$ ,  $v_y = 28.37187$   
Sextupoles: SF = -.15194, SD = +.3111 for  $\xi_x = \xi_y = 1$   
 $\beta_x^* = \beta_y^* = 3.00001$ 

#### Special Families of Sextupoles

In Figure 5 we show the distribution of the sextupole. There are four families in one arc with a total number of eight families. The dashed lines in all the figures correspond to the values of sextupoles that we next are giving and obviously they have to be optimized by Harmon<sup>3</sup> or SYNCH<sup>4</sup>. On the other hand, the scheme must be optimized to reduce the phase space distortions related with the linear contribution to the W-vector introduced by Guignard<sup>5</sup>. For the moment, the four families per arc is in accordance with the number he suggests for a 90° lattice.

For one arc:

Family	one:	S80=D2=H2=Z2= -•04934
Family	two:	B2=F2=J2= .045
Family	three:	A2=E2=I2= .03533
Family	four:	C2=G2=K2= -•15268

For the other arc:

Family	five:	S8I=D1=H1=Z1= .01953
Family	six:	B1=F1=J1= .0037
Family	seven:	A1=E1=I1= -•075
Family	eight:	C1=G1=K1= .1120

The effective sextupole will be SF or SD plus one of the above ones. We see that the larger sextupole is about 1.5 larger than SD. For these values, SF and SD have to be readjusted to keep  $\xi_x = \xi_y = 1$ . The new values are

```
SF = -.1301
SD = +.3052
```

The change with respect to the original values is small.

- 3. G. Guignard reported at Sardinia School very good results using Harmon.
- 4. J. Claus, private communication.
- 5. G. Guignard. Lecture given at Sardinia School, March 1985.

<sup>1.</sup> A. Antillon, RHIC-8, BNL (1985).

<sup>2.</sup> S. Y. Lee, private communication.



Figure 1.



Figure 2.  $\Delta\beta/\beta$  maximum in insertions.





Figure 4. Dispersion at crossing points.

-







Figure 7.  $\beta_x$ ,  $\beta_y$  for  $\Delta p/p = +1.4$  and special families.

0

. 83

