

Location of Sextupole Magnets

H. Hahn

May 1986

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Location of Sextupole Magnets

H. Hahn

BNL

May 12, 1986

Location of Sextupole Magnets

H. Hahn

The correction of the chromatic effects in the RHIC lattice is achieved with sextupoles at each arc quadrupole.^{1,2} There are 25 quadrupole units per arc (23 arc quadrupoles + 1/2 Q9 + 1/2 Q9). However, in order to avoid half-length sextupoles, 24 sextupole units of equal length will be installed in each arc. One has therefore, several options for the location of the sextupoles. The following ground rules were applied to find a unique solution:

1. The 2 rings look identical with respect to its circulating beam.
2. The antisymmetry with respect to the crossing point is retained.
3. The sextupoles of inner and outer rings are at the same azimuthal location.
4. The 24 sextupoles within each arc are uniformly spaced.
5. The four Q9 quadrupoles at the injection area (6 o'clock) and consequently at 2 and 10 o'clock remain without sextupoles.
6. The dipole correctors at the Q9 quadrupoles in the 2, 6, and 10 o'clock insertions are on the arc side, thereby locating all correctors and sextupoles in the entire arc.

-
1. G. Parzen, "Correction of chromatic effects in the β -functions in RHIC", RHIC-AP-28 (1986).
 2. S. Y. Lee, "Chromatic correction for the RHIC lattice", RHIC-17 (1986).