

Some Comparisons Of HIJET With Data

T. Ludlam

February 1986

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

SOME COMPARISONS OF HIJET
WITH DATA

T. Ludlam

February 24, 1986

SOME COMPARISONS OF HIJET
WITH DATA

Thomas W. Ludlam

I. Introduction

This note compares HIJET 1.0 with some data samples from p-p, p-A and p-nucleus collisions, with the purpose of trying to gain a comprehensive view of how well the model reproduces well-known features of high energy interactions. We are not trying to break new physics ground here, nor are we trying to make detailed fits (no parameters are adjusted).

II. Proton-proton Collisions

Here we are looking strictly at ISAJET, which does the nucleon-nucleon interactions in HIJET. HIJET uses the MINBIAS routine of ISAJET, and uses it at much lower energies than the collider range that ISAJET was designed for and for which it has been extensively tested by high energy users. For HIJET the interesting range of nucleon-nucleon collisions includes laboratory momenta roughly from 1 to 100 GeV/c.

In Fig. 1 we see that the ISAJET charged particle multiplicity agrees very well with data over this momentum range. Figure 2 shows that the higher moments of the multiplicity distributions are also in good agreement with experiment, agreeing well with the empirical KNO scaling curve.

Figure 3 shows the x distribution for inclusive proton production over the interesting range of energies. The model gives a good accounting of the data except near $x=1.0$, where ISAJET does not include the diffractive peak seen in the data. Figures 4a, b show the invariant cross section for proton production as a function of x and P_T . Again, ISAJET gives good agreement with the data, with the exception of the diffractive peak at $x=1$.

The rapidity spectra for charged particle production are compared with data at 200 GeV/c in Fig. 5. The data [de Marzo et al., Phys. Lett. 112B, 173 (1982)] are well reproduced.

Finally, Fig. 6 shows the E_T spectrum measured by the NA5 calorimeter at the CERN SPS. This is well reproduced by ISAJET.

We may conclude that the essential dynamics of nucleon-nucleon interactions are very well described by ISAJET, down to the lowest energies of interest for HIJET. The kinematic aspects are done precisely: energy and momentum are rigorously conserved in each collision. The detailed accounting of quantum number flow is not done in ISAJET's MINBIAS routine, except in an average way. Charge and strangeness need not balance exactly in a given event (baryon number will). The K/π ratio is roughly right at high energies

of interest for HIJET. The kinematic aspects are done precisely: energy and momentum are rigorously conserved in each collision. The detailed accounting of quantum number flow is not done in ISAJET's MINBIAS routine, except in an average way. Charge and strangeness need not balance exactly in a given event (baryon number will). The K/π ratio is roughly right at high energies ($\sqrt{s} \geq 50$ GeV), but there will be too many strange particles to agree with the data at fixed-target energies.

III. Proton-Nucleus Collisions

The mean charged-particle multiplicity for proton-nucleus collisions is shown in Fig. 7, for two different energies, as a function of A . Figures 8 and 9 show the charged particle rapidity distributions obtained in the NA5 streamer chamber at the CERN SPS, with argon and xenon, respectively, as the target nuclei [de Marzo et al. Phys. Rev. D26, 1019 (1982)]. HIJET agrees remarkably well with these data. It is this agreement which has been the primary motivation for the widespread use of HIJET as a guide to the expected particle production in the design of high energy nuclear beams experiments.

Figure 10 shows the mean number of collisions per p-nucleus interaction as a function of A . This is somewhat larger in HIJET than the numbers usually extracted from the data (solid line), the effect being greater at small A than at large A . This is due in part to the hard sphere, uniform density distribution employed in HIJET.

An important issue is the so-called "stopping power": the distribution of leading baryons in p-nucleus collisions. In Fig. 11 the HIJET result is compared with the much-studied Fermilab data sample of Barton et al. It will be seen that the agreement is very good for the heavy nuclear targets. For the light nucleus (carbon) the HIJET result is high by about a factor of two in the range of x covered by the data. Note (Fig. 4) that for a proton target the agreement is again very good.

In general, HIJET seems to do very well in reproducing the average values of measured quantities in p-nucleus collisions. It does not do so well in reproducing the fluctuations seen in the data. This is evident in Fig. 12, where the dispersion of the multiplicity distribution is plotted against the mean. The calculated result is in complete agreement with the data for the proton-proton case, but the multiplicity distributions from HIJET are significantly narrower than the data for the heavier nuclei. A similar, and quite possibly related effect is seen in Fig. 13. Here the transverse energy (E_T) spectrum is shown in 200 GeV/c p-pb collisions as measured by the HELIOS collaboration at the SPS. This distribution is very broad (and therefore very interesting!), and not at all well reproduced by HIJET. The p-p spectrum shown in this figure is the same data as in Fig. 6, scaled to the rapidity interval of the HELIOS experiment. Here, as seen in Fig. 6, the HIJET (ISAJET) result reproduces the data extremely well.

IV. Nucleus-Nucleus Collisions


There is very little data with which to evaluate the HIJET extrapolation from the p-nucleus case of nucleus-nucleus collisions. The comparisons which can be made are encouraging. Two examples are shown here. Figure 14 shows the pseudo-rapidity distribution for charged particles produced in collisions of α -particles, accelerated in the CERN PS accelerator, with an emulsion target. Central collisions are selected on the basis of the number of grey tracks observed. The HIJET result, calculated for impact parameters less than .5 fermi, agrees quite well with both the shape (angular distribution) and area (total charged particle multiplicity) of the measured curve.

Figure 15 shows a well-known cosmic ray event observed by the JACEE collaboration (Phys. Rev. Lett. 50, 2062). The pseudo-rapidity distribution is plotted for the 1015 charged particles produced in this collision, for which the projectile was identified as a silicon nucleus with an energy of about 5000 GeV/amu. The HIJET result shown is an average over 50 such collisions, with impact parameters less than .5 fermi. The mean number of charged particles per event in this sample is 946, and it is evident that these HIJET events are very similar to the observed event.

FIG. 1

AVERAGE CHARGED PARTICLE Multiplicity
IN P-P COLLISIONS

— EMPIRICAL FIT TO DATA [THOMÉ ET AL., Nucl. Phys. B. 129, 365]
X HIJET (ISAJET)




KNO SCALING

EMPIRICAL FIT
TO PP DATA

- (ISAJET)
HIJET
- x $\sqrt{s} = 13.8 \text{ GeV}$ [$P_{\text{LAB}} = 100 \text{ GeV}/c$]
 - o $\sqrt{s} = 6.3 \text{ GeV}$ [$P_{\text{LAB}} = 20 \text{ GeV}/c$]
 - $\sqrt{s} = 4.5 \text{ GeV}$ [$P_{\text{LAB}} = 10 \text{ GeV}/c$]



FIG. 2



DATA →

WHITMORE,
PHYS REP. 10, 273

FIG. 3



INCLUSIVE PROTON SPECTRA:
PP → P + ...




Fig. 50 Invariant cross sections for the reaction $pp \rightarrow p + \text{anything}$ as a function of $x = 2p_L^*/\sqrt{s}$ for fixed values of the transverse momentum.

J. WHITMORE, *PHYS. REP.* 10, 273 (1974)

Fig. 4a

PP → P + ANYTHING

100 GeV/c

$\sqrt{s} = 13.8$

HIJET
(ISAJET)

$E \frac{d^3\sigma}{dP^3} (mb/GeV^2)$







FIG. 5

200 GeV/c PP



46 6213

PHOTODUPLICATION SERVICE, NATIONAL BUREAU OF STANDARDS, GAITHERSBURG, MARYLAND 20899



P-A: CHARGED PARTICLE MULTIPLICITY

- Δ CERN SPS (de MARZO ET AL.)
- \circ FERMI LAB (ELIAS ET AL.)
- X HIJET





FIG. 7

200 GeV/c p-Am




FIG. 8

200 GeV/c P-Xe





FIG. 9




FIG. 10

$$E \frac{d^3 \psi}{dr^3} \left[\frac{mb}{(G_1 v/c)^2} \right]$$




FIG. 11

200 GeV/c P-A
 CHARGED PARTICLE Multiplicity DISTRIBUTION:
 DISPERSION VS. MEAN

- DATA [NA-5]
- x HIJET




FIG. 12

FIG. 13

HELIOS TEST 1984

CERN SPS

P(200 GeV) PG

$\frac{d\sigma}{dE_T} [mb/GeV]$

HIJET MONTE CARLO

P-P DATA \rightarrow

P-P $\sqrt{s} = 19.4$

P-(5m) $\sqrt{s} = 45$





Fig. 14

α -Ag 12A GeV/c Pseudo Rapidity
CENTRAL COLLISIONS

x HIJET
— DATA: LUND/INDIA EMULSION COLLAB.



JACEE Collab.


Si + Emulsion (AgBr)

1 EVENT

1015 Charged particles

$$\sqrt{s}/A =$$

50 + 50 GeV/nucleon



~~X X~~ HIJET Si + Ag
AVERAGE OVER 50
CENTRAL COLLISIONS
<N_{ch}> = 946

Fig. 15