

## Errors in Siberian Snakes

E. D. Courant

August 1993

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

BNL-49436  
AD/RHIC-121  
Informal Report

## **Errors in Siberian Snakes**

E. D. Courant

August 1993

## **R H I C   P R O J E C T**

Brookhaven National Laboratory  
Associated Universities, Inc.  
Upton, NY 11973

Under Contract No. DE-AC02-76CH00016 with the  
UNITED STATES DEPARTMENT OF ENERGY

# ERRORS IN SIBERIAN SNAKES

E D Courant

Aug. 18, 1993

In an accelerator or storage ring for polarized protons, "Siberian snakes" may be inserted in the lattice in order to stabilize the spin dynamics, i.e. to prevent betatron oscillations and orbit distortions from depolarizing the beam.

This is accomplished by inserting an odd number  $N_p$  of pairs of snakes, spaced  $\pi/N_p$  apart in orbit azimuth; each snake is a spin rotator that rotates the spin by exactly 180 degrees about a precession axis in the horizontal plane. The precession axes of the two snakes of a pair should be exactly 90 degrees apart (e.g. longitudinal and transverse horizontal, or  $\pm 45$  degrees from longitudinal). The preferred choice in most cases is  $\pm 45$  degrees.

For RHIC, one pair of snakes ( $N_p = 1$ ) is planned, 180 degrees apart with  $\pm 45$  degree axes; in the case of the Tevatron,  $N_p = 3$  (six snakes) is envisaged, with the snakes spaced 60 degrees apart.

The spin rotation matrix for a snake with its axis at  $\pm 45$  degrees is

$$S_{\pm} = i \frac{(\sigma_1 \pm \sigma_2)}{\sqrt{2}} = i\sigma_1 e^{\pm i\pi\sigma_3/4} \quad (1)$$

while the matrix for an arc with bending angle  $\vartheta$  is

$$A = e^{i\gamma G \vartheta \sigma_3/2} \quad (2)$$

with  $\vartheta = \pi/N_p$  and  $G = 1.7928$  the proton anomalous moment, so that the matrix for the whole ring (with no imperfections) is

$$M = (S_+ A S_- A)^{N_p} = (i\sigma_3)^{N_p} \quad (3)$$

independent of energy. The spin tune  $\nu_{sp}$  is determined by

$$\cos \pi \nu_{sp} = \text{Tr } M / 2 \quad (4)$$

which equals 0, i.e.  $\nu_{sp} = \frac{1}{2}$  when  $M$  is given by (3) with odd  $N_p$ ; the stable spin direction is exactly vertical.

Now what if the snakes are not perfect? We consider three types of imperfections:

(a) The rotation angle of each snake is not exactly 180 degrees, but rather  $\pi + \delta$ .

(b) The precession axes of the snakes are not exactly at 45 degrees from the longitudinal, but rather  $\pm\pi/4 + \beta$ .

(c) The bending arcs between the snakes are not exactly  $\pi/N_p$ , i. e. each snake is azimuthally displaced by a bending angle  $\epsilon$ .

In case (a) (incorrect precession angle) the matrix for the  $k$ -th snake becomes

$$-\sin \delta_k/2 + i \frac{(\sigma_1 \pm \sigma_2)}{\sqrt{2}} \cos \delta_k/2 \quad (5)$$

To lowest order in the  $\delta$ 's, the overall matrix is then

$$i\sigma_3 + \frac{i}{2}\delta_1 \frac{(\sigma_1 - \sigma_2)}{\sqrt{2}} + \frac{i}{2}\delta_2 \frac{(\sigma_1 + \sigma_2)}{\sqrt{2}} e^{i\pi\gamma G\sigma_3} \quad (6)$$

for the case of two snakes (RHIC), and

$$-i\sigma_3 - \frac{i}{2}(\delta_1 + \delta_3 + \delta_5) \frac{(\sigma_1 - \sigma_2)}{\sqrt{2}} - \frac{i}{2}(\delta_2 + \delta_4 + \delta_6) \frac{(\sigma_1 + \sigma_2)}{\sqrt{2}} e^{i\pi\gamma G\sigma_3} \quad (7)$$

for the case of six snakes (Tevatron). Thus, to lowest order in the deviations from 180 degree in the rotators, the spin tune is still exactly  $\frac{1}{2}$ , but the axis of the stable spin direction is tilted from the vertical by an angle of order  $\delta$ . Thus if the rotators are off by a degree, the spin axis is tilted by a similar angle, which should be tolerable.

In case (b) (precession axes deviate from 45 degrees by  $\beta$ ) the snake matrices become, instead of (1),

$$S_k = i\sigma_1 e^{i(\pm\pi/4 + \beta_k)\sigma_3} \quad (8)$$

and the overall matrix is

$$(i\sigma_3)^{N_p} e^{-i(\beta_1 - \beta_2 + \beta_3 - \beta_4 + \beta_5 - \beta_6)\sigma_3} \quad (9)$$

The trace of this matrix deviates from zero, and we find that the spin tune becomes

$$\nu_{sp} = \frac{1}{2} \pm (\beta_1 - \beta_2 + \beta_3 - \beta_4 + \beta_5 - \beta_6)/\pi, \quad (10)$$

i.e. the spin tune deviates from  $\frac{1}{2}$  by an amount of the order of  $\beta$  (about 1% if the axes are off by a degree, which should be tolerable).

Finally, in case (c), where the azimuthal locations of the snakes are not evenly spaced: If the location of a snake is off by a bending angle  $\epsilon$ , the spin matrix for the displaced snake is

$$e^{i\gamma G \epsilon_k \sigma_3/2} S_{\pm} e^{-i\gamma G \epsilon_k \sigma_3/2} = S_{\pm} e^{-i\gamma G \epsilon_k \sigma_3} \quad (11)$$

and the spin tune is

$$\nu_{sp} = \frac{1}{2} \pm (\epsilon_1 - \epsilon_2 + \epsilon_3 - \epsilon_4 + \epsilon_5 - \epsilon_6) \gamma G / \pi . \quad (12)$$

Because of the factor  $\gamma G$  the spin tune is very sensitive to this error at high energy. If we wish the spin tune to be within .05 units of  $1/2$ , then for RHIC ( $\gamma G = 480$  at 250 GeV) we must have  $|\epsilon_1 - \epsilon_2| < .02$  degrees = 0.3 milliradians; for the Tevatron ( $\gamma G = 1920$ ) the alternating sum of the position errors must be .005 degrees or .08 milliradians of bending. These tolerances sound tight, but they will be met automatically if the orbit correction system of the ring corrects the closed orbit errors to better than a few millimeters.