

Luminosity Loss Rate Due To Intrabeam Scattering

A. G. Ruggiero

December 1983

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

LUMINOSITY LOSS RATE
DUE
TO INTRABEAM SCATTERING

A. G. RUGGIERO

(BNL, December 11, 1983)

RHIC-PG-21

luminosity loss rate
due
to Intra-beam Scattering

A. G. Ruggiers

BNL, December 11, 1983

In the limit of short interaction region the luminosity can be very well represented by

$$L = \frac{N^2 B f_{rev}}{4\pi \sigma_v^* \sigma_H^*}$$

where, for identical colliding beams,

- N , number of particles per bunch
- B , number of bunches per beam
- f_{rev} , revolution frequency
- σ_v^* , vertical rms beam size at the collision point
- σ_H^* , horizontal rms beam size at the collision point

assuming crossing ~~at~~ at an angle on the horizontal plane

$$f = \sqrt{1 + p^2}$$

and

$$p = \frac{\alpha \sigma_e}{2 \sigma_H^*}$$

(2)

α , total crossing angle
 σ_e , rms bunch length.

Intrabeam scattering has the effect of having σ_v^* , σ_H^* and σ_E , the r.m.s. energy spread to vary, generally by growing.

~~Obviously~~ Define the luminosity loss rate

$$\tau_L^{-1} = -\frac{1}{L} \frac{dL}{dt}$$

Obviously:

$$\tau_L^{-1} = \frac{1}{\sigma_v^*} \frac{d\sigma_v^*}{dt} + \frac{1}{\sigma_H^*} \frac{d\sigma_H^*}{dt} + \frac{1}{f} \frac{df}{dt}$$

One can work out that

$$\frac{1}{f} \frac{df}{dt} = \frac{P^2}{f^2} \left[\frac{1}{\sigma_e} \frac{d\sigma_e}{dt} - \frac{1}{\sigma_H^*} \frac{d\sigma_H^*}{dt} \right]$$

Assumptions:

1. The bunch length increases (or decreases) at the same rate the energy spread does, that is

$$\frac{1}{\sigma_e} \frac{d\sigma_e}{dt} = \tau_e^{-1}$$

2. We assume that the two modes of oscillations (H and V) are fully coupled to each other on a time shorter than the intrabeam scattering diffusion ~~is~~ times, that is

$$\frac{1}{\sigma_v^\alpha} \frac{d\sigma_v^\alpha}{dt} = \frac{1}{\sigma_H^\alpha} \frac{d\sigma_H^\alpha}{dt} = \tau_H^{-1} + \tau_V^{-1}$$

Here τ_e^{-1} , τ_H^{-1} and τ_V^{-1} are the diffusion (or damping rates) from intrabeam scattering respectively in energy, horizontal (betatron) size and vertical (betatron) size.

In conclusion

$$\tau_L^{-1} = \frac{p^2}{f^2} \tau_E^{-1} + \left(2 - \frac{p^2}{f^2}\right) (\tau_H^{-1} + \tau_V^{-1})$$

For head-on collision $p=0$, $f=1$ and

$$\tau_L^{-1} = 2 (\tau_H^{-1} + \tau_V^{-1})$$