

BNL-101744-2014-TECH RHIC/AP/89;BNL-101744-2013-IR

Comments on the a1, b1 Correction System

G. Parzen

March 1990

Collider Accelerator Department Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No.DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

AD/RHIC/AP-89

RHIC PROJECT Brookhaven National Laboratory

) 1

.

Comments on the a_1 , b_1 Correction System

G. Parzen

March 1990

COMMENTS ON THE a_1, b_1 CORRECTION SYSTEM

G. Parzen Accelerator Development Department March 20, 1990

1. a_1 Correction System

ν -Shift Effect

Coupling introduces normal modes ν_1, ν_2 .

Before Correction

$$|\nu_1 - \nu_2|_{max} = 228 \times 10^{-3} \quad \beta^* = 2$$
$$|\nu_1 - \nu_2|_{max} = 80 \times 10^{-3} \quad \beta^* = 6$$

$$|\nu_1 - \nu_2|_{max} = 80 \times 10^{-3} \ \beta^* = 6$$

(G. Parzen, AD/RHIC-AP-81, 1989.)

Global Correction System

 a_1 correctors in insertions at Q2 and Q5. Two families for each β^* . Q2 and Q5 for $\beta^* = 6$. Q2 for $\beta^* = 2$.

<u>After Correction</u> (Residual $\Delta \nu$)

 $\Delta \nu_{11}$ Stopband = 0 (Correct to make stopband = 0)

$$|\nu_1 - \nu_2|_{max} = 15 \times 10^{-3} \quad \beta^* = 6$$

 $|\nu_1 - \nu_2|_{max} = 20 \times 10^{-3} \quad \beta^* = 2$

<u>Empirical Correction</u> (Correct to minimize $|\nu_1 - \nu_2|$)

$$|\nu_1 - \nu_2| = 7 \times 10^{-3} \ \beta^* = 6$$

 $|\nu_1 - \nu_2| = 12 \times 10^{-3} \ \beta^* = 2$

Some additional a_1 correctors needed to reduce $|\nu_1 - \nu_2|$.

Note the advantage of setting correctors empirically, rather than setting them to cancel the $\nu_x = \nu_y$ stopband.

<u>Betatron Distortion due to Random a_1 </u>

Indicated by large $\beta_1, \beta_2 - \Delta\beta/\beta \simeq 60\%$ found.

Betatron distortion measured by coupling distribution factor, CDF

$$CDF = \frac{X_{max}(s)}{X_{max}(s)_{a_1=0}} \text{ for given } x_o, x'_o, y_o, y'_o$$

Usual assumption, $CDF \simeq 1.4$ for $\epsilon_{x,o} = \epsilon_{y,o}$. Computer study gives

$$(CDF)_{max} \simeq 2$$

Consequences of Large CDF – Aperture Loss

- a) <u>Linear Aperture Loss</u> Calculations of aperture required, e.g. the extraction magnet, can be off by 40%.
- b) <u>Dynamic Aperture Loss</u> $(A_{SL} \text{ loss})$

At some QF, A_{SL} can be reduced by 40%. Average loss A_{SL} about 15%. (G. Parzen, AD/RHIC-AP-80, 1989.)

<u>Residual</u> $|\nu_1 - \nu_2|$ and CDF Correction

<u>Proposal</u>: Separately excited a_1 near each high β quad in the insertions. Twelve separately excited a_1 near Q2.

This was also suggested by Correction System Review Committee.

This may correct both residual $|\nu_1 - \nu_2|$ and the CDF.

 a_1 correctors near QD in arc may also be helpful but may not be necessary.

The above 3 effects, (1) the residual $|\nu_1 - \nu_2|$, (2) the high CDF and (3) the loss in A_{SL} , deserve careful consideration before giving up the a_1 correctors in the arcs.

The a_1 at Q2 are excited in two families to generate the cos and sin of the $\nu_x + \nu_y$ harmonic. This gives a total of 4 knobs in the a_1 correction system to control $|\nu_1 - \nu_2|$ and the CDF.

Note that again the Q2 correctors are set empirically to reduce the CDF and the residual $|\nu_1 - \nu_2|$. They are not set to cancel out a $|\nu_x + \nu_y|$ stopband.

Are two knobs sufficient to control the CDF and the residual $|\nu_1 - \nu_2|$? Some judgment is needed as to what level of correction is sufficient.

 $\Delta \beta / \beta$ Effects of Random b_1

$$\beta^* = 6$$
 $(\Delta \beta / \beta)_{max} = 0.36$
 $\beta^* = 2$ $(\Delta \beta / \beta)_{max} = 0.90$

 $(\Delta\beta/\beta)_{max} = 0.20$ comes from the arcs. Largest effect from Q1, Q2, Q3, $\beta^* = 2$. (G. Parzen, AD/RHIC-AP-71, 1988.)

Above is a large effect and needs correction. Most of this effect can be corrected using b_1 corrections in the insertion quads.

<u>Proposal</u>: Use the b_1 corrections in the insertion quads to also correct the 20% effect due the arc magnets, as well as the large effect due to Q1, Q2, Q3.

Note that the a_1 effects and b_1 effects are similar and of the same order. The b_1 effects only appear smaller because of the availability of correction coils that are already there for other reasons. Both a_1 and b_1 produce a ν -shift, a betatron distortion, and a loss in dynamic aperture.

Using the b_1 coils in the insertions makes the b_1 correction system and the a_1 correction look similar.

<u>Proposal</u>: Use 4 knobs in the Q2, Q3 magnets to control the effects of the $2\nu_x$ and $2\nu_y$ resonances on $\Delta\beta_x/\beta_x$ and $\Delta\beta_y/\beta_y$. This will produce a certain level of correction which may be sufficient. More knobs could be added, but would be difficult to use.

<u>Why Correct b_1 ?</u>

 b_1 effects may be larger than expected because of

- a) difficulty in correcting Q1, Q2, Q3
- b) closed orbit errors in sextupoles
- c) other various sources
- d) operating near the integer ν -value.

3. Dispersion Correction

$$\frac{\Delta X_p}{X_p} = 0.25 \text{ at QF}$$
$$\frac{\Delta Y_{p,max}}{Y_p} = 0.31 \text{ at QD}$$

Mostly from arc magnets. (G. Parzen, AD/RHIC-AP-71, 1988.)

Reasons for Correcting X_p, Y_p

i

- 1) Effect on beam-beam interaction
- 2) To be able to operate near integer, $\nu \simeq 29$, where effect maybe about 3 times larger.

Proposal:

- 1) Use a_1 correctors at QD in arcs to correct Y_p . Possible a_1 corrector near Q9 as a back-up correction.
- 2) Use b_1 correctors at QF in arcs and b_1 in insertion quads to correct X_p .

The only new correctors required by the proposals in this note are the a_1 correctors at each Q2 in the insertions.