

Very High Energy Probes Of The Quark-Gluon Plasma

T. Ludlam

March 1984

Collider Accelerator Department
Brookhaven National Laboratory

U.S. Department of Energy

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

VERY HIGH ENERGY PROBES
OF THE
QUARK-GLUON PLASMA

T. Ludlam and F. Paige

Brookhaven National Laboratory

March 1984

VERY HIGH ENERGY PROBES OF THE QUARK-GLUON PLASMA

T. Ludlam and F. Paige

Brookhaven National Laboratory
Upton, New York 11973

L. Madansky
Johns Hopkins University
Baltimore, Maryland 21218

I. INTRODUCTION

Among the penetrating probes of nuclear matter the most frequently discussed have been those which involve the detection of photons or leptons with $m_T \approx P_T < 3$ GeV. This is the expected range of emission from a hot, thermalized plasma of quarks and gluons. The suggestion has been made that in very high energy collisions of nuclei the properties of high P_T jets may also reflect the characteristics of the nuclear medium through which the parent partons have propagated just after the collision.^{1,2} In this note we expand on the possible uses of such a probe.

II. HIGH P_T JETS

Consider a high energy nuclear collision in which a pair of partons suffers a hard scattering event of the sort which produces the hadronic jets observed at large transverse momentum in pp and pp collisions. One or both of the observed jets may be the hadronization of a quark or gluon which has propagated through a substantial volume of quark-gluon plasma produced in the collision. The average properties of such jets may be measurably different from those of "ordinary" jets. We have in mind two possible effects:

1) The hadronization of ordinary jets proceeds as $q\bar{q}$ pairs are materialized from the QCD vacuum (the "sea"). It may be that a quark or gluon propagating through the real sea of a deconfined plasma hadronizes with quite different properties which could signify the presence of a plasma and provide clues to its characteristics. The parameters of jets in which these differences might show up include the fragmentation function $f(Z)$, the transverse momenta of fragmentation products relative to the jet axis, and the flavor content of the leading particles. Specifically, if chiral symmetry is restored in the plasma we may expect an enhanced fraction of strangeness and heavier flavors among the fragmentation products of the jet. In addition, one should be sensitive to a possible halo of soft particles accompanying the jet, which may arise from "radiative" effects as the parent parton traverses the plasma. Such a component of soft particles in excess of the minimum bias background has been observed in multi-jet events seen in the UA-1 experiment at the CERN pp collider.³

2) The correlation within pairs of jets should be examined for evidence of the characteristic energy loss behavior suggested by Bjorken.¹ High energy quarks and gluons traversing the plasma may suffer energy loss comparable to the initial P_T , giving rise to a transverse momentum imbalance in the jet pair which is not

seen in pp collisions. In the extreme case a hard scatter occurring near the periphery of the plasma volume results in one jet escaping unattenuated while the other is completely absorbed.

To do these experiments we must detect jets and, in many instances, resolve individual particles within jets. From experience at the CERN collider, we expect that this will require $P_T(\text{jet}) > 20 \text{ GeV}$, and that these experiments will only be carried out in high energy colliding beam machines. Unlike most of the contemplated probes of a quark-gluon plasma, which involve the detection of soft particles amongst the very high multiplicity background, these measurements allow for very selective calorimetric triggers on local transverse energy deposition. Fig. 1 shows a possible detector arm covering 2 units of rapidity centered at $y = 0$, and 90° of azimuth. This detector arm was a design study⁴ for the detection of heavy quark jets at CBA. The detector is based on a well-segmented, high resolution calorimeter, and includes tracking chambers capable of resolving individual tracks within jets. The device was designed to have good efficiency for identifying both electrons (transition radiation detectors) and muons associated with jets. The correlation experiment would require two such arms. We have not examined in detail the performance of such a detector arm in a heavy ion collider, but we believe it would be well suited. Again, this is because the studies we are interested in here allow an effective anti-selection of the low P_T particles which give the most formidable backgrounds in these events.

To estimate the rate for observing jets we take the cross section for p-p collisions and assume this scales as A^2 . The $p+p \rightarrow \text{jet} + x$ cross section is given by ISAJET.⁵ The number of jets produced in a year of running is:

$$N = L T \sigma A^2$$

where σ is the cross section in pp collisions, and we take for the luminosity L and time T :

$$L = 10^{27} \text{ cm}^{-2} \text{ sec}^{-1}$$

$$T = 10^7 \text{ sec/machine year}$$

The number of jets per year for various P_T thresholds is shown in Fig. 2 as a function of the machine energy.

For heavy ions, the minimum P_T for observable jets is higher than in p-p collisions because of the many soft particles produced. For pp collisions at accelerator energies where jets are observed, the particle density (charged plus neutral) at mid-rapidity is:

$$\frac{dn}{dy} \approx 4.$$

The KNO multiplicity distribution gives:

$$\frac{(\langle n^2 \rangle - \langle n \rangle^2)^{1/2}}{\langle n \rangle} \approx \frac{1}{2}$$

With $\langle E_T \rangle \approx .4$ GeV/particle, we obtain:

$$\frac{dE_T}{dy} = (1.6 \pm 0.8) \text{ GeV},$$

where the fluctuation comes just from the multiplicity distribution.

Now consider A + A collisions. Treating these as A independent pp collisions we obtain:

$$\frac{dE_T^{(AA)}}{dy} = (1.6 A \pm .8 \sqrt{A}) \text{ GeV}.$$

For A = 200:

$$\frac{dE_T^{(A=200)}}{dy} = (320 \pm 11.5) \text{ GeV}.$$

Since the highest energy heavy ion collider presently under discussion has $\sqrt{s}/A \approx 100 + 100$ GeV/amu, no jet will ever dominate the total E_T distribution. In order to be observed, the P_T of the jet must be larger than the E_T in a cone large enough to contain it fully, say $\Delta y = 1$, $\Delta\phi = 45^\circ$. Then for A = 200:

$$P_T(\text{jet}) > \frac{dE_T}{dy} \Delta y \frac{\Delta\phi}{2\pi} = 40 \text{ GeV}/c.$$

Alternatively, we might ask that 2 P_T (jet) be larger than the fluctuations of E_T in the central $\Delta y = 2$ interval of rapidity. Hence for A = 200, requiring $> 4\sigma$:

$$P_T(\text{jet}) > \frac{1}{2} \left(\frac{dE_T}{dy} \right) \Delta y \times 4 = 45 \text{ GeV}/c.$$

These results imply that the observation of jets is possible in high energy collisions of heavy nuclei, but that for these experiments there is a premium on reaching the highest possible energy. Referring to Fig. 2, taking the curve for $P_T > 40$ GeV/c, we see that jets should be visible in usable numbers for a 100 + 100 GeV/amu collider.

III. DEEP INELASTIC SCATTERING VIA DRELL-YAN PAIRS

It may be that high P_T leptons produced in the massive Drell-Yan continuum can be exploited as a probe of the nuclear medium. Such a probe would be of interest if there were reason to expect observable radiative effects associated with the propagation of these leptons through the plasma.

A fundamental difference may be the effectively short multiple scattering path compared to the coherence length of the radiative effect (see paper on so-called Landau-Pomeranchuk-Migdal effect).⁶ Just how this phenomenon scales in a quark-gluon plasma would be an interesting study. If, however, such an effect is not too small, then one might look for photon pairs or jets that accompany the leptons.


This work was supported by the U.S. Department of Energy under contract DE-AC02-76CH00016 and DE-AC02-76ER03274.

REFERENCES

1. J.D. Bjorken, Fermilab Pub. 82/59/59-THY (1982).
2. L. Mc Lerran, BNL Report (1983).
3. G. Pancheri and C. Rubbia, Proc. III International Conference on Ultra Relativistic Nucleus-Nucleus Collisions, Nucl. Phys. A418, 117c (1984).
4. G. Benenson, et al., Proc. 1983 DPF Workshop on Collider Detectors, LBL-15973 (1983).
5. F. Paige and S. Protopopescu, BNL-29777; ISAJET Version 4.02 is used, with Eichten, Hinchcliffe, Lane, Quiqq (1984) Structure Functions.
6. T. Stanev, Ch. Vankov, R.E. Streitmatter, R.W. Ellsworth, T. Bowen, Phys. Rev. D25, 1291 (1982).

FIGURE CAPTIONS

- Fig. 1. Single arm heavy quark detector at $\theta = 90^\circ$, covering ± 1 units of rapidity and $\pm 45^\circ$ in azimuth (Ref. 4). The calorimeter is envisioned to be segmented in tower geometry, with ≈ 100 cells in the hadronic part and ≈ 1000 cells in the EM portion (the first 20 radiation lengths). TRD refers to transition radiation detectors.
- Fig. 2. The rates for two-jet events with the indicated P_T thresholds, as a function of collider energy. The calculation assumes a luminosity of $10^{27} \text{ cm}^{-2} \text{ sec}^{-1}$ integrated for 10^7 seconds.



1 METER

Fig. 1




Fig. 2