

# The Partial Beam Lifetime at RHIC due to Coulomb Dissociation of the Nucleus

M. J. Rhoades-Brown

July 1992

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

BNL-47806  
AD/RHIC-110  
Informal Report

**The Partial Beam Lifetime at RHIC due to Coulomb  
Dissociation of the Nucleus**

M. J. Rhoades-Brown and J. Weneser

July 1992

**R H I C   P R O J E C T**

Brookhaven National Laboratory  
Associated Universities, Inc.  
Upton, NY 11973

Under Contract No. DE-AC02-76CH00016 with the  
UNITED STATES DEPARTMENT OF ENERGY

The partial beam lifetime at RHIC due to Coulomb dissociation of the nucleus

M.J. Rhoades-Brown and J. Weneser\*

RHIC Project and Physics Department\*

Brookhaven National Laboratory

## Introduction

During beam crossing at RHIC, the Lorentz contracted Coulomb interaction between the heavy ions will excite internal modes of the nucleus. The subsequent decay of these modes is predominately via single or multiple nucleon emission. Changing the atomic mass of the beam ion will eventually cause beam intensity loss at RHIC for the radius of the ion orbit is sensitive to changes of the ratio  $Z/A$ .

While calculations for this beam loss mechanism have been made<sup>1</sup>, it is now clear that these earlier theoretical studies underestimated the Coulomb dissociation loss rate for they appear to have included only a limited range (first dipole resonance region, up to  $\sim 30$  MeV) of internal nuclear excitation energy.

In this report we reexamine the question of Coulomb dissociation cross sections at RHIC by including internal excitation energies up to thousands of GeV. In addition, we utilize experimental photonuclear absorption cross sections when evaluating the dissociation cross section. Also, internal excitation of a nucleus in one beam will result in both energy loss and transverse momentum change of an ion in the colliding beam. These recoil effects will be examined in detail to determine if there is an additional loss rate for ions out of the rf bucket or a non-negligible change in the ion's betatron momentum.

## Coulomb Dissociation Cross Section

The Coulomb dissociation cross section is given by the Weizsacker-Williams expression (whose accuracy is explored in Ref. 2)

$$\sigma = \frac{2\alpha Z_p^2}{\pi (\hbar c)^2 \gamma_{eff}^2} \int d\omega \omega \sigma_{ph}(\omega) \int_{b_{min}}^{\infty} b db K_1^2 \left( \left| \frac{b\omega}{\hbar c \gamma_{eff}} \right| \right) \quad (1)$$

where  $Z_p$  is the ion atomic number,  $\gamma_{eff}$  is the *fixed target frame* Lorentz parameter ( $\gamma_{eff} = 2\gamma^2 - 1$ ,  $\gamma = 108$  for  $^{197}\text{Au}^{79+}$  at RHIC),  $\alpha$  is the fine structure constant,  $K_1$  is a Macdonald or modified Bessel function,  $b$  is the impact parameter between the two

colliding ions, and  $\sigma_{ph}(\omega)$  is the target ion photonuclear cross section at photon energy  $\omega$  MeV. In this report the values of  $\sigma_{ph}(\omega)$  are taken from experimental data.

Equation (1) is an expression for the perturbative transition amplitude squared that is correct<sup>2</sup> to order  $1/\gamma_{\text{eff}}^2$ . The cut off in impact parameter is determined by the asymptotic behavior of  $K_1(z)$ . For  $z \gg 1$ ,  $K_1(z)$  drops faster then exponentially [as  $(\pi/2z)^{1/2}e^{-z}$ ]. The integration over  $b$  was carried out analytically using the expression,

$$\int^z t K_1(kt) K_1(kt) dt = (z^2/2) [K_1^2(kz) - K_0(kz) K_2(kz)] \quad . \quad (2)$$

In all results presented here, the lower integration limit  $b_{\text{min}}$  was taken to be 15fm, i.e. the grazing impact parameter between two Au ions. The dependence on  $b_{\text{min}}$  is weak.

The experimental values for  $\sigma_{ph}(\omega)$  were taken from several sources. Starting with the nuclear separation energy of 8 MeV,  $\sigma_{ph}(\omega)$  for Au ions and the range 8 MeV  $\leq \omega \leq 28$  MeV were taken from the measurements of Veyssiere<sup>3</sup> et al. For the range 28 MeV  $\leq \omega \leq 103$  MeV, the work of Lepretre<sup>4</sup> et al. was used. This data is for <sup>208</sup>Pb, and  $\sigma_{ph}$  was scaled as 197/208 to obtain Au values. For the range 103 MeV  $\leq \omega \leq 440$  MeV the scaled values (Pb data) of Carlos<sup>5</sup> et al. were used. For  $\omega$  values in the range 440 MeV  $\leq \omega \leq 2$  GeV there are no experimental measurements of  $\sigma_{ph}$  for Au. In this region scaled values of the  $\gamma$ , p and  $\gamma$ , n experimental cross section<sup>6</sup> were utilized, i.e.,  $\sigma_{ph}(\omega) = Z_T \sigma_{\gamma,p}(\omega) + (A_T - Z_T) \sigma_{\gamma,n}(\omega)$ . For the range 2 GeV  $\leq \omega \leq 9.51$  GeV the Au data of Michalowski<sup>7</sup> et al. was used. For the range 9.51 GeV  $< \omega < 17.84$  GeV the scaled values of the  $\gamma$ ,p and  $\gamma$ , n experimental cross sections<sup>6</sup> were once again utilized. Beyond the range of the table<sup>6</sup> ( $\omega > 17.84$  GeV) it was assumed that the  $\gamma$ , p and  $\gamma$ , n cross sections are constant in  $\omega$ , with a value taken from the tables at 17.84 GeV. Inspection of the tabulated data between 10 and 17 GeV indicates that this is a very reasonable assumption.

We note that using scaled  $\gamma$ ,p and  $\gamma$ ,n data for the range 440 MeV  $\leq \omega \leq 2$  GeV underestimates the  $\sigma_{ph}(\omega)$  cross section for Au. Comparing with known Au measurements at 440 MeV and 2 GeV suggests an underestimate in the contributions from this  $\omega$  region of approximately 15%. By contrast, studies by Michalowski et al. indicate that for  $\omega$  values greater than 9.51 GeV it is necessary to multiply  $\sigma$  by a shadowing factor<sup>7</sup>  $S$ , i.e. a factor to take into account the fact that not all nucleons contribute to the photonuclear

cross section. For Au ions with  $\omega > 9.51$  GeV we have taken a value<sup>7</sup> for S of 0.75. We also note, that except for the regions  $8 \text{ MeV} \leq \omega \leq 12 \text{ MeV}$  and  $28 \text{ MeV} \leq \omega \leq 103 \text{ MeV}$ , the photonuclear cross sections were given as tabulated experimental data. When actual tabulated data was unavailable, it was necessary to read published graphs.

In Table I, the relative contributions to  $\sigma$  are given for different ranges of  $\omega$ .

The error estimates for  $\sigma(b)$  were calculated from equation (1) using the experimental errors for the photonuclear cross sections. For the  $\omega$  range  $\omega > 9.51 \text{ GeV}$ , the overall uncertainties in the extrapolation suggested an error value of 1.0 b.

**Table 1:** Relative Contributions to  $\sigma$

$\omega$ Range (MeV)				$\sigma(b)$
8	$\leq$	$\omega$	$\leq$ 28	$61.5 \pm 6.1$
28	$\leq$	$\omega$	$\leq$ 103	$4.9 \pm 0.8$
103	$\leq$	$\omega$	$\leq$ 440	$12.4 \pm 0.6$
440	$\leq$	$\omega$	$\leq$ 2000	$8.5 \pm 1.0$
2000	$\leq$	$\omega$	$\leq$ 9510	$2.7 \pm 0.08$
9510	$\leq$	$\omega$	$\leq$ 25000	$2.3 \pm 1.0$

It can be guessed from this table where the quoted value<sup>1</sup> of 60b came from. Including higher excitation energies gives a total value of  $\sigma = (92.3 \pm 6.3)b$ , a  $\sim 54\%$  increase. The total error was calculated from a sum of squares analysis of the errors in Table 1.

#### Momentum Recoil out of the rf bucket

Including higher internal excitation energies in one nucleus, it is necessary to consider possible beam loss due to momentum recoil of the other colliding ion out of the rf bucket.

We label the two ions as 1 and 2, and use X to denote the laboratory or colliding frame. If  $\Delta E$  represents the energy loss of ion 2 in frame X,

$$\begin{aligned} E_2 &\rightarrow E_2 + \Delta E \\ p_2 &\rightarrow p_2 + \frac{\partial p}{\partial E} \Delta E \end{aligned} \tag{3}$$

Hence in the frame of ion 1 (denoted prime),

$$\Delta E'_2 + E'_2 = \gamma \left[ \left( E_2 + \Delta E - \beta \left( -p_2 - \frac{\partial p}{\partial E} \Delta E \right) \right) \right] \tag{4}$$

or

$$\Delta E'_2 = \gamma \left[ \Delta E + \frac{\partial p}{\partial E} \beta \Delta E \right] \quad (5)$$

$$= 2\gamma \Delta E \quad (6)$$

where the relation  $\beta = p/E$  (momentum of ion 1,  $p'_1 \equiv 0$ ) has been used.

Hence, if we assume an rf voltage of 6.7 MV, this gives a bucket area<sup>8</sup> of 1.31 eV sec/u and a  $\Delta E/E$  value for the bucket of  $\Delta E/E = 2 \times 10^{-3}$ . Hence for 100 GeV/u  $^{197}\text{Au}^{79}$  beams at RHIC, equation (6) gives

$$\Delta E'_2 = 851 \text{ GeV} \quad , \quad (7)$$

as the value of energy transfer beyond which the recoiling ion will be lost.

It is obvious from both the tabulated photonuclear cross sections<sup>6</sup>, and the cut-off properties of  $K_1(z)$  discussed earlier, that partial cross section contributions for  $\omega \geq 851$  GeV will be negligible. Recoil of an ion out of the rf bucket following Coulomb excitation of the colliding ion will not occur appreciably.

#### Momentum Recoil Effects on Betatron Motion

During beam crossing at RHIC, the r.m.s. value for the transverse component of the betatron momentum  $\langle p_\perp \rangle_{rms}$  is related to the longitudinal component of momenta  $p_\parallel$  via,

$$\langle p_\perp \rangle_{rms} = \frac{1}{\sqrt{2}} \sqrt{\frac{E_N}{\pi (\beta\gamma) \beta^*}} p_\parallel \quad , \quad (8)$$

where  $E_N$  is the normalized transverse emittance (taken to be  $10\pi \times 10^{-6}$  m rad) and  $\beta^*$  is the value of the beta function at the beam crossing point (taken to be 2m). With these values,

$$\langle p_\perp \rangle_{rms} = 1.5 \times 10^{-4} p_\parallel \equiv 3000 \text{ MeV} \quad , \quad (9)$$

where for Au ions  $p_\parallel = 100 \text{ GeV A/c}$ .

For Coulomb excitation of the ions, the energy transfer between the ions corresponds almost entirely to a longitudinal momentum transfer. The transverse momentum (as given by perturbation theory in the appendix) is of the order  $\omega/\gamma_{eff}$ . Since  $\omega \leq 25 \text{ GeV}$  and

$\gamma \sim 2.3 \times 10^4$ , the accompanying transverse momentum push is of the order  $10^{-3}$  GeV or less. This value is well within the betatron momentum aperture and hence we can assume the recoil accompanying Coulomb excitation to have a negligible effect on the betatron motion of the ions.

#### Partial Lifetime at RHIC

Using the values of luminosity in the Design Manual,<sup>1,9</sup> i.e.,  $L = 8.40 \times 10^{26} \text{ cm}^{-2} \text{ sec}^{-1}$ , the intensity decay rate  $\lambda_3$  for Coulomb dissociation of Au beams is now modified from  $19 \times 10^{-3} \text{ h}^{-1}$  to a value

$$\lambda_3 = (29.2 \pm 2.0) \times 10^{-3} \text{ h}^{-1} \quad (10)$$



### References

1. Conceptual Design of the Relativistic Heavy Ion Collider RHIC, p. 119, May 1989, BNL 52195.
2. A.J. Baltz, M.J. Rhoades-Brown, J. Weneser, Phys. Rev. A44 5569 (1991).
3. A. Veyssiere, H. Beil, R. Bergere, P. Carlos and L. Lepretre, Nucl. Phys. A159, 561 (1970).
4. A. Lepretre, H. Beil, R. Gergere, P. Carlos, J. Fagot, A. De Miniac and A. Veyssiere, Nucl. Phys. A367, 237 (1981).
5. P. Carlos, H. Beil, R. Bergere, J. Fagot, A. Lepretre, A. De Miniac and A. Veyssiere, Nucl Phys. A431, 573 (1984).
6. Total cross-sections for reactions of high energy particles, Volume 12, Landolt- Bornsein, edited by H. Schopper.
7. S. Michalowski, D. Andrews, J. Eickmeyer, T. Gentile, N. Mistry, R. Talman, K. Ueno, Phys. Rev. Lett. 39, 737 (1977).
8. H. Hahn, private communication.
9. In Ref. (1), p. 117–119, it is stated that a luminosity value for Au ions from Table IV. 3–1 was used in estimating the beam loss rate. Two values are given in Table IV. 3–1,  $L = 2 \times 10^{26} \text{cm}^{-2} \text{sec}^{-1}$  and  $L = 1.1 \times 10^{27} \text{cm}^{-2} \text{sec}^{-1}$ . However, closer inspection of the loss rates quoted indicates a luminosity value of  $L = 8.40 \times 10^{26} \text{cm}^{-2} \text{sec}^{-1}$  was used. Of course, a self-consistent analysis of time dependent luminosity and beam loss rate that includes intra-beam scattering is required.

## APPENDIX

In order to calculate the transverse recoil momentum given to the projectile ion it is useful to re-work the Weizsacker–Williams cross section so as to make explicit the momentum transfers. Distinguishing between projectile ion and target ion by the subscripts  $P, T$ , we can write the **transverse**–photon transfer cross section:

$$\frac{d\sigma}{d\Omega} = 2\pi \int dE_F \frac{p_{P_f} E_{P_f}}{(2\pi\hbar)^3 v_{P_0}} \left| \langle p_{P_f} | \vec{j} \cdot \hat{e} e^{-i\vec{q} \cdot \vec{r}_P} | p_{P_0} \rangle \right|^2 \frac{(4\pi)^2}{[\vec{q}^2 - q_0^2]^2} \left| \langle T_f | \vec{j} \cdot \hat{e} e^{i\vec{q} \cdot \vec{r}_T} | T_0 \rangle \right|^2$$

The  $\vec{q}$  represents the momentum transfer,  $\vec{q} = \vec{p}_{P_0} - \vec{p}_{P_f}$ ;  $q_0$  is the corresponding energy transfer,  $(E_{P_0} - E_{P_f}) = \omega$ . It is assumed here that the projectile ion is **not** internally excited, and that it is sufficient to represent the initial and final projectile states as plane waves. The target states need not be explicitly represented since the target matrix element will be swept into the effective photon cross section as in the usual Weizsacker–Williams approximations. It can be shown (not here) that the Coulomb or longitudinal photon contribution is of lower order in  $M_P/E_P = 1/\gamma$ , and that the transverse part outlined above suffices for the leading  $(\ln \gamma)$  approximation.

The matrix element for the projectile transition, taken as that of a spinless particle, leads to the vertex strength

$$\begin{aligned} \sum_{\hat{e}} \left| \langle p_{P_f} | \vec{j} \cdot \hat{e} e^{-i\vec{q} \cdot \vec{r}_P} | p_{P_0} \rangle \right|^2 &= \alpha Z_P^2 \sum \left| \frac{\overrightarrow{p_{P_f} + p_{P_0}} \cdot \hat{e}}{2\sqrt{E_{P_f} \cdot E_{P_0}}} \right|^2 \\ &= \frac{\left( \overrightarrow{p_{P_f} + p_{P_0}} \right)^2 - \left( \left( \overrightarrow{p_{P_f} + p_{P_0}} \right) \cdot \vec{q} \right)^2 / q^2}{4E_{P_0}E_{P_f}} \\ &= \frac{2 \left( p_{P_0} + p_{P_f} \right)^2}{E_{P_0}E_{P_f} (\vec{q})^2} (\chi - \chi^2/2), \\ \chi &= 1 - \cos \theta = \theta^2/2 + \dots; \end{aligned}$$

$\theta$  is the projectile scattering angle. For high energies and relatively small energy losses this is

$$\cong \frac{2p_{P_0}}{q^2} (\chi - \chi^2/2).$$

Similarly

$$\begin{aligned}
\vec{q}^2 &= \left( \overrightarrow{p_{P_0} - p_{P_f}} \right)^2 = \left( p_{P_0} - p_{P_f} \right)^2 + 2p_{P_0}p_{P_f}\chi \\
&= 2p_{P_0}p_{P_0}\chi + \omega^2 + \omega^2 \frac{M_P^2}{p_{P_0}^2} + \left( \text{higher order in } 1/p_{P_0} \right) \\
\vec{q}^2 - q_0^2 &= 2p_{P_0}p_{P_f}\chi + \omega^2 \frac{M_P^2}{p_{P_0}^2} + \left( \text{higher order in } 1/p_{P_0} \right) \\
&\cong 2p_{P_0}p_{P_f}\chi + \omega^2/\gamma^2 \\
\vec{q}^2 &\cong 2p_{P_0}p_{P_f}\chi + \omega^2, \quad \gamma \gg 1.
\end{aligned}$$

This allows us to write  $d\sigma/d\Omega$  as:

$$\frac{d\sigma}{d\Omega} = 8\alpha Z_p^2 \int dE_{P_f} \frac{p_{P_f} E_{P_f}}{v_{p_0}} \frac{p_{P_0}^2 (\chi - \chi^2/2)}{\left[ \omega^2 + 2p_{P_0}p_{P_f}\chi \right] \left[ \omega^2/\gamma^2 + 2p_{P_0}p_{P_f}\chi \right]^2} \left| \langle T_f | j_\perp e^{i\vec{q} \cdot \vec{r}_T} | T_0 \rangle \right|^2$$

The equivalent of the Weizsacker–Williams approximation is made by noting that the denominator factor,  $[\omega^2/\gamma^2 + 2p_{P_0}p_{P_f}\chi]^2$ , weights  $\chi$  so that  $(2p_{P_0}p_{P_f}\chi)$  is of order  $\omega^2/\gamma^2$ . Then,  $\vec{q}^2$  differs from  $\omega^2$  only by the negligible quantity  $\omega^2/\gamma^2$ . This, in turn, means that, to this order, the intermediate virtual photon is on the mass shell; that is, since  $|\vec{q}| \simeq \omega$ , the matrix element  $\langle T_f | j_\perp e^{i\vec{q} \cdot \vec{r}_T} | T_0 \rangle$  is well approximated by the real photon matrix element and we can take over from experiment the photo cross section:

$$\sigma_\gamma(T_0 \rightarrow T_F) = \frac{4\pi^2}{\omega} \left| \langle T_f | j_\perp e^{i\vec{q} \cdot \vec{r}_T} | T_0 \rangle \right|^2.$$

Then

$$\frac{d\sigma}{d\Omega} = 8\alpha Z_p^2 \int dE_{P_f} \frac{p_{P_f} E_{P_f}}{v_{p_0}} \frac{p_{P_0}^2 (\chi - \chi^2/2)}{\left[ \omega^2 + 2p_{P_0}p_{P_f}\chi \right] \left[ \omega^2/\gamma^2 + 2p_{P_0}p_{P_f}\chi \right]^2} \frac{\omega}{4\pi^2} \sigma_\gamma(T_0 \rightarrow T_F).$$

Then, with the necessary neglect of  $\chi^2/2$ ,  $p_{P_f} \simeq E_{P_f} \simeq p_{P_0}$ ,  $v_{p_0} \sim 1$ :

$$\begin{aligned}
\sigma &= \frac{4}{\pi} p_{P_0}^4 \alpha Z_p^2 \int dE_{P_f} \omega \int d\chi \frac{\chi}{\left[ \omega^2 + 2p_{P_0}^2 \chi \right] \left[ \omega^2/\gamma^2 + 2p_{P_0}^2 \chi \right]^2} \sigma_\gamma(T_0 \rightarrow T_F) \\
&= \frac{2}{\pi} \int d\omega \frac{1}{\omega} \left[ \ln \gamma - 1/2 + \theta \left( \frac{\omega^2}{p_{P_0}^2} \ln \frac{p_{P_0}}{\omega} \right) \right] \sigma_\gamma(T_0 \rightarrow T_F),
\end{aligned}$$

displaying the leading  $\ln \gamma$  of the Weizsacker–Williams form. We do not attempt to recover the subsequent terms.

It is necessary to note the limitations inherent in the above derivation. In order to take over the photo cross section, the closeness of  $|\vec{q}|$  to  $\omega$  is vital. This is, in fact, correct for the part that leads to the  $\ln \gamma$  dependence. It is, however, less true for the higher orders. In fact, for the  $\chi^2$  term it is not at all valid, since if  $\chi$  is replaced by  $\chi^2$  the weighting is no longer such that  $2p_{P_0}p_{P_f}\chi \sim \mathcal{O}(\omega^2/\gamma^2)$ , but is rather spread over  $\mathcal{O}(\omega^2)$ ; then  $\vec{q}^2$  is no longer just  $\omega^2$ , but  $\omega^2 + \mathcal{O}(\omega^2)$ . This does **not** ruin the leading  $(\ln \gamma)$  approximation since it can easily be seen that the involving term  $\chi^2$  is small, of order  $\omega^2/p_{P_0}^2 (\ln(p_{P_0}/\omega)/\ln \gamma)$  relative to the leading term, if one makes the reasonable assumption that the matrix element,  $\langle T_F | j_\perp e^{i\vec{q}\cdot\vec{r}} | T_0 \rangle$ , is of the same order for the larger range of  $|\vec{q}|$ . If, however, we wish to calculate some average value of momentum transfer, as say

$$p_{P_0} \sin \theta \simeq \sqrt{2\chi} p_{P_0},$$

the extra weighting involved in the calculation of

$$\int d\chi \frac{d\sigma}{d\Omega} \sqrt{\chi} / \int d\chi \frac{d\sigma}{d\Omega},$$

moves  $|\vec{q}|$  into the larger domain. Therefore, without bringing in the full form factor of the photo reaction no close approximation is possible. It can, however, provide an order of magnitude estimate. For our purposes here, we need not go into it.

It is sufficient to note that the leading order of the excitation cross section is contained within the domain roughly specified by

$$2p_{P_0}p_{P_f}\chi \sim \mathcal{O}(\omega^2/\gamma^2).$$

Since the transverse component is defined by

$$p_{P_\perp} = p_P \sin \theta \sim p_P \theta,$$

we have that the effective range of transverse momenta is given as:

$$p_{P_\perp} = p_P \sin \theta \sim \mathcal{O}(\omega/\gamma).$$

the order of magnitude used in the text.