

## AGS Modifications for High Transition Energy

E. D. Courant

May 1996

Collider Accelerator Department  
**Brookhaven National Laboratory**

**U.S. Department of Energy**

USDOE Office of Science (SC)

Notice: This technical note has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. The publisher by accepting the technical note for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this technical note, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

# AGS MODIFICATIONS FOR HIGH TRANSITION ENERGY

E. D. Courant and D. Trbojevic

May 14, 1996

## 1 Introduction

The transition energy in the AGS is at approximately 8 GeV ( $\gamma \cong 8.5$ ), right in the middle of the acceleration range. This inevitably causes some beam loss, especially at high intensity. If the transition energy could be moved outside the acceleration range these problems could be avoided; this would be advantageous for high-intensity operation, and especially for operation with heavy ions.

The transition energy  $\gamma_t$  is related to the momentum compaction as:  $\alpha_0 = 1/\gamma_t^2$ , while the momentum compaction defined to the lowest order in  $\delta$  is related to the horizontal dispersion function  $D_x$  by:

$$\alpha_0 = \frac{1}{C_0} \oint \frac{D_x(s)}{\rho(s)} ds, \quad (1.1)$$

where  $\rho$  is the radius of curvature and  $s$  is the longitudinal path length measured along the reference orbit with a circumference  $C_0$ .

## 2 An Example of the AGS Lattice Without Transition

A value of the transition energy could be adjusted by additional quadrupoles which introduce oscillations of the dispersion within dipoles between positive and negative values. We have found a way of accomplishing this by inserting extra quadrupoles in some of the "5 foot" straight sections of the existing AGS lattice; however, several of the dipoles in each superperiod have to be shifted azimuthally. The drawbacks are that the new orbit is moved radially compared to the old one (but still fits well within the tunnel), and that the peak beta functions and dispersion function are considerably larger than in the standard AGS.

The modified lattice is obtained from the standard one by the following modifications (in each of the 12 superperiods):

Move magnets 15 and 16 by 3.7 mm away from the symmetry point, e.g. lengthen the straight section 15 by 7.4 mm to 1.5314 m. Move magnets 14 and 17 by 41.84 cm away from the symmetry point:

Straight sections 14 and 16 lengthened from 0.6096 to 1.0243 m.

Table I: New and Old AGS Betatron Lattice Functions

	Original	Modified
Transition $\gamma$	8.427	30.59
H. Tune $\nu_x$	8.687	11.232
V. Tune $\nu_y$	8.780	7.436
max $\beta_x$	22.33	22.53
min $\beta_x$	10.23	3.19
max $\beta_y$	22.33	32.64
min $\beta_y$	10.13	10.13
max $\eta$	2.17	3.88
min $\eta$	1.48	-3.90

Move magnets 13 and 18 by 14.74 cm toward symmetry point:


Straight sections 13 and 17 shortened from 1.524 to 0.9582 m.

Insert quadrupole (H-focusing) in center of straight sections 13 and 17; length 0.2 m, gradient 25.905 T/m at peak energy ( $B\rho = 98.144$  T-m).

Insert trim quadrupole (0.2m, 60 Gauss/m) in straight sections 3 and 7. (This one is so weak that it should be possible to find a solution where it is absent). With these modifications the lattice functions are presented in Table I as follows:

The lattice functions and layout are pictured for the original AGS and the modified lattice in Figs. 1 and 2; the figures show one superperiod starting with magnet 6 and ending with magnet 5 of the next superperiod.

The main drawback of this scheme is that it requires moving six magnets per superperiod (72 in all), which implies a major rebuilding of the AGS. And with the increase maximum dispersion function and vertical  $\beta$  function some aperture problems may become more severe. Furthermore some of the existing sextupoles might have to be moved.



Dispersion max/min: 2.16942/ 1.47640m,

$\gamma: ( 8.43, 0.00)$


$\beta_x$  max/min: 22.53/10.23229m,  $v_x: 8.68689$ ,  $\xi_x: -8.93$ ,

Module length: 807.1104m

$\beta_y$  max/min: 22.33/10.12588m,  $v_y: 8.78018$ ,  $\xi_y: -9.29$ ,

Total bend angle: 6.28319311 rad

**Figure 1: Orbit Functions in standard AGS Lattice**



Dispersion max/min: 3.88070/-3.90000m,

$\beta_x$  max/min: 22.53/3.18925m,  $\nu_x$ : 11.23200,  $\xi_x$ : -11.95,

$\beta_y$  max/min: 32.64/10.12588m,  $\nu_y$ : 7.43633,  $\xi_y$ : -10.32,

$\gamma_t$ : ( 30.59, 0.00)

Module length: 807.1104m

Total bend angle: 6.28319311 rad

**Figure 2: Orbit Functions in AGS lattice with shifted magnets and quads**